

## SENIOR SECONDARY INTERVENTION PROGRAMME 2013



**education**

Department: Education

**GAUTENG PROVINCE**



**GRADE 12**

**MATHEMATICS**

**TEACHER NOTES**

The SSIP is supported by



# TABLE OF CONTENTS

## TEACHER NOTES

SESSION	TOPIC	PAGE
1	Topic 1: Logarithms Topic 2: Factorisation of 3 <sup>rd</sup> degree polynomials	3 – 13
2	Sequences & Series	14 – 25
3	Financial Maths	26 – 31
4	Financial Maths	32 – 39
5	Trigonometry Revision	40 – 47
6	Trigonometry	48 – 52
7	Trigonometry	53 – 59
8	Consolidation: Topic 1: Number patterns, sequences and series, Topic 2: Financial mathematics & Trigonometry	60 – 65
9	Consolidation: Topic 1: Number patterns, sequences and series, Topic 2: Financial mathematics	66 – 75
10	Transformations	76 – 83
11	Functions and graphs	84 – 90
12	Inverse graphs	91 – 97
13	Calculus	98 – 111
14	Topic 1: Calculus Graphs Topic 2: Linear Programming	112 - 123

**TOPIC 1: LOGARITHMS**

**Teacher Note:** Changing from exponential to logarithmic form in real world problems is the most important concept in this section. This concept is particularly useful in Financial Maths when learners are required to solve for  $n$ .

**LESSON OVERVIEW**

- 1. Introduction session: 5 minutes
- 2. Typical exam questions:
  - Question 1: 10 minutes
  - Question 2: 5 minutes
  - Question 3: 10 minutes
- 3. Discussion of solutions: 30 minutes

**SECTION A: TYPICAL EXAM QUESTIONS**

**QUESTION 1** (Used with permission from Maths Grade 12 Allcopy Mind Action Series textbook)

A colony of an endangered species originally numbering 1000 was predicted to have a population  $N$  after  $t$  years given by the equation  $N = 1000(0,9)^t$ .

- (a) Estimate the population after 1 year. (2)
  - (b) Estimate the population after 2 years. (2)
  - (c) After how many years will the population decrease to 200? (5)
- [9]

**QUESTION 2** (DoE Nov 2008)

R1 570 is invested at 12% per annum. compound interest. After how many years will the investment be worth R23 000? [4]


**QUESTION 3** (Link to inverse graphs)

Given:  $g(x) = \left(\frac{1}{2}\right)^x$

- (a) Write the inverse of  $g$  in the form  $g^{-1}(x) = \dots$  (2)
  - (b) Sketch the graph of  $g^{-1}$  (2)
  - (c) Determine graphically the values of  $x$  for which  $\log_{\frac{1}{2}} x < 0$  (1)
- [5]

## SECTION B: SOLUTIONS AND HINTS TO SECTION A

1(a)	At $t = 0$ , $N = 50 \cdot 2^0 = 50$ "super-bugs"	<ul style="list-style-type: none"> <li>✓ <math>t = 0</math></li> <li>✓ <math>N = 50 \cdot 2^0 = 50</math></li> </ul> <p style="text-align: right;">(2)</p>
1(b)	At $t = 5$ , $N = 50 \cdot 2^5 = 1600$ "super-bugs"	<ul style="list-style-type: none"> <li>✓ <math>N = 50 \cdot 2^5</math></li> <li>✓ <math>= 1600</math></li> </ul> <p style="text-align: right;">(2)</p>
1(c)	Here $N = 10\,000$ $N = 50 \cdot 2^t$ $\therefore 10\,000 = 50 \cdot 2^t$ $\therefore 200 = 2^t$ $\therefore \log_2 200 = t$ $\therefore t = 7,64385619$ $\therefore t \approx 8$ days	<ul style="list-style-type: none"> <li>✓ <math>10\,000 = 50 \cdot 2^t</math></li> <li>✓ <math>200 = 2^t</math></li> <li>✓ <math>\log_2 200 = t</math></li> <li>✓ <math>t = 7,64385619</math></li> <li>✓ <math>t \approx 8</math> days</li> </ul> <p style="text-align: right;">(5)</p>
2	$A = P(1+i)^n$ $\therefore 23000 = 1570(1+0,12)^n$ $\therefore \frac{23000}{1570} = (1,12)^n$ $\therefore 14,64968153\dots\dots = (1,12)^n$ $\therefore \log_{1,12} 14,64968153\dots\dots = n$ $\therefore n = 23,68701789\dots$ years $\therefore n = 23$ years 8 months	<ul style="list-style-type: none"> <li>✓ formula</li> <li>✓ substitution</li> <li>✓ apply log function</li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p>The 8 months was calculated by multiplying the decimal 0,68701789 by 12</p> </div>
3(a)	$y = \left(\frac{1}{2}\right)^x$ $x = \left(\frac{1}{2}\right)^y$ $\therefore \log_{\frac{1}{2}} x = y$ $\therefore g^{-1}(x) = \log_{\frac{1}{2}} x$	<div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p>Remember that the inverse of a graph is determined by interchanging x and y in the equation of the original graph.</p> </div> <ul style="list-style-type: none"> <li>✓ <math>x = \left(\frac{1}{2}\right)^y</math></li> <li>✓ <math>g^{-1}(x) = \log_{\frac{1}{2}} x</math></li> </ul> <p style="text-align: right;">(2)</p>

3(b)	 <div style="border: 1px solid black; padding: 5px; margin: 10px auto; width: fit-content;">                 Don't forget to indicate the coordinates of the intercept with the axes.                  The y-axis is the asymptote of the graph.             </div>	✓ shape ✓ (1;0) (2)
3(c)	$\log_{\frac{1}{2}} x < 0$ for $x > 1$	✓ $x > 1$ (1)

**SECTION C: HOMEWORK**

**QUESTION 1** (Used with permission from Maths Grade 12 Mind Action Series textbook)  
 Archaeologists use a specific formula when determining the age of fossils. This formula is the carbon dating formula and is given by:  $P = \left(\frac{1}{2}\right)^{\frac{n}{5700}}$  where P is the percentage of carbon-14 remaining in the fossils after  $n$  years. Calculate the approximate age of a certain fossil discovered if the percentage of carbon-14 in the fossil is 12,5%. [4]

**QUESTION 2** (Used with permission from Maths Grade 12 Mind Action Series textbook)

(a) Determine how many years it would take for the value of a car to depreciate to 50% of its original value if the rate of depreciation, based on the reducing balance method, is 8% per annum. (3)

(b) How long will it take for an amount of R50 000 to double if the interest rate is 18% per annum compounded monthly? (6)  
 [9]

**QUESTION 3**

The graph of  $f : x \rightarrow \log_a x$  passes through the point (16 ; 2).


(a) Calculate the value of  $a$ . (3)

(b) Write down the equation of the inverse in the form  $f^{-1}(x) = \dots$  (2)

(c) Sketch the graphs of  $f$  and  $f^{-1}$  on the same set of axes. (4)  
 [9]

**SECTION D: SOLUTIONS TO HOMEWORK**

1	$0,125 = \left(\frac{1}{2}\right)^{\frac{n}{5700}}$ $\therefore 0,125 = (0,5)^{\frac{n}{5700}}$ $\therefore \log_{0,5} 0,125 = \frac{n}{5700}$ $\therefore 3 = \frac{n}{5700}$ $\therefore n = 17100 \text{ years}$	<p>✓ 0,125</p> <p>✓ <math>0,125 = (0,5)^{\frac{n}{5700}}</math></p> <p>✓ <math>\log_{0,5} 0,125 = \frac{n}{5700}</math></p> <p>✓ <math>n = 17100 \text{ years}</math></p> <p style="text-align: right;"><b>[4]</b></p>
2(a)	$0,5x = x(1 - 0,08)^n$ $\therefore 0,5 = 0,92^n$ $\therefore \log_{0,92} 0,5 = n$ $\therefore n = 8,312950414$ <p>8 years and 4 months</p>	<p>✓ correct substitution into formula</p> <p>✓ use of logs</p> <p>✓ answer</p> <p style="text-align: right;">(3)</p>
2(b)	$A = P \left(1 + \frac{i}{12}\right)^{12n}$ $100\ 000 = 50\ 000 \left(1 + \frac{0,18}{12}\right)^{12n}$ $\therefore 2 = (1,015)^{12n}$ $\therefore \log_{1,015} 2 = 12n$ $\therefore 12n = 46,55552563$ $\therefore n = 3,879627136$ <p>3 years 11 months</p> <div style="border: 1px solid black; padding: 5px; display: inline-block; margin-left: 100px;"> <p>Divide <math>i</math> by 12 and multiply <math>n</math> by 12</p> </div>	<p>✓ <math>12n</math></p> <p>✓ <math>\frac{0,18}{12}</math></p> <p>✓ <math>2 = (1,015)^{12n}</math></p> <p>✓ <math>\log_{1,015} 2 = 12n</math></p> <p>✓ <math>n = 3,879627136</math></p> <p>✓ 3 years 11 months</p> <p style="text-align: right;">(6) <b>[9]</b></p>
3(a)	$\therefore 2 = \log_a 16$ $\therefore a^2 = 16$ $\therefore a = 4$	<p>✓ <math>2 = \log_a 16</math></p> <p>✓ <math>a^2 = 16</math></p> <p>✓ <math>a = 4</math></p> <p style="text-align: right;">(3)</p>

<p>3(b)</p>	$y = \log_4 x$ $\therefore x = \log_4 y$ $\therefore 4^x = y$ $\therefore f^{-1}(x) = 4^x$	<p>✓ <math>x = \log_4 y</math></p> <p>✓ <math>f^{-1}(x) = 4^x</math></p> <p>(2)</p>
<p>3(c)</p>		<p>✓ Graph of <math>f(x)</math></p> <p>✓ y-intercept of <math>f</math></p> <p>✓ Graph of <math>f^{-1}(x)</math></p> <p>✓ x-intercept of <math>f^{-1}</math></p> <p>(4)</p> <p>[9]</p>



**TOPIC 2: FACTORISATION OF THIRD DEGREE POLYNOMIALS**

**Teacher Note:** The factorisation of third degree polynomials is essential when sketching the graphs of cubic functions. The  $x$ -intercepts of a cubic graph can be determined by factorising cubic polynomials.

**LESSON OVERVIEW**

1. Introduction session: 5 minutes
2. Typical exam questions:
  - Question 1: 20 minutes
  - Question 2: 10 minutes
3. Discussion of solutions: 25 minutes

**SECTION A: TYPICAL EXAM QUESTIONS****QUESTION 1**

Solve the following equations:



- (a)  $x^3 - x^2 - 22x + 40 = 0$  (4)
- (b)  $x^3 + 9x^2 + 26x + 24 = 0$  (4)
- (c)  $3x^3 - 7x^2 + 4 = 0$  (4)
- (d)  $4x^3 - 19x - 15 = 0$  (4)
- (e)  $x^3 - x^2 - 4x + 4 = 0$  (4)
- [20]

**QUESTION 2**

Solve the following equations:

- (a)  $x^3 - 27 = 0$  (4)
- (b)  $x^3 + 27 = 0$  (4)
- [8]



**SECTION B: SOLUTIONS AND HINTS TO SECTION A**

1(a)	$x^3 - x^2 - 22x + 40 = 0$ $\therefore (x-2)(x^2 + x - 20) = 0$ $\therefore (x-2)(x+5)(x-4) = 0$ $\therefore x = 2$ or $x = -5$ or $x = 4$	✓ $(x-2)$ ✓ $(x^2 + x - 20)$ ✓ $(x+5)(x-4)$ ✓ $x = 2$ or $x = -5$ or $x = 4$ (4)
1(b)	$x^3 + 9x^2 + 26x + 24 = 0$ $\therefore (x+3)(x^2 + 6x + 8) = 0$ $\therefore (x+3)(x+4)(x+2) = 0$ $\therefore x = -3$ or $x = -4$ or $x = -2$	✓ $(x+3)$ ✓ $(x^2 + 6x + 8)$ ✓ $(x+4)(x+2)$ ✓ $x = -3$ or $x = -4$ or $x = -2$ (4)
1(c)	$3x^3 - 7x^2 + 4 = 0$ $\therefore (x-1)(3x^2 - 4x - 4) = 0$ $\therefore (x-1)(3x+2)(x-2) = 0$ $\therefore x = 1$ or $x = -\frac{2}{3}$ or $x = 2$	✓ $(x-1)$ ✓ $(3x^2 - 4x - 4)$ ✓ $(3x+2)(x-2)$ ✓ $x = 1$ or $x = -\frac{2}{3}$ or $x = 2$ (4)
1(d)	$4x^3 - 19x - 15 = 0$ $\therefore (x+1)(4x^2 - 4x - 15) = 0$ $\therefore (x+1)(2x-5)(2x+3) = 0$ $\therefore x = -1$ or $x = \frac{5}{2}$ or $x = -\frac{3}{2}$	✓ $(x+1)$ ✓ $(4x^2 - 4x - 15)$ ✓ $(2x-5)(2x+3)$ ✓ $x = -1$ or $x = \frac{5}{2}$ or $x = -\frac{3}{2}$ (4)
1(e)	$x^3 - x^2 - 4x + 4 = 0$ $\therefore x^2(x-1) - 4(x-1) = 0$ $\therefore (x-1)(x^2 - 4) = 0$ $\therefore (x-1)(x+2)(x-2) = 0$ $\therefore x = 1$ or $x = -2$ or $x = 2$	✓ $(x-1)$ ✓ $(x^2 - 4)$ ✓ $(x+2)(x-2)$ ✓ $x = 1$ or $x = -2$ or $x = 2$ (4)

[20]

## MATHEMATICS

## GRADE 12

## SESSION 1

## (TEACHER NOTES)

<p>2(a)</p>	$x^3 - 27 = 0$ $\therefore (x-3)(x^2 + 3x + 9) = 0$ $\therefore x = 3 \quad \text{or} \quad x^2 + 3x + 9 = 0$ $x = \frac{-3 \pm \sqrt{(3)^2 - 4(1)(9)}}{2(1)}$ $x = \frac{-3 \pm \sqrt{-27}}{2}$ <p>non-real solution</p> <p>OR</p> $x^3 - 27 = 0$ $\therefore x^3 = 27$ $\therefore x = 3$	$\checkmark (x-3)(x^2 + 3x + 9) = 0$ $\checkmark x = 3$ $\checkmark x = \frac{-3 \pm \sqrt{-27}}{2}$ $\checkmark x = 3 \text{ is the only real solution} \quad (4)$ $\checkmark\checkmark x^3 = 27$ $\checkmark\checkmark x = 3$
<p>2(b)</p>	$x^3 + 27 = 0$ $\therefore (x+3)(x^2 - 3x + 9) = 0$ $\therefore x = -3 \quad \text{or} \quad x^2 - 3x + 9 = 0$ $x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(9)}}{2(1)}$ $x = \frac{3 \pm \sqrt{-27}}{2}$ <p>non-real solution</p> <p>OR</p> $x^3 + 27 = 0$ $\therefore x^3 = -27$ $\therefore x = -3$	$\checkmark (x+3)(x^2 - 3x + 9) = 0$ $\checkmark x = -3$ $\checkmark x = \frac{3 \pm \sqrt{-27}}{2}$ $\checkmark x = -3 \text{ is the only real solution} \quad (4)$ $\checkmark\checkmark x^3 = -27$ $\checkmark\checkmark x = -3$

[8]

**SECTION C: HOMEWORK**

**QUESTION 1**

Solve the following equations, rounded off to two decimal places where appropriate:

(a)  $x^3 - 6x^2 - x - 6 = 0$  (4)

(b)  $x^3 - 2x^2 + 16 = 0$  (5)

(c)  $2x^3 - 5x^2 - 4x + 3 = 0$  (4)

(d)  $-x^3 + 4x^2 - 2x - 4 = 0$  (6)

(e)  $x^3 - 5x^2 - 3x + 9 = 0$  (4)

[23]

**QUESTION 2**

(a) Determine the coordinates of the  $x$ -intercepts of the graph of  $f(x) = x^3 - 8x^2 + 19x - 12$  (6)

(b) Show that the graph of  $f(x) = x^3 - x^2 - x - 2$  cuts the  $x$ -axis at one point only. (5)

[11]

**SECTION D: SOLUTIONS TO HOMEWORK**

1(a)	$x^3 - 6x^2 - x - 6 = 0$ $\therefore (x-1)(x^2 - 5x - 6) = 0$ $\therefore (x-1)(x-6)(x+1) = 0$ $\therefore x = \pm 1 \text{ or } x = 6$	$\checkmark (x-1)$ $\checkmark (x^2 - 5x - 6)$ $\checkmark (x-6)(x+1)$ $\checkmark x = \pm 1 \text{ or } x = 6$	(4)
------	---	---	-----

## MATHEMATICS

## GRADE 12

## SESSION 1

## (TEACHER NOTES)

1(b)	$x^3 - 2x^2 + 16 = 0$ $\therefore (x+2)(x^2 - 4x + 8) = 0$ $\therefore x = -2 \text{ or } x = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(8)}}{2(1)}$ $x = \frac{4 \pm \sqrt{-16}}{2}$ <p style="text-align: center;">non-real</p>	<ul style="list-style-type: none"> <li>✓ <math>(x+2)</math></li> <li>✓ <math>(x^2 - 4x + 8)</math></li> <li>✓ <math>x = -2</math></li> <li>✓ <math>x = \frac{4 \pm \sqrt{-16}}{2}</math></li> <li>✓ <math>x = -2</math> is the only real solution (5)</li> </ul>
1(c)	$2x^3 - 5x^2 - 4x + 3 = 0$ $\therefore (x+1)(2x^2 - 7x + 3) = 0$ $\therefore (x+1)(2x-1)(x-3) = 0$ $\therefore x = -1 \text{ or } x = \frac{1}{2} \text{ or } x = 3$	<ul style="list-style-type: none"> <li>✓ <math>(x+1)</math></li> <li>✓ <math>(2x^2 - 7x + 3)</math></li> <li>✓ <math>(2x-1)(x-3)</math></li> <li>✓ <math>x = -1 \text{ or } x = \frac{1}{2} \text{ or } x = 3</math> (4)</li> </ul>
1(d)	$-x^3 + 4x^2 - 2x - 4 = 0$ $\therefore x^3 - 4x^2 + 2x + 4 = 0$ $\therefore (x-2)(x^2 - 2x - 2) = 0$ $\therefore x = 2 \text{ or } x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(1)(-2)}}{2(1)}$ $x = \frac{2 \pm \sqrt{12}}{2}$ $x = 2,73 \text{ or } x = -0,73$	<ul style="list-style-type: none"> <li>✓ <math>(x-2)</math></li> <li>✓ <math>(x^2 - 2x - 2)</math></li> <li>✓ <math>x = 2</math></li> <li>✓ <math>x = \frac{2 \pm \sqrt{12}}{2}</math></li> <li>✓ <math>x = 2,73</math></li> <li>✓ <math>x = -0,73</math> (6)</li> </ul>
1(e)	$x^3 - 5x^2 - 3x + 9 = 0$ $\therefore (x+1)(x^2 - 6x + 9) = 0$ $\therefore (x+1)(x-3)^2 = 0$ $\therefore x = -1 \text{ or } x = 3$	<ul style="list-style-type: none"> <li>✓ <math>(x+1)</math></li> <li>✓ <math>(x^2 - 6x + 9)</math></li> <li>✓ <math>(x-3)^2</math></li> <li>✓ <math>x = -1 \text{ or } x = 3</math> (4)</li> </ul> <p>[23]</p>
2(a)	$0 = x^3 - 8x^2 + 19x - 12$ $\therefore (x-1)(x^2 - 7x + 12) = 0$ $\therefore (x-1)(x-4)(x-3) = 0$ $\therefore x = 1 \text{ or } x = 4 \text{ or } x = 3$ $(1; 0) \quad (4; 0) \quad (3; 0)$	<ul style="list-style-type: none"> <li>✓ <math>0 = x^3 - 8x^2 + 19x - 12</math></li> <li>✓ <math>(x-1)</math></li> <li>✓ <math>(x^2 - 7x + 12)</math></li> <li>✓ <math>(x-4)(x-3)</math></li> <li>✓ <math>x = 1 \text{ or } x = 4 \text{ or } x = 3</math></li> <li>✓ <math>(1; 0) \quad (4; 0) \quad (3; 0)</math> (6)</li> </ul>

<p>2(b)</p>	$0 = x^3 - x^2 - x - 2$ $\therefore (x - 2)(x^2 + x + 1) = 0$ $\therefore x = 2 \quad \text{or} \quad x = \frac{-1 \pm \sqrt{(1)^2 - 4(1)(1)}}{2(1)}$ $x = \frac{-1 \pm \sqrt{-3}}{2}$ <p>non-real</p>	<ul style="list-style-type: none"> <li>✓ <math>(x - 2)</math></li> <li>✓ <math>(x^2 + x + 1)</math></li> <li>✓ <math>x = 2</math></li> <li>✓ <math>x = \frac{-1 \pm \sqrt{-3}}{2}</math></li> <li>✓ <math>x = 2</math> is the only <math>x</math>-intercept</li> </ul> <p>(5)</p>
-------------	--	---

[11]




**TOPIC 1: SEQUENCES AND SERIES**

**Teacher Note:** Sequences and series is an exciting part of the curriculum. Make sure the learners know the difference between arithmetic and geometric sequences. They must also know the relevant formulae for finding specific terms and the sum of a certain number of terms. The sum to infinity is an important concept, as well as real world applications of the formulae.

**LESSON OVERVIEW**

1. Introduction session: 5 minutes
2. Typical exam questions:
  - Question 1: 10 minutes
  - Question 2: 10 minutes
  - Question 3: 10 minutes
3. Discussion of solutions: 25 minutes

**SECTION A: TYPICAL EXAM QUESTIONS****QUESTION 1**

- (a). Consider the sequence  $-2; 3; 8; 13; 18; 23; 28; 33; 38; \dots$
- (1) Determine the 100<sup>th</sup> term. (2)
  - (2) Determine the sum of the first 100 terms. (2)
- (b) The 13<sup>th</sup> and 7<sup>th</sup> terms of an arithmetic sequence are 15 and 51 respectively.  
Which term of the sequence is equal to  $-21$  (6)
- [10]

**QUESTION 2**

In a geometric sequence, the 6<sup>th</sup> term is 243 and the 3<sup>rd</sup> term is 72.

Determine:

- (a) the constant ratio. (4)
  - (b) the sum of the first 10 terms. (4)
- [8]

MATHEMATICS GRADE 12 SESSION 2 (TEACHER NOTES)

**QUESTION 3** (DoE Nov 2008 Paper 1)

Consider the sequence:  $\frac{1}{2}; 4; \frac{1}{4}; 7; \frac{1}{8}; 10; \dots$

- (a) If the pattern continues in the same way, write down the next TWO terms in the sequence. (2)
- (b) Calculate the sum of the first 50 terms of the sequence. (7)  
[9]

**QUESTION 4**

- (a) Calculate the value of  $\sum_{k=1}^{100} (2k - 1)$  (4)
- (b) Write the following series in sigma notation:  $2 + 5 + 8 + 11 + 14 + 17$  (4)
- (c) Calculate the value of  $\sum_{k=1}^{10} 5\left(\frac{1}{5}\right)^{k-1}$  (4)
- (d) Calculate the value of  $n$  if  $\sum_{k=1}^n 2^k = 2046$  (5)
- (e) Calculate:  $\sum_{k=1}^{\infty} 5\left(\frac{1}{5}\right)^{k-1}$  (5)  
[22]



**QUESTION 5**

Given the geometric series:  $8x^2 + 4x^3 + 2x^4 + \dots$

- (a) Determine the  $n^{\text{th}}$  term of the series. (3)
- (b) For what value(s) of  $x$  will the series converge? (2)
- (c) Calculate the sum of the series to infinity if  $x = \frac{3}{2}$ . (3)  
[8]

**SECTION B: SOLUTIONS AND HINTS TO SECTION A**

1(a)(1)	$T_n = a + (n - 1)d$ $\therefore T_{100} = -2 + (100 - 1)(5) = 493$	$\checkmark T_n = a + (n - 1)d$ $\checkmark T_{100} = 493$ (2)
---------	--	--

<p>1(a)(2)</p>	$S_n = \frac{n}{2}[2a + (n-1)d]$ $\therefore S_{100} = \frac{100}{2}[2(-2) + (100-1)(5)]$ $\therefore S_{100} = 24550$	$\checkmark S_n = \frac{n}{2}[2a + (n-1)d]$ $\checkmark T_{100} = 493$ <p style="text-align: right;">(2)</p>
<p>1(b)</p>	$T_{13} = 15$ $\therefore a + 12d = 15$ $\therefore a + 12d = 15 \dots A$ $\therefore a + 6d = 51 \dots B$ $\therefore 6d = -36 \quad A - B$ $\therefore d = -6$ $\therefore a + 12(-6) = 15$ $\therefore a - 72 = 15$ $\therefore a = 87$ $T_n = -21$ $\therefore a + (n-1)d = -21$ $\therefore 87 + (n-1)(-6) = -21$ $\therefore 87 - 6n + 6 = -21$ $\therefore n = 19$ $\therefore T_{19} = -21$	$\checkmark a + 12d = 15$ $\checkmark a + 6d = 51$ $\checkmark d = -6$ $\checkmark a = 87$ $\checkmark 87 + (n-1)(-6) = -21$ $\checkmark n = 19$ <p style="text-align: right;">(6)</p>



[10]

<p>2(a)</p>	$T_6 = 243 \quad \text{AND} \quad T_3 = 72$ $a.r^5 = 243 \dots A$ $a.r^5 = 243 \dots A$ $a.r^2 = 72 \dots B$ $\therefore r^3 = \frac{27}{8} \dots A \div B$ $\therefore r = \frac{3}{2}$	$\checkmark a.r^5 = 243$ $\checkmark a.r^2 = 72$ $\checkmark r^3 = \frac{27}{8}$ $\checkmark r = \frac{3}{2}$ <p style="text-align: right;">(4)</p>
<p>2(b)</p>	$\therefore a \left(\frac{3}{2}\right)^5 = 243$ $\therefore a = 32$ $\therefore S_{10} = \frac{32 \left( \left(\frac{3}{2}\right)^{10} - 1 \right)}{\frac{3}{2} - 1} = 3626,5625$	$\checkmark a \left(\frac{3}{2}\right)^5 = 243$ $\checkmark a = 32$ $\checkmark S_{10} = \frac{32 \left( \left(\frac{3}{2}\right)^{10} - 1 \right)}{\frac{3}{2} - 1}$ $\checkmark \text{answer}$ <p style="text-align: right;">(4) [8]</p>




MATHEMATICS GRADE 12 SESSION 2 (TEACHER NOTES)

3(a)	$\frac{1}{16}; 13$	✓ ✓ answers (2)
3(b)	$S_{50} = 25 \text{ terms of } 1^{\text{st}} \text{ sequence} + 25 \text{ terms of } 2^{\text{nd}} \text{ sequence}$ $S_{50} = \left( \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots \text{to } 25 \text{ terms} \right) + (4 + 7 + 10 + 13 + \dots \text{to } 25 \text{ terms})$  $S_{50} = \frac{\frac{1}{2} \left( \left( \frac{1}{2} \right)^{25} - 1 \right)}{\frac{1}{2} - 1} + \frac{25}{2} [2(4) + 24(3)]$  $S_{50} = 0,999999\dots + 1000$ $S_{50} = 1001,00$	✓ separating into an arithmetic and geometric series  $\frac{1}{2} \left( \left( \frac{1}{2} \right)^{25} - 1 \right)$ $\frac{1}{2} - 1$ ✓ correct formulae ✓ ✓ $\frac{25}{2} [2(4) + 24(3)]$ ✓ answer (7)

[9]

4(a)	$\sum_{k=1}^{100} (2k-1) = [2(1)-1] + [2(2)-1] + [2(3)-1] + [2(4)-1] + \dots + [2(100)-1]$ $= 1 + 3 + 5 + 7 + \dots + 199$  We can use the formula $S_n = \frac{n}{2} [a+l]$ to calculate the sum: $S_{100} = \frac{100}{2} [1+199]$ $\therefore S_{100} = 10000$	✓ expanding ✓ correct formula ✓ substitution ✓ answer (4)
4(b)	The series is arithmetic. There are also 6 terms in the series. $a = 2 \quad d = 3 \quad n = 6$ We can determine the general term as follows: $T_n = a + (n-1)d$ $\therefore T_n = 2 + (n-1)(3)$ $\therefore T_n = 2 + 3n - 3$ $\therefore T_n = 3n - 1$ We can now write the series in sigma notation as follows: $\sum_{n=1}^6 (3n-1)$	✓ correct formula ✓ $T_n = 3n - 1$ ✓ $n = 6$ ✓ $\sum_{n=1}^6 (3n-1)$ (4)
4(c)	$\sum_{k=1}^{10} 5 \left( \frac{1}{5} \right)^{k-1} = 5 \left( \frac{1}{5} \right)^0 + 5 \left( \frac{1}{5} \right)^1 + 5 \left( \frac{1}{5} \right)^2 + \dots + 5 \left( \frac{1}{5} \right)^9$ $= 5 \left( \frac{1}{5} \right)^0 + 5 \left( \frac{1}{5} \right)^1 + 5 \left( \frac{1}{5} \right)^2 + \dots + 5 \left( \frac{1}{5} \right)^9$ $= 5 + 1 + \frac{1}{5} + \dots + \frac{1}{5^8}$	✓ expanding ✓ correct formula ✓ substitution ✓ answer (4)

	<p>This is a geometric series with:</p> $a = 5 \quad \text{and} \quad r = \frac{1}{5}$ $S_{10} = \frac{5 \left[ 1 - \left( \frac{1}{5} \right)^{10} \right]}{1 - \frac{1}{5}} = \frac{25}{4} \left[ 1 - \left( \frac{1}{5} \right)^{10} \right] = 6,25$	
<p>4(d)</p>	<p><math>\sum_{k=1}^n 2^k = 2^1 + 2^2 + 2^3 + \dots</math></p> <p><math>= 2 + 4 + 8 + 16 + \dots</math></p> <p>This is a geometric series with:</p> <p><math>a = 2 \quad r = 2 \quad S_n = 2046</math></p> $S_n = \frac{a(r^n - 1)}{r - 1}$ $\therefore 2046 = \frac{(2)(2^n - 1)}{2 - 1}$ $\therefore 2046 = (2)(2^n - 1)$ $\therefore 1023 = 2^n - 1$ $\therefore 1024 = 2^n$ $\therefore 2^{10} = 2^n$ $\therefore n = 10$ 	<p>✓ expanding</p> <p>✓ correct formula</p> <p>✓ <math>2046 = \frac{(2)(2^n - 1)}{2 - 1}</math></p> <p>✓ <math>1024 = 2^n</math></p> <p>✓ <math>n = 10</math></p> <p>(5)</p>
<p>4(e)</p>	$\sum_{k=1}^{\infty} 5 \left( \frac{1}{5} \right)^{k-1}$ $= 5 \left( \frac{1}{5} \right)^{1-1} + 5 \left( \frac{1}{5} \right)^{2-1} + 5 \left( \frac{1}{5} \right)^{3-1} + \dots$ $= 5 \left( \frac{1}{5} \right)^0 + 5 \left( \frac{1}{5} \right)^1 + 5 \left( \frac{1}{5} \right)^2 + \dots$ $= 5 + 1 + \frac{1}{5} + \dots$ <p>The series is clearly geometric with the constant ratio lying in the interval <math>-1 &lt; r &lt; 1</math>.</p> <p>For this convergent geometric series <math>a = 5</math> and <math>r = \frac{1}{5}</math></p> $S_{\infty} = \frac{a}{1 - r} = \frac{5}{1 - \frac{1}{5}} = \frac{5}{\frac{4}{5}} = \frac{25}{4}$	<p>✓ expanding</p> <p>✓ stating that <math>-1 &lt; r &lt; 1</math></p> <p>✓ correct formula</p> <p>✓ substitution</p> <p>✓ answer</p> <p>(5)</p>

5(a)	$T_n = (8x^3)\left(\frac{1}{2}x\right)^{n-1}$	✓ correct formula ✓ $a = 8x^3$ ✓ $r = \frac{1}{2}x$	(3)
5(b)	$-1 < \frac{x}{2} < 1$ $= -2 < x < 2$	✓ $-1 < \frac{x}{2} < 1$ ✓ $-2 < x < 2$	(2)
5(c)	$S_\infty = \frac{a}{1-r}$ $\therefore S_\infty = \frac{8x^2}{1-\frac{x}{2}}$ $\therefore S_\infty = \frac{8\left(\frac{3}{2}\right)^2}{1-\frac{1}{2}\left(\frac{3}{2}\right)}$ $\therefore S_\infty = 72$	✓ correct formula ✓ substitution ✓ answer	(3)
			<b>[8]</b>

**SECTION C: HOMEWORK**



**QUESTION 1**

The 19<sup>th</sup> term of an arithmetic sequence is 11, while the 31<sup>st</sup> term is 5.

- (a) Determine the first three terms of the sequence. (5)
- (b) Which term of the sequence is equal to -29? (3)
- [8]

**QUESTION 2**

Given:  $\frac{1}{181} + \frac{2}{181} + \frac{3}{181} + \frac{4}{181} + \dots + \frac{180}{181}$

- (a) Calculate the sum of the given series. (4)
- (b) Hence calculate the sum of the following series:

$$\left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{2}{3}\right) + \left(\frac{1}{4} + \frac{2}{4} + \frac{3}{4}\right) + \dots + \left(\frac{1}{181} + \frac{2}{181} + \dots + \frac{180}{181}\right)$$

(4)

[8]

MATHEMATICS

GRADE 12

SESSION 2

(TEACHER NOTES)

**QUESTION 3** (DoE Feb 2009 Paper 1)

The following is an arithmetic sequence:  $1 - p ; 2p - 3 ; p + 5 ; \dots$

- (a) Calculate the value of  $p$ . (3)
- (b) Write down the value of:
- (1) The first term of the sequence. (1)
  - (2) The common difference. (2)
- (c) Explain why none of the numbers in this arithmetic sequence are perfect squares. (2)
- [8]

**QUESTION 4**

In a geometric sequence in which all terms are positive, the sixth term is  $\sqrt{3}$  and the eighth term is  $\sqrt{27}$ . Determine the first term and constant ratio. [7]

**QUESTION 5**

- (a) Determine  $n$  if  $\sum_{r=1}^n (6r - 1) = 456$  (7)

- (b) Prove that  $\sum_{k=3}^n (2k - 1)n = n^3 - 4n$ . [8]

[13]

**QUESTION 6**

Consider the series  $\sum_{n=1}^{\infty} 2\left(\frac{1}{2}x\right)^n$

- (a) For which values of  $x$  will the series converge? (3)
- (b) If  $x = \frac{1}{2}$ , calculate the sum to infinity of this series. (3)

[6]

**QUESTION 7**

(DoE Feb 2009 Paper 1)

A sequence of squares, each having side 1, is drawn as shown on the following page. The first square is shaded, and the length of the side of each shaded square is half the length of the side of the shaded square in the previous diagram.

MATHEMATICS

GRADE 12

SESSION 2

(TEACHER NOTES)



DIAGRAM 1




DIAGRAM 2




DIAGRAM 3




DIAGRAM 4

- (a) Determine the area of the unshaded region in DIAGRAM 3. (2)  
 (b) What is the sum of the areas of the unshaded regions on the first seven squares? (5)  
 [7]

**QUESTION 8**

A plant grows 1,5 m in 1<sup>st</sup> year. Its growth each year thereafter is  $\frac{2}{3}$  of its growth in the previous year. What is the greatest height it can reach? [3]

**SOLUTIONS TO HOMEWORK: SEQUENCES & SERIES (A)**

1(a)	$T_{19} = a + 18d = 11$ $T_{31} = a + 30d = 5$ $\therefore 12d = -6$ $\therefore d = -\frac{1}{2}$ $\therefore a + 18\left(-\frac{1}{2}\right) = 11$ $\therefore a = 20$ <div style="text-align: center; margin-top: 10px;">   <math>\therefore 20; 19\frac{1}{2}; 19\dots</math> </div>	<ul style="list-style-type: none"> <li>✓ <math>T_{19} = a + 18d = 11</math></li> <li>✓ <math>T_{31} = a + 30d = 5</math></li> <li>✓ <math>d = -\frac{1}{2}</math></li> <li>✓ <math>a = 20</math></li> <li>✓ sequence</li> </ul> <p style="text-align: right;">(5)</p>
1(b)	$20 + (n-1)\left(-\frac{1}{2}\right) = -29$ $\therefore (n-1)\left(-\frac{1}{2}\right) = -49$ $\therefore n-1 = 98$ $\therefore n = 99$ $\therefore T_{99} = -29$	<ul style="list-style-type: none"> <li>✓ Substitution into formula</li> <li>✓ equating to -29</li> <li>✓ <math>n = 99</math></li> </ul> <p style="text-align: right;">(3)</p>
<b>[8]</b>		

MATHEMATICS

GRADE 12

SESSION 2

(TEACHER NOTES)

<p>2(a)</p>	$\frac{1}{181} + \frac{2}{181} + \frac{3}{181} + \frac{4}{181} + \dots + \frac{180}{181}$ $a = \frac{1}{181} \quad d = \frac{1}{181} \quad n = 180$ $S_{180} = \frac{180}{2} \left[ 2 \left( \frac{1}{181} \right) + (179) \frac{1}{181} \right] = 90[1] = 90$	<ul style="list-style-type: none"> <li>✓ correct <math>a</math> and <math>d</math></li> <li>✓ correct <math>n</math></li> <li>✓ <math>S_n</math> formula</li> <li>✓ correct answer</li> </ul> <p style="text-align: right;">(4)</p>
<p>2(b)</p>	$\left( \frac{1}{2} \right) + \left( \frac{1}{3} + \frac{2}{3} \right) + \left( \frac{1}{4} + \frac{2}{4} + \frac{3}{4} \right) + \dots + \left( \frac{1}{181} + \frac{2}{181} + \dots + \frac{180}{181} \right)$ $= \frac{1}{2} + 1 + 1\frac{1}{2} + 2 + \dots + 90 \quad [a = \frac{1}{2} \quad d = \frac{1}{2} \quad T_n = 90]$ $\therefore \frac{1}{2} + (n-1)\frac{1}{2} = 90$ $\therefore 1 + n - 1 = 180$ $\therefore n = 180$ $\therefore S_{180} = \frac{180}{2} \left[ \frac{1}{2} + 90 \right] = 90 \left[ 90\frac{1}{2} \right] = 8145$	<ul style="list-style-type: none"> <li>✓ simplifying fractions to get series</li> <li>✓ <math>\frac{1}{2} + (n-1)\frac{1}{2} = 90</math></li> <li>✓ <math>n = 180</math></li> <li>✓ substitution into <math>S_n</math> formula to get 8145</li> </ul> <p style="text-align: right;">(4)</p>

[8]

<p>3(a)</p>	$(2p-3) - (1-p) = (p+5) - (2p-3)$ $2p-3-1+p = p+5-2p+3$ $3p-4 = -p+8$ $4p = 12$ $p = 3$	<ul style="list-style-type: none"> <li>✓ <math>T_2 - T_1 = T_3 - T_2</math></li> <li>✓ <math>2p-3-1+p = p+5-2p+3</math></li> <li>✓ <math>p = 3</math></li> </ul> <p style="text-align: right;">(3)</p>
<p>3(b)(1)</p>	$T_1 = 1 - (3) = -2$	<ul style="list-style-type: none"> <li>✓ answer</li> </ul> <p style="text-align: right;">(1)</p>
<p>3(b)(2)</p>	$d = T_2 - T_1 = (2p-3) - (1-p)$ $\therefore d = (2(3)-3) - (1-3)$ $\therefore d = 3 - (-2)$ $\therefore d = 5$	<ul style="list-style-type: none"> <li>✓ <math>(2p-3) - (1-p)</math></li> <li>✓ <math>p = 3</math></li> </ul> <p style="text-align: right;">(2)</p>
<p>3(c)</p>	<p>The sequence is <math>-2; 3; 8; 13; 18; 23; 28; 33; 38; \dots</math></p> <p>After the first term <math>-2</math>, all the other terms end in either a 3 or an 8.</p> <p>Perfect squares never end in a 3 or an 8.</p>	<ul style="list-style-type: none"> <li>✓✓ answer</li> </ul> <p style="text-align: right;">(2)</p> <p style="text-align: right;">[8]</p>

4	$ar^5 = \sqrt{3}$ $ar^7 = \sqrt{27}$ $\therefore \frac{ar^7}{ar^5} = \frac{\sqrt{27}}{\sqrt{3}}$ $\therefore r^2 = \sqrt{\frac{27}{3}}$ $\therefore r^2 = \sqrt{9}$ $\therefore r^2 = 3$ $\therefore r = \sqrt{3} \quad (\text{terms are positive})$ $\therefore a(\sqrt{3})^5 = \sqrt{3}$ $\therefore a = \frac{\sqrt{3}}{(\sqrt{3})^5}$ $\therefore a = \frac{1}{(\sqrt{3})^4}$ $\therefore a = \frac{1}{(3^{\frac{1}{2}})^4}$ $\therefore a = \frac{1}{9}$	<ul style="list-style-type: none"> <li>✓ <math>ar^5 = \sqrt{3}</math></li> <li>✓ <math>ar^7 = \sqrt{27}</math></li> <li>✓ <b>dividing</b></li> <li>✓ <math>r^2 = 3</math></li> <li>✓ <math>r = \sqrt{3}</math></li> <li>✓ <b>correct working with surds</b></li> <li>✓ <math>a = \frac{1}{9}</math></li> </ul>
---	---	--




[7]

**SOLUTIONS TO HOMEWORK: SEQUENCES & SERIES (B)**

5(a)	$\sum_{r=1}^n (6r-1) = [6(1)-1] + [6(2)-1] + [6(3)-1] + \dots + [6(n)-1] = 456$ $= 5 + 11 + 17 + \dots + (6n-1) = 456$ $S_n = \frac{n}{2}(2a + (n-1)d)$ $\therefore 456 = \frac{n}{2}(2a + (n-1)d)$ $\therefore 456 = \frac{n}{2}(2(5) + (n-1)(6))$ $\therefore 456 = \frac{n}{2}(10 + 6n - 6)$ $\therefore 456 = \frac{n}{2}(4 + 6n)$ $\therefore 456 = 2n + 3n^2$ $\therefore 0 = 3n^2 + 2n - 456$	<ul style="list-style-type: none"> <li>✓ expanding</li> <li>✓ correct formula</li> <li>✓ <math>456 = \frac{n}{2}(2a + (n-1)d)</math></li> <li>✓ <math>0 = 3n^2 + 2n - 456</math></li> <li>✓ <math>(3n+38)(n-12) = 0</math></li> <li>✓ <math>n = -\frac{38}{3}</math> or <math>n = 12</math></li> <li>✓ <math>\therefore n = 12</math></li> </ul>
------	--	--

(7)

	$\therefore (3n+38)(n-12) = 0$ $\therefore 3n = -38 \text{ or } n = 12$ $\therefore n = -\frac{38}{3} \text{ or } n = 12$ $\therefore n = 12$	
5(b)	$\sum_{k=3}^n [(2k-1)n] = 5n + 7n + 9n + \dots + [(2n-1)n]$ $\therefore a = 5n, d = 2n \text{ and number of terms} = n - 2$ $\therefore S_{n-2} = \frac{n-2}{2} [2a + (n-2-1)d]$ $\therefore S_{n-2} = \frac{n-2}{2} [2(5n) + (n-3)(2n)]$ $\therefore S_{n-2} = \frac{n-2}{2} [10n + 2n^2 - 6n]$ $\therefore S_{n-2} = \frac{n-2}{2} [2n^2 + 4n]$ $\therefore S_{n-2} = 2n(n-2) + n^2(n-2)$ $\therefore S_{n-2} = 2n^2 - 4n + n^3 - 2n^2 = n^3 - 4n$	<ul style="list-style-type: none"> <li>✓ expanding</li> <li>✓ <math>a = 5n, d = 2n</math></li> <li>✓ number of terms = <math>n - 2</math></li> <li>✓ correct formula</li> <li>✓ substitution</li> <li>✓ answer</li> </ul> <p style="text-align: right;">(6)</p> <p style="text-align: right;"><b>[13]</b></p>
6(a)	$\sum_{n=1}^{\infty} 2\left(\frac{1}{2}x\right)^n$  $= 2\left(\frac{1}{2}x\right)^1 + 2\left(\frac{1}{2}x\right)^2 + 2\left(\frac{1}{2}x\right)^3 + 2\left(\frac{1}{2}x\right)^4 + \dots$ $= x + \frac{1}{2}x^2 + \frac{1}{4}x^3 + \frac{1}{8}x^4 + \dots$ <p>The series converges for:</p> $-1 < \frac{1}{2}x < 1$ $\therefore -2 < x < 2$	<ul style="list-style-type: none"> <li>✓ <math>r = \frac{1}{2}x</math></li> <li>✓ <math>-1 &lt; \frac{1}{2}x &lt; 1</math></li> <li>✓ <math>-2 &lt; x &lt; 2</math></li> </ul> <p style="text-align: right;">(3)</p>
6(b)	$a = \frac{1}{2} \quad r = \frac{1}{2} \left( \frac{1}{2} \right) = \frac{1}{4}$ $\therefore S_{\infty} = \frac{\frac{1}{2}}{1 - \frac{1}{4}} = \frac{2}{3}$	<ul style="list-style-type: none"> <li>✓ <math>a</math> and <math>r</math></li> <li>✓ <math>S_{\infty}</math> formula</li> <li>✓ <math>\frac{2}{3}</math></li> </ul> <p style="text-align: right;">(3)</p> <p style="text-align: right;"><b>[6]</b></p>



Question 7:




DIAGRAM 1




DIAGRAM 2




DIAGRAM 3




DIAGRAM 4

7(a)	Area of unshaded square = Area of large square – Area of small shaded square $= (1)(1) - \left(\frac{1}{4}\right)\left(\frac{1}{4}\right)$ $= 1 - \frac{1}{16} = \frac{15}{16}$	$\checkmark (1)(1) - \left(\frac{1}{4}\right)\left(\frac{1}{4}\right)$ $\checkmark \frac{15}{16}$ <p style="text-align: right;">(2)</p>
7(b)	Sum of the unshaded areas of the first seven squares: $= (1-1) + \left(1 - \frac{1}{4}\right) + \left(1 - \frac{1}{4^2}\right) + \dots + \left(1 - \frac{1}{4^6}\right)$ $= 7 - \left(1 + \frac{1}{4} + \frac{1}{4^2} + \dots + \frac{1}{4^6}\right)$ $= 7 - \left(\frac{1\left(1 - \left(\frac{1}{4}\right)^7\right)}{1 - \frac{1}{4}}\right)$ $= 7 - 1,333251953\dots$ $= 5,67$	$\checkmark \checkmark$ Getting the pattern for the unshaded areas $\checkmark$ correct formula $\checkmark$ substitution $\checkmark$ answer <p style="text-align: right;">(5)</p> <p style="text-align: right;">[7]</p>
8	$\therefore S_{\infty} = \frac{1,5}{1 - \left(\frac{2}{3}\right)}$ $\therefore S_{\infty} = 4,5 \text{ m}$ Thus the greatest height is 4,5 m	$\checkmark$ correct formula $\checkmark$ substitution $\checkmark$ answer <p style="text-align: right;">[3]</p>

**FINANCIAL MATHEMATICS**

**Teacher Note:** This session on Financial Mathematics will deal with **future and present value annuities**. A future value annuity is a savings plan for the future, whereas a present value annuity is a loan. There are two types of loans dealt with in this session: The first loan is one in which a learner borrows money from a bank and has to pay a certain number of equal repayments with interest. The second type of loan is one in which the learner deposits a large sum of money into the bank and the bank pays the learner equal amounts with interest over a given time period.

**LESSON OVERVIEW**

- |    |                                |            |
|----|--------------------------------|------------|
| 1. | Review Homework from Session 2 | 10 minutes |
| 2. | Introduction session:          | 10 minutes |
| 3. | Typical exam questions:        |            |
|    | Question 1:                    | 15 minutes |
|    | Question 2:                    | 15 minutes |
| 4. | Discussion of solutions:       | 35 minutes |
| 5. | Discussion of Homework         | 5 minutes  |

**SECTION A: TYPICAL EXAM QUESTIONS****QUESTION 1      15 minutes**

- (a) Suppose that at the beginning of the month, R1000 is deposited into a bank. At the end of that month, a further R1000 is deposited and a further R1000 at the end of the next month. This continues for eight years. If the interest rate is 6% per annum compounded monthly, how much will have been saved after the eight year period? (4)
- (b) Patrick decided to start saving money for a period of eight years starting on 31 December 2009. At the end of January 2010 (in one month's time), he deposited R2300 into the savings plan. Thereafter, he continued making deposits of R2300 at the end of each month for the planned eight year period. The interest rate remained fixed at 10% per annum compounded monthly. (4)
- (1) How much will he have saved at the end of his eight year plan which started on the 31 December 2009? (4)
- (2) If Patrick leaves the accumulated amount in the bank for a further three months, what will the investment then be worth? (4)

[16]

**QUESTION 2 15 minutes**

(a) David takes out a bank loan to pay for his new car. He repays the loan by means of monthly payments of R4000 for a period of five years starting one month after the granting of the loan. The interest rate is 24% per annum compounded monthly. Calculate the purchase price of his new car. (4)

(b) Peter inherits R400 000 from his father. He invests the money at an interest rate of 12% per annum compounded monthly. He wishes to earn a monthly salary

from the investment for a period of twenty years starting in one month's time.


How much will he receive each month? (4)

[8]

**SECTION B: SOLUTIONS AND HINTS TO SECTION A**


**QUESTION 1**

(a) Draw a timeline




<p>(a)</p> $F = \frac{x[(1+i)^n - 1]}{i}$ $\therefore F = \frac{1000 \left[ \left( 1 + \frac{0,06}{12} \right)^{97} - 1 \right]}{\frac{0,06}{12}}$ $\therefore F = R124\,442,68$	<p>In this example, the duration of the loan is 8 years (96 months). However, the number of payments made is 97 because of the first payment being made immediately. This means that in the first month, two payments of R1000 were made.</p>	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>n = 97</math></li> <li>✓ <math>\frac{0,06}{12}</math></li> <li>✓ answer</li> </ul> <p>(4)</p>
--	---	---

(b)(1) Draw a timeline



(b)(1)	$F = \frac{x[(1+i)^n - 1]}{i}$ $\therefore F = \frac{2300 \left[ \left(1 + \frac{0,10}{12}\right)^{96} - 1 \right]}{\frac{0,10}{12}}$ $\therefore F = R336\,216,47$	<div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;">                 In this example, the duration of the loan is 8 years (96 months). However, the number of payments made is 96 because of the first payment being made one month after the start of the savings plan.             </div> <ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>n = 96</math></li> <li>✓ <math>\frac{0,10}{12}</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p>
--------	---	---

(b)(2) Draw a timeline




(b)(2)	$A = P(1+i)^n$ $\therefore A = 336\,216,47 \left(1 + \frac{0,10}{12}\right)^3$ $\therefore A = R344\,692,12$	<div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;">                 Since there will no longer be any further payments of R2300 into the annuity, all we now need to do is grow the R336 216,47 for three months using the formula <math>A = P(1+i)^n</math> to calculate the future value of the investment after the further three months.             </div> <ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>n = 3</math></li> <li>✓ <math>P = 336\,216,47</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p>
--------	--	---

[16]


QUESTION 2

(a) Draw a timeline



(a)	$P = \frac{x[1 - (1+i)^{-n}]}{i}$ $\therefore P = \frac{4000 \left[ 1 - \left( 1 + \frac{0,24}{12} \right)^{-60} \right]}{\frac{0,24}{12}}$ $\therefore P = R139043,55$	<div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;">                 There must always be a gap between the loan (P) and the first payment in order for the present value formula to work.             </div> <ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>n = 60</math></li> <li>✓ <math>\frac{0,24}{12} = 0,02</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p>
-----	---	--

(b) Draw a timeline



(b)	$400000 = \frac{x \left[ 1 - \left( 1 + \frac{0,12}{12} \right)^{-240} \right]}{\frac{0,12}{12}}$ $\therefore \frac{400000 \times 0,01}{\left[ 1 - (1,01)^{-240} \right]} = x$ $\therefore x = R4404,34$	<div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;">                 This is an example of a loan where you pay money into a bank and the bank pays you monthly amounts with interest.             </div> <ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>n = 240</math></li> <li>✓ <math>\frac{0,12}{12} = 0,01</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p>
-----	--	---

[8]

### SECTION C: HOMEWORK

#### QUESTION 1

Mpho takes out a retirement annuity that will supplement his pension when he retires in thirty years' time. He estimates that he will need R2 500 000 in this retirement fund at that stage. The interest rate he earns is 9% per annum compounded monthly.

- (a) Calculate his monthly payment into this fund if he starts paying immediately and makes his final payment in 30 years' time. (5)

- (b) The retirement fund does not pay out the R2 500 000 million when Mpho retires. Instead he will be paid monthly amounts, for a period of twenty years, starting one month after his retirement. If the interest that he earns over this period is calculated at 7% per annum compounded monthly, determine the monthly payments he will receive. (5)

[10]

**QUESTION 2**


Simphiwe takes out a twenty year loan of R100 000. She repays the loan by means of equal monthly payments starting **three months** after the granting of the loan. The interest rate is 18% per annum compounded monthly. Calculate the monthly payments. [5]

**SECTION D: SOLUTIONS TO SESSION 3 HOMEWORK: TOPIC 1: FINANCIAL MATHEMATICS A**

**QUESTION 1**

(a)	$2\,500\,000 = \frac{x \left[ (1,0075)^{361} - 1 \right]}{0,0075}$ $\therefore \frac{2\,500\,000 \times 0,0075}{\left[ (1,0075)^{361} - 1 \right]} = x$ $\therefore x = \text{R}1354,67$	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>n = 361</math></li> <li>✓ <math>\frac{0,09}{12} = 0,0075</math></li> <li>✓ <math>F = 2\,500\,000</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(5)</p>
(b)	$2\,500\,000 = \frac{x \left[ 1 - \left( 1 + \frac{0,07}{12} \right)^{-240} \right]}{\left( \frac{0,07}{12} \right)}$ $\therefore \frac{2\,500\,000 \times \left( \frac{0,07}{12} \right)}{\left[ 1 - \left( 1 + \frac{0,07}{12} \right)^{-240} \right]} = x$ $\therefore x = \text{R}19\,382,47$	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>n = 240</math></li> <li>✓ <math>\frac{0,07}{12}</math></li> <li>✓ <math>P = 2\,500\,000</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(5)</p> <p style="text-align: right;"><b>[10]</b></p>

QUESTION 2



(a)	$\therefore 100\,000(1,015)^2 = \frac{x[1 - (1,015)^{-238}]}{0,015}$ $\therefore 100\,000(1,015)^2 \times 0,015 = x[1 - (1,015)^{-238}]$ $\therefore \frac{100\,000(1,015)^2 \times 0,015}{[1 - (1,015)^{-238}]} = x$ $\therefore x = \text{R}1591,35$	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>n = 238</math></li> <li>✓ <math>\frac{0,18}{12} = 0,015</math></li> <li>✓</li> <li><math>100\,000(1,015)^2</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">[5]</p>
-----	--	--







(2) One month after purchasing his present tractor, the farmer deposited  $x$  rands into an account that pays interest at a rate of 12% p.a., compounded monthly. He continued to deposit the same amount at the end of each month for a total of 60 months. At the end of sixty months he has exactly the amount that is needed to purchase a new tractor, after he trades in his present tractor.

Calculate the value of  $x$ . (6)


(b) Suppose that twelve months after the purchase of the present tractor and every twelve months thereafter, he withdraws R5 000 from his account, to pay for maintenance of the tractor. If he makes five such withdrawals, what will the new monthly deposit be?

(4)

[13]


**SECTION B: SOLUTIONS AND HINTS TO SECTION A**


1(a) Draw a timeline



<p>1(a)</p> $300\,000 = \frac{4000 \left[ (1,0125)^n - 1 \right]}{0,0125}$ $\therefore \frac{300\,000 \times 0,0125}{4000} = (1,0125)^n - 1$ $\therefore \frac{300\,000 \times 0,0125}{4000} + 1 = (1,0125)^n$ $\therefore 1,9375 = (1,0125)^n$ $\therefore \log_{1,0125} 1,9375 = n$ $\therefore n = 53,24189314$ <p>It will take Anna 53 months and approximately 7-8 days depending on the number of days in the 54<sup>th</sup> month.</p>	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>\frac{0,15}{12} = 0,0125</math></li> <li>✓ <math>1,9375 = (1,0125)^n</math></li> <li>✓ <math>\log_{1,0125} 1,9375 = n</math></li> <li>✓ <math>n = 53,24189314</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(6)</p>
--	---

1(b) Draw a timeline




1(b)	$500\,000 = \frac{8000 [1 - (1,015)^{-n}]}{0,015}$ $\therefore \frac{500\,000 \times 0,015}{8000} = 1 - (1,015)^{-n}$ $\therefore (1,015)^{-n} = 1 - \frac{500\,000 \times 0,015}{8000}$ $\therefore (1,015)^{-n} = 0,0625$ $\therefore 0,0625 = (1,015)^{-n}$ $\therefore \log_{1,015} 0,0625 = -n$ $\therefore -186,2221025 = -n$ $\therefore n = 186,2221025$ <p style="text-align: center;"></p> <p>There will be 186 payments of R8000 into the annuity. The decimal here indicates that there will be a final payment which is less than R8000.</p>	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>\frac{0,18}{12} = 0,015</math></li> <li>✓ <math>(1,015)^{-n} = 0,0625</math></li> <li>✓</li> <li><math>\log_{1,015} 0,0625 = -n</math></li> <li>✓ <math>n = 186,2221025</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(6)</p>
------	--	---

[12]


2(a)(1)	$A = 800000(1,08)^5$ $\therefore A = 1175462,46$ $\therefore 1175462,46 - 200\,000$ $= R975462,46$	<ul style="list-style-type: none"> <li>✓ <math>800000(1,08)^5</math></li> <li>✓ <math>\log_{1,12} \frac{23000}{1570} = n</math></li> <li>✓ <math>n = 23,69</math> years</li> </ul> <p style="text-align: right;">(3)</p>
---------	--	--

2(a) (2) Draw a time line



<p>2(a)(2)</p>	$F = \frac{x[(1+i)^n - 1]}{i}$ $\therefore 975462,46 = \frac{x[(1,01)^{60} - 1]}{0,01}$ $\therefore \frac{975462,46 \times 0,01}{[(1,01)^{60} - 1]} = x$ $\therefore x = R\ 11944,00$	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>n = 60</math></li> <li>✓ <math>\frac{0,12}{12} = 0,01</math></li> <li>✓ <math>F = 975462,46</math></li> <li>✓✓</li> <li><math>x = R\ 11944,00</math></li> </ul> <p style="text-align: right;">(6)</p>
----------------	---	---

2(b) Draw a time line



<p>2(b)</p>	<p>Services</p> $= [5000(1,01)^{48} + 5000(1,01)^{36} + 5000(1,01)^{24} + 5000(1,01)^{12} + 5000]$ $= 32197,77$ $975462,46 + \text{services} = x \frac{[1,01]^{60} - 1}{0,01}$ $975462,46 + 32197,77 = 81,66966986x$ $x = R12338,24$	<ul style="list-style-type: none"> <li>✓ services</li> <li>✓ <math>975462,46 + \text{services}</math></li> <li>✓ <math>\frac{x[1,01]^{60} - 1}{0,01}</math></li> <li>✓ <math>x = R12338,24</math></li> </ul> <p style="text-align: right;">(4)</p>
-------------	--	--

[13]

**SECTION C: HOMEWORK****QUESTION 1**

Mark's small business, called Postal Emporium, purchases a photocopying machine for R200 000. The photocopy machine depreciates in value at 20% per annum on a reducing balance. Mark's business wants to buy a new machine in five years' time. A new machine will cost much more in the future and its cost will escalate at 16% per annum effective. The old machine will be sold at scrap value after five years. A sinking fund is set up immediately in order to save up for the new machine. The proceeds from the sale of the old machine will be used together with the sinking fund to buy the new machine. The small business will pay equal monthly amounts into the sinking fund, and the interest earned is 18% per annum compounded monthly. The first payment will be made immediately, and the last payment will be made at the end of the five year period.

- (a) Find the scrap value of the old machine. (2)
- (b) Find the cost of the new machine in five years' time. (2)
- (c) Find the amount required in the sinking fund in five years' time. (1)
- (d) Find the equal monthly payments made into the sinking fund. (4)
- (e) Suppose that the business decides to service the machine at the end of each year for the five year period. R3000 will be withdrawn from the sinking fund at the end of each year starting one year after the original machine was bought.
- (1) Calculate the reduced value of the sinking fund at the end of the five year period due to these withdrawals. (3)
- (2) Calculate the increased monthly payment into the sinking fund which will yield the original sinking fund amount as well as allow for withdrawals from the fund for the services. (5)
- [17]

**QUESTION 2**

- (a) R2000 is immediately deposited into a savings account. Six months later and every six months thereafter, R2000 is deposited into the account. The interest rate is 16% p.a. compounded half-yearly. How long will it take to accumulate R100 000? (7)
- (b) How long will it take to repay a loan of R400 000 if the first quarterly payment of R17000 is made three months after the granting of the loan and the interest rate

is 16% per annum compounded quarterly? (6)


[13]

**SECTION D: SOLUTIONS TO HOMEWORK: FINANCIAL MATHEMATICS**

**QUESTION 1**


1(a)	$A = 200\,000(1 - 0,2)^5$ $\therefore A = R65\,536$	✓ correct formula ✓ answer (2)
1(b)	$A = 200\,000(1 + 0,16)^5$ $\therefore A = R420\,068,33$	✓ correct formula ✓ answer (2)
1(c)	Sinking fund = $420\,068,33 - 65\,536$ Sinking fund = $354\,532,33$	✓ answer (1)

1(d) Draw a time line



1(d)	$354\,532,33 = \frac{x[(1,015)^{61} - 1]}{0,015}$ $\frac{354\,532,33 \times 0,015}{[(1,015)^{61} - 1]} = x$ $\therefore x = R3593,55$	✓ correct formula ✓ $F = 354\,532,33$ ✓ $n = 61$ ✓ $\frac{0,18}{12} = 0,015$ ✓ $x = R3593,55$ (4)
------	---	--

1(e) (1) Draw a time line



<p>1(e)(1)</p>	<p>Future value of the withdrawals:</p> $3000\left(1 + \frac{0,18}{12}\right)^{48} + 3000\left(1 + \frac{0,18}{12}\right)^{36}$ $+ 3000\left(1 + \frac{0,18}{12}\right)^{24} + 3000\left(1 + \frac{0,18}{12}\right)^{12} + 3000$ $= R22\ 133,22$ <p>The reduced value of the sinking fund will be:</p> $R354\ 532,33 - R22\ 133,22 = R332\ 399,11$	<ul style="list-style-type: none"> <li>✓ services</li> <li>✓ R22 133,22</li> <li>✓ reduced value (3)</li> </ul>
<p>1(e)(2)</p>	<p>If we add R22 133,22 to the original sinking fund amount of R354 532,33, then it will be possible not only to receive the sinking fund amount of R354 532,33 at the end of the five year period, but also be able to make the service withdrawals at the end of each year for the five year period.</p> $354\ 532,33 + 22\ 133,22 = \frac{x[(1,015)^{61} - 1]}{0,015}$ $\therefore 376\ 665,55 = \frac{x[(1,015)^{61} - 1]}{0,015}$ $\therefore \frac{376\ 665,55 \times 0,015}{[(1,015)^{61} - 1]} = x$ $\therefore x = R3817,90$	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>F = 376\ 665,55</math></li> <li>✓ <math>n = 61</math></li> <li>✓ <math>\frac{0,18}{12} = 0,015</math></li> <li>✓ <math>x = R3817,90</math></li> </ul> <p style="text-align: right;">(5)</p> <p style="text-align: right;">[17]</p>

<p>2(a)</p> $100\,000 = \frac{2000 \left[ (1,08)^{n+1} - 1 \right]}{0,08}$ $\therefore \frac{100\,000 \times 0,08}{2000} = (1,08)^{n+1} - 1$ $\therefore \frac{100\,000 \times 0,08}{2000} + 1 = (1,08)^{n+1}$ $\therefore 5 = (1,08)^{n+1}$ $\therefore \log_{1,08} 5 = n + 1$ $\therefore n = 19,91237188$ $\therefore n \approx 20 \text{ half-years}$ $\therefore n \approx 10 \text{ years}$ <p>(since there are 20 half years in a ten-year period)</p>	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>\frac{0,16}{2} = 0,08</math></li> <li>✓ time period <math>= n + 1</math></li> <li>✓ <math>5 = (1,08)^{n+1}</math></li> <li>✓ <math>\log_{1,08} 5 = n + 1</math></li> <li>✓ <math>n = 19,91237188</math></li> <li>✓ <math>n \approx 10 \text{ years}</math></li> </ul> <p style="text-align: right;">(7)</p>
---	---

<p>2(b)</p> $400\,000 = \frac{17000 \left[ 1 - (1,04)^{-n} \right]}{0,04}$ $\therefore \frac{400\,000 \times 0,04}{17000} = 1 - (1,04)^{-n}$ $\therefore (1,04)^{-n} = 1 - \frac{400\,000 \times 0,04}{17000}$ $\therefore (1,04)^{-n} = 0,05882352941$ $\therefore 0,05882352941 = (1,04)^{-n}$ $\therefore \log_{1,04} 0,05882352941 = -n$ $\therefore n = 72,23768046$ <p>18 years 3 months</p>	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>\frac{0,16}{4} = 0,04</math></li> <li>✓</li> <li><math>0,0588... = (1,04)^{-n}</math></li> <li>✓</li> <li><math>\log_{0,0625} 1,015 = -n</math></li> <li>✓ <math>n = 72,23768046</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(6)</p>
--	--

[13]

## TRIGONOMETRY (REVISION)

## SECTION A: TYPICAL EXAM QUESTIONS

## QUESTION 1

(a) If  $2 \tan \theta - 3 = 0$ , then determine by means of a diagram and without using a calculator, the value of  $13 \sin^2 \theta - \frac{2}{3} \tan \theta$ , if it is given that  $180^\circ < \theta < 360^\circ$ . (6)

(b) (1) If  $\sin 18^\circ = t$ , use a diagram to determine the following in terms of  $t$ :

$$\frac{\cos^2 18^\circ \cdot \tan^2 18^\circ}{\sin 18^\circ} \quad (6)$$

(2) By using identities, verify your answer. (3)

## QUESTION 2

Simplify the following without using a calculator:

(a)  $\sin 150^\circ \cdot \cos 240^\circ \cdot \tan 315^\circ$  (5)

(b)  $\frac{\sin(180^\circ + \theta)}{\cos 360^\circ \cdot \cos(360^\circ - \theta)}$  (4)

(c)  $\sin^2 130^\circ + \sin^2 320^\circ$  (4)

## QUESTION 3

(a) If  $\tan A = p$ ,  $p > 0$  and  $A \in [0^\circ; 90^\circ]$  determine with the aid of a diagram the value of the following in terms of  $p$ .

(1)  $\sin A$  (3)

(2)  $\cos A$  (1)

(b) Simplify the following without using a calculator:

(1)  $\frac{\tan(-480^\circ) \cdot \sin 300^\circ \cdot \cos 14^\circ \cdot \sin(-135^\circ)}{\sin 104^\circ \cdot \cos 225^\circ}$  (7)




(2)  $\tan(90^\circ + x) \cdot \sin(-x - 180^\circ)$  (6)




QUESTION 4

Prove the following by using identities:  $\frac{1 + \sin \theta}{\cos \theta} + \frac{\cos \theta}{1 + \sin \theta} = \frac{2}{\cos \theta}$  (5)

SECTION B: SOLUTIONS AND HINTS

<p>1(a)</p>	$\tan \theta = \frac{3}{2} = \frac{-3}{-2}$  $(-2)^2 + (-3)^2 = r^2$ $\therefore r^2 = 13$ $\therefore r = \pm\sqrt{13}$ <p>But <math>r &gt; 0</math></p> $\therefore r = \sqrt{13}$ $13\sin^2 \theta - \frac{2}{3}\tan \theta$ $= 13\left(\frac{-3}{\sqrt{13}}\right)^2 - \frac{2}{3}\left(\frac{-3}{-2}\right)$  $= 13\left(\frac{9}{13}\right) - \frac{2}{3}\left(\frac{3}{2}\right)$ $= 9 - 1$ $= 8$	<ul style="list-style-type: none"> <li>✓ <math>\tan \theta = \frac{-3}{-2}</math></li> <li>✓ correct quadrant</li> <li>✓ <math>r = \sqrt{13}</math></li> <li>✓ correct substitution:</li> </ul> $13\left(\frac{-3}{\sqrt{13}}\right)^2 - \frac{2}{3}\left(\frac{-3}{-2}\right)$ <ul style="list-style-type: none"> <li>✓ correct answer</li> </ul> <p>(6)</p>
<p>1(b)(1)</p>	 $\sin 18^\circ = \frac{t}{1}$ $x^2 = r^2 - y^2$ $\therefore x^2 = 1^2 - t^2$ $\therefore x^2 = 1 - t^2$ $\therefore x = \sqrt{1 - t^2}$ $\frac{\cos^2 18^\circ \cdot \tan^2 18^\circ}{\sin 18^\circ}$ $= \frac{\left(\frac{\sqrt{1-t^2}}{1}\right)^2 \left(\frac{t}{\sqrt{1-t^2}}\right)^2}{t} = \frac{(1-t^2) \left(\frac{t^2}{1-t^2}\right)}{t} = t$	<ul style="list-style-type: none"> <li>✓ correct quadrant</li> <li>✓ <math>x = \sqrt{1-t^2}</math></li> <li>✓ <math>\frac{\sqrt{1-t^2}}{1}</math></li> <li>✓ <math>\frac{t}{\sqrt{1-t^2}}</math></li> <li>✓ squaring</li> <li>✓ answer</li> </ul> <p>(6)</p>

1(b)(2)	$\frac{\cos^2 18^\circ \cdot \tan^2 18^\circ}{\sin 18^\circ}$ $= \frac{\cos^2 18^\circ \cdot \frac{\sin^2 18^\circ}{\cos^2 18^\circ}}{\sin 18^\circ}$ $= \frac{\sin^2 18^\circ}{\sin 18^\circ}$ $= \sin 18^\circ$ $= t$	<ul style="list-style-type: none"> <li>✓ <math>\frac{\sin^2 18^\circ}{\cos^2 18^\circ}</math></li> <li>✓ <math>\sin 18^\circ</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(3)</p>
2(a)	$\sin 150^\circ \cdot \cos 240^\circ \cdot \tan 315^\circ$ $= (\sin 30^\circ)(-\cos 60^\circ)(-\tan 45^\circ)$ $= \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)(-1)$ $= \frac{1}{4}$	<ul style="list-style-type: none"> <li>✓ <math>\sin 30^\circ</math></li> <li>✓ <math>\cos 360^\circ = 1</math></li> <li>✓ <math>-\tan 45^\circ</math></li> <li>✓ evaluating special angles</li> <li>✓ answer</li> </ul> <p style="text-align: right;">(5)</p>
2(b)	$\frac{\sin(180^\circ + \theta)}{\cos 360^\circ \cdot \cos(360^\circ - \theta)}$ $= \frac{-\sin \theta}{(1)(\cos \theta)}$ $= -\tan \theta$	<ul style="list-style-type: none"> <li>✓ <math>-\sin \theta</math></li> <li>✓ <math>-\cos 60^\circ</math></li> <li>✓ <math>\cos \theta</math></li> <li>✓ <math>-\tan \theta</math></li> </ul> <p style="text-align: right;">(4)</p>
2(c)	$\sin^2 130^\circ + \sin^2 320^\circ$ $= \sin^2 50^\circ + \sin^2 40^\circ$ $= \sin^2 50^\circ + \cos^2 50^\circ$ $= 1$	<ul style="list-style-type: none"> <li>✓ <math>\sin^2 50^\circ</math></li> <li>✓ <math>\sin^2 40^\circ</math></li> <li>✓ <math>\cos^2 50^\circ</math></li> <li>✓ 1</li> </ul> <p style="text-align: right;">(4)</p>

<p>3(a)(1)</p>	$\tan A = \frac{p}{1}$ $r^2 = x^2 + y^2$ $r^2 = 1^2 + p^2$ $r^2 = p^2 + 1$ $r = \sqrt{p^2 + 1}$ $\sin A = \frac{p}{\sqrt{p^2 + 1}}$ 	<ul style="list-style-type: none"> <li>✓ correct quadrant</li> <li>✓ <math>r = \sqrt{1 + p^2}</math></li> <li>✓ <math>\sin A = \frac{p}{\sqrt{p^2 + 1}}</math></li> </ul> <p style="text-align: right;">(3)</p>
<p>3(a)(2)</p>	$\cos A = \frac{1}{\sqrt{p^2 + 1}}$	<ul style="list-style-type: none"> <li>✓ <math>\cos A = \frac{1}{\sqrt{p^2 + 1}}</math></li> </ul> <p style="text-align: right;">(1)</p>
<p>3(b)(1)</p>	$\frac{\tan(-480^\circ) \cdot \sin 300^\circ \cdot \cos 14^\circ \cdot \sin(-135^\circ)}{\sin 104^\circ \cdot \cos 225^\circ}$ $= \frac{(-\tan 480^\circ)(-\sin 60^\circ)(\cos 14^\circ)(-\sin 135^\circ)}{(\sin 76^\circ)(-\cos 45^\circ)}$ $= \frac{(-\tan 120^\circ)(-\sin 60^\circ)(\cos 14^\circ)(-\sin 45^\circ)}{(\sin 76^\circ)(-\cos 45^\circ)}$ $= \frac{(-(-\tan 60^\circ))(-\sin 60^\circ)(\cos 14^\circ)(-\sin 45^\circ)}{(\sin(90^\circ - 14^\circ))(-\cos 45^\circ)}$ $= \frac{(\tan 60^\circ)(-\sin 60^\circ)(\cos 14^\circ)(-\sin 45^\circ)}{(\cos 14^\circ)(-\cos 45^\circ)}$ $= \frac{(\sqrt{3}) \cdot \left(-\frac{\sqrt{3}}{2}\right) \cdot \left(-\frac{\sqrt{2}}{2}\right)}{\left(-\frac{\sqrt{2}}{2}\right)} = -\frac{3}{2}$	<ul style="list-style-type: none"> <li>✓ <math>\tan 60^\circ</math></li> <li>✓ <math>-\sin 60^\circ</math></li> <li>✓ <math>-\sin 45^\circ</math></li> <li>✓ <math>\sin(90^\circ - 14^\circ) = \cos 14^\circ</math></li> <li>✓ <math>-\cos 45^\circ</math></li> <li>✓ evaluating special angles</li> <li>✓ answer</li> </ul> <p style="text-align: right;">(7)</p>
<p>3(b)(2)</p>	$\tan(90^\circ + x) \cdot \sin(-x - 180^\circ)$ $= \frac{\sin(90^\circ + x)}{\cos(90^\circ + x)} \sin[-(x + 180^\circ)]$ $= \frac{\cos x}{-\sin x} \times -\sin(180^\circ + x)$ $= \frac{\cos x}{-\sin x} \times -(-\sin x)$ $= \frac{\cos x}{-\sin x} \times \sin x$ $= -\cos x$	<ul style="list-style-type: none"> <li>✓ <math>\frac{\sin(90^\circ + x)}{\cos(90^\circ + x)}</math></li> <li>✓ <math>-\sin(180^\circ + x)</math></li> <li>✓✓ <math>\frac{\cos x}{-\sin x}</math></li> <li>✓ <math>\sin x</math></li> <li>✓ <math>-\cos x</math></li> </ul> <p style="text-align: right;">(6)</p>

4	$\begin{aligned} \text{LHS} &= \frac{1 + \sin \theta}{\cos \theta} + \frac{\cos \theta}{1 + \sin \theta} \\ &= \frac{(1 + \sin \theta)^2 + \cos^2 \theta}{\cos \theta(1 + \sin \theta)} \\ &= \frac{1 + 2 \sin \theta + \sin^2 \theta + \cos^2 \theta}{\cos \theta(1 + \sin \theta)} \\ &= \frac{1 + 2 \sin \theta + 1}{\cos \theta(1 + \sin \theta)} \\ &= \frac{2 + 2 \sin \theta}{\cos \theta(1 + \sin \theta)} = \frac{2(1 + \sin \theta)}{\cos \theta(1 + \sin \theta)} = \frac{2}{\cos \theta} = \text{RHS} \end{aligned}$	$\begin{aligned} &\checkmark \cos \theta(1 + \sin \theta) \\ &\checkmark \\ &1 + 2 \sin \theta + \sin^2 \theta + \cos^2 \theta \\ &\checkmark \sin^2 \theta + \cos^2 \theta = 1 \\ &\checkmark 2(1 + \sin \theta) \\ &\checkmark \frac{2}{\cos \theta} \end{aligned} \quad (5)$
---	--	---

**SECTION C: HOMEWORK****QUESTION 1**

- (a) If  $\frac{5 \sin A}{2} = \sqrt{6}$  and  $A \in [90^\circ; 360^\circ]$  calculate without the use of a calculator and with the aid of a diagram the value of  $5 \tan A \cdot \cos A$ .
- (b) If  $\sin 32^\circ = m$ , use a diagram to express the following in terms of  $m$ :
- (1)  $\sin 328^\circ$  (4)
  - (2)  $\cos 58^\circ$  (2)
  - (3)  $\tan 212^\circ$  (2)

**QUESTION 2**

Simplify the following without using a calculator:

- (a)  $\frac{\cos 210^\circ \cdot \tan^2 315^\circ}{\sin 300^\circ \cdot \cos 120^\circ}$  (5)
- (b)  $\frac{\sin^2(180^\circ + \theta) \cdot \sin(360^\circ - \theta)}{\cos^2(90^\circ - \theta) \cdot \cos(90^\circ + \theta)}$  (5)
- (c)  $\cos^2(360^\circ - x) - \sin(180^\circ - x) \cos(90^\circ + x) - \cos^2(180^\circ + x)$  (5)

**QUESTION 3**

(a) Simplify:

$$\frac{\tan(180^\circ + x)\cos(360^\circ - x)}{\sin(180^\circ - x)\cos(90^\circ + x) + \cos(540^\circ + x)\cos(-x)} \quad (9)$$

(b) If  $\cos 26^\circ = p$ , express the following in terms of  $p$ :



$$\frac{\cos(-64^\circ)\tan(-244^\circ)\sin^2 334^\circ}{\cos 566^\circ} \quad (8)$$



**QUESTION 4**

Prove that:

$$\cos^2 x \left[ \frac{1}{\sin x - 1} + \frac{1}{\sin x + 1} \right] = 2 \quad (5)$$

**SECTION D: SOLUTIONS TO HOMEWORK**

<p>1(a)</p>	<p><math>\sin A = \frac{2\sqrt{6}}{5}</math></p> <p><math>x^2 = r^2 - y^2</math></p> <p><math>x^2 = 5^2 - (2\sqrt{6})^2</math></p> <p><math>x^2 = 1</math></p> <p><math>x = \pm 1</math></p> <p><math>\therefore x = -1</math></p> <p><math>\therefore 5 \tan A \cdot \cos A = 5 \left( \frac{2\sqrt{6}}{-1} \right) \cdot \left( \frac{-1}{5} \right) = 2\sqrt{6}</math></p>	<p></p> <p></p> <p>✓ <math>\sin A = \frac{2\sqrt{6}}{5}</math></p> <p>✓ correct quadrant</p> <p>✓ <math>x = -1</math></p> <p>✓✓ correct substitution:  <math>5 \left( \frac{2\sqrt{6}}{-1} \right) \cdot \left( \frac{-1}{5} \right)</math></p> <p>✓ correct answer</p> <p>(6)</p>
-------------	---	---

<p>1(b)(1)</p>	$\sin 32^\circ = \frac{m}{1}$ $x^2 + m^2 = 1$ $\therefore x = \sqrt{1 - m^2}$ $\sin 328^\circ = \sin 32^\circ = m$		<ul style="list-style-type: none"> <li>✓ correct quadrant</li> <li>✓ <math>x = \sqrt{1 - m^2}</math></li> <li>✓ <math>\sin 328^\circ = \sin 32^\circ</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p>
<p>1(b)(2)</p>			<ul style="list-style-type: none"> <li>✓ <math>\sin 32^\circ</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(2)</p>
<p>1(b)(3)</p>	$\tan 212^\circ$ $= \tan 32^\circ$ $= \frac{m}{\sqrt{1 - m^2}}$		<ul style="list-style-type: none"> <li>✓ <math>\tan 32^\circ</math></li> <li>✓ <math>\frac{m}{\sqrt{1 - m^2}}</math></li> </ul> <p style="text-align: right;">(2)</p>
<p>2(a)</p>	$\frac{\cos 210^\circ \cdot \tan^2 315^\circ}{\sin 300^\circ \cdot \cos 120^\circ}$ $= \frac{(-\cos 30^\circ)(-\tan 45^\circ)^2}{(-\sin 60^\circ)(-\cos 60^\circ)}$ $= \frac{\left(-\frac{\sqrt{3}}{2}\right)(-1)^2}{\left(-\frac{\sqrt{3}}{2}\right)\left(-\frac{1}{2}\right)}$ $= -2$		<ul style="list-style-type: none"> <li>✓ <math>-\cos 30^\circ</math></li> <li>✓ <math>(-\tan 45^\circ)^2</math></li> <li>✓ <math>-\sin 60^\circ</math></li> <li>✓ <math>-\cos 60^\circ</math></li> <li>✓ evaluating special angles</li> <li>✓ answer</li> </ul> <p style="text-align: right;">(6)</p>
<p>2(b)</p>	$\frac{\sin^2(180^\circ + \theta) \cdot \sin(360^\circ - \theta)}{\cos^2(90^\circ - \theta) \cdot \cos(90^\circ + \theta)}$ $= \frac{(\sin^2 \theta)(-\sin \theta)}{(\sin^2 \theta)(-\sin \theta)}$ $= 1$		<p>Numerator:</p> <ul style="list-style-type: none"> <li>✓ <math>\sin^2 \theta</math></li> <li>✓ <math>-\sin \theta</math></li> </ul> <p>Denominator:</p> <ul style="list-style-type: none"> <li>✓ <math>\sin^2 \theta</math></li> <li>✓ <math>-\sin \theta</math></li> <li>✓ 1</li> </ul> <p style="text-align: right;">(5)</p>
<p>2(c)</p>	$\cos^2(360^\circ - x) - \sin(180^\circ - x)\cos(90^\circ + x) - \cos^2(180^\circ + x)$ $= \cos^2 x - (\sin x)(-\sin x) - \cos^2 x$ $= \sin^2 x$		<ul style="list-style-type: none"> <li>✓✓✓✓ reductions</li> <li>✓ <math>\sin^2 x</math></li> </ul> <p style="text-align: right;">(5)</p>

<p>3(a)</p>	$\frac{\tan(180^\circ + x) \cos(360^\circ - x)}{\sin(180^\circ - x) \cos(90^\circ + x) + \cos(540^\circ + x) \cos(-x)}$ $= \frac{(\tan x)(\cos x)}{(\sin x)(-\sin x) - (\cos x)(\cos x)}$ $= \frac{\frac{\sin x}{\cos x} \cos x}{-\sin^2 x - \cos^2 x}$ $= \frac{\sin x}{-(\sin^2 x + \cos^2 x)}$ $= -\sin x$	<p>Numerator:</p> <ul style="list-style-type: none"> <li>✓✓ <math>(\tan x)(\cos x)</math></li> <li>✓ <math>\frac{\sin x}{\cos x}</math></li> </ul> <p>Denominator:</p> <ul style="list-style-type: none"> <li>✓✓✓✓</li> <li><math>(\sin x)(-\sin x) - (\cos x)(\cos x)</math></li> <li>✓ <math>-(\sin^2 x + \cos^2 x)</math></li> <li>✓ <math>-\sin x</math></li> </ul> <p style="text-align: right;">(9)</p>
<p>3(b)</p>	$\frac{\cos(-64^\circ) \tan(-244^\circ) \sin^2 334^\circ}{\cos 566^\circ}$ $= \frac{(\cos 64^\circ)(-\tan 244^\circ)(-\sin 26^\circ)^2}{\cos 206^\circ}$ $= \frac{(\cos 64^\circ)(-\tan 64^\circ) \sin^2 26^\circ}{-\cos 26^\circ}$ $= \frac{(\cos 64^\circ) \left( \frac{-\sin 64^\circ}{\cos 64^\circ} \right) \sin^2 26^\circ}{-\sin 64^\circ}$ $= \sin^2 26^\circ$ $= 1 - \cos^2 26^\circ$ $= 1 - p^2$	<ul style="list-style-type: none"> <li>✓ <math>\cos 64^\circ</math></li> <li>✓ <math>-\tan 64^\circ</math></li> <li>✓ <math>\sin^2 26^\circ</math></li> <li>✓ <math>-\cos 26^\circ</math></li> <li>✓ <math>\frac{-\sin 64^\circ}{\cos 64^\circ}</math></li> <li>✓ <math>-\sin 64^\circ</math></li> <li>✓ <math>1 - \cos^2 26^\circ</math></li> <li>✓ <math>1 - p^2</math></li> </ul> <p style="text-align: right;">(8)</p>
<p>4</p>	$\cos^2 x \left[ \frac{1}{\sin x - 1} + \frac{1}{\sin x + 1} \right]$ $= \cos^2 x \left[ \frac{(1 + \sin x) + (1 - \sin x)}{(1 + \sin x)(1 - \sin x)} \right]$ $= \cos^2 x \left[ \frac{2}{1 - \sin^2 x} \right]$ $= \cos^2 x \left[ \frac{2}{\cos^2 x} \right]$ $= 2$	<ul style="list-style-type: none"> <li>✓ <math>(1 + \sin x) + (1 - \sin x)</math></li> <li>✓ <math>(1 + \sin x)(1 - \sin x)</math></li> <li>✓ <math>\frac{2}{1 - \sin^2 x}</math></li> <li>✓ <math>\frac{2}{\cos^2 x}</math></li> <li>✓ 2</li> </ul> <p style="text-align: right;">(5)</p>

## TRIGONOMETRY

**Teacher Note:** Trigonometry is an extremely important and large part of Paper 2. Ensure that learners master all the basic rules and definitions and be able to apply these rules in many different types of questions. In this session learners will concentrate on Grade 12 Trigonometry which involves compound and double angles. These Grade 12 concepts will be integrated with the trigonometry studied in Grade 11. Before attempting the typical exam questions, learners must familiarise themselves with the basics in Section B of the learner notes.

## LESSON OVERVIEW

- |    |                                   |            |
|----|-----------------------------------|------------|
| 1. | Introduction session:             | 10 minutes |
|    | Discuss Homework from Session 5   | 10 minutes |
| 2. | Typical exam questions:           |            |
|    | Question 1                        | 15 minutes |
|    | Question 2:                       | 10 minutes |
|    | Question 3:                       | 5 minutes  |
| 3. | Discussion of solutions:          | 35 minutes |
| 4. | Discuss Homework for this session | 5 minutes  |

## SECTION A: TYPICAL EXAM QUESTIONS

## QUESTION 1

Simplify the following without using a calculator:

$$(a) \frac{\tan(-60^\circ) \cos(-156^\circ) \cos 294^\circ}{\sin 492^\circ} \quad (7)$$

$$(b) \frac{\cos^2 375^\circ - \cos^2(-75^\circ)}{\sin(-50^\circ) \sin 230^\circ - \sin 40^\circ \cos 310^\circ} \quad (7)$$

[14]

## QUESTION 2

$$(a) \text{ Show that } \cos(60^\circ + \theta) - \cos(60^\circ - \theta) = -\sqrt{3} \sin \theta \quad (5)$$

$$(b) \text{ Hence, evaluate } \cos 105^\circ + \cos 15^\circ \text{ without using a calculator.} \quad (5)$$

[10]



QUESTION 3

Rewrite  $\cos 3\theta$  in terms of  $\cos \theta$ .

[6]

**SECTION B: SOLUTIONS AND HINTS**

<p>1(a)</p>	$\frac{\tan(-60^\circ)\cos(-156^\circ)\cos 294^\circ}{\sin 492^\circ}$ $= \frac{(-\tan 60^\circ)(\cos 156^\circ)(-\cos 66^\circ)}{(\sin 132^\circ)}$ $= \frac{(-\sqrt{3})(-\cos 24^\circ)(-\sin 24^\circ)}{(\sin 48^\circ)}$ $= \frac{(-\sqrt{3})(-\cos 24^\circ)(-\sin 24^\circ)}{2\sin 24^\circ\cos 24^\circ}$ $= -\frac{\sqrt{3}}{2}$	<ul style="list-style-type: none"> <li>✓ <math>(-\tan 60^\circ)(\cos 156^\circ)</math></li> <li>✓ <math>-\cos 66^\circ</math></li> <li>✓ <math>\sin 48^\circ</math></li> <li>✓ <math>-\sqrt{3}</math></li> <li>✓ <math>-\sin 24^\circ</math></li> <li>✓ <math>2\sin 24^\circ\cos 24^\circ</math></li> <li>✓ <math>\frac{\sqrt{3}}{2}</math></li> </ul> <p>(7)</p>
<p>1(b)</p>	$\frac{\cos^2 375^\circ - \cos^2(-75^\circ)}{\sin(-50^\circ)\sin 230^\circ + \sin 40^\circ\cos 310^\circ}$ $= \frac{\cos^2 15^\circ - \cos^2 75^\circ}{(-\sin 50^\circ)(-\sin 50^\circ) + (\sin 40^\circ)(\cos 50^\circ)}$ $= \frac{\cos^2 15^\circ - \sin^2 15^\circ}{\sin^2 50^\circ + (\cos 50^\circ)(\cos 50^\circ)}$ $= \frac{\cos^2 15^\circ - \sin^2 15^\circ}{\sin^2 50^\circ + \cos^2 50^\circ}$ $= \frac{\cos 2(15^\circ)}{1}$ $= \cos 30^\circ$ $= \frac{\sqrt{3}}{2}$	<ul style="list-style-type: none"> <li>✓ <math>\cos^2 15^\circ</math></li> <li>✓ <math>\cos^2 75^\circ</math></li> <li>✓ <math>\sin^2 50^\circ</math></li> <li>✓ <math>\cos^2 50^\circ</math></li> <li>✓ <math>\cos 30^\circ</math></li> <li>✓ 1</li> <li>✓ <math>\frac{\sqrt{3}}{2}</math></li> </ul> <p>(7)</p> <p>[14]</p>
<p>2(a)</p>	$\cos(60^\circ + \theta) - \cos(60^\circ - \theta)$ $= \cos 60^\circ \cdot \cos \theta - \sin 60^\circ \cdot \sin \theta - (\cos 60^\circ \cdot \cos \theta + \sin 60^\circ \cdot \sin \theta)$ $= \left(\frac{1}{2}\right) \cdot \cos \theta - \left(\frac{\sqrt{3}}{2}\right) \cdot \sin \theta - \left(\frac{1}{2}\right) \cdot \cos \theta - \left(\frac{\sqrt{3}}{2}\right) \cdot \sin \theta$ $= -\sqrt{3} \sin \theta$	<ul style="list-style-type: none"> <li>✓ <math>\cos 60^\circ \cdot \cos \theta - \sin 60^\circ \cdot \sin \theta</math></li> <li>✓ <math>\cos 60^\circ \cdot \cos \theta + \sin 60^\circ \cdot \sin \theta</math></li> <li>✓ <math>\frac{\sqrt{3}}{2}</math></li> <li>✓ <math>\frac{1}{2}</math></li> <li>✓ <math>-\sqrt{3} \sin \theta</math></li> </ul> <p>(5)</p>

2(b)	$\begin{aligned} & \cos 105^\circ + \cos 15^\circ \\ &= \cos(60^\circ + 45^\circ) - \cos(60^\circ - 45^\circ) \\ &= -\sqrt{3} \sin 45^\circ \\ &= -\sqrt{3} \left( \frac{\sqrt{2}}{2} \right) \\ &= \frac{-\sqrt{6}}{2} \end{aligned}$	$\begin{aligned} & \checkmark \cos(60^\circ + 45^\circ) \\ & \checkmark \cos(60^\circ - 45^\circ) \\ & \checkmark -\sqrt{3} \sin 45^\circ \\ & \checkmark \frac{\sqrt{2}}{2} \\ & \checkmark \frac{-\sqrt{6}}{2} \end{aligned} \quad (5)$ <p style="text-align: right;"><b>[10]</b></p>
3	$\begin{aligned} & \cos 3\theta \\ &= \cos(2\theta + \theta) \\ &= \cos 2\theta \cdot \cos \theta - \sin 2\theta \cdot \sin \theta \\ &= (2\cos^2 \theta - 1) \cdot \cos \theta - (2\sin \theta \cos \theta) \cdot \sin \theta \\ &= 2\cos^3 \theta - \cos \theta - 2\sin^2 \theta \cdot \cos \theta \\ &= 2\cos^3 \theta - \cos \theta - 2(1 - \cos^2 \theta) \cos \theta \\ &= 2\cos^3 \theta - \cos \theta - 2(\cos \theta - \cos^3 \theta) \\ &= 2\cos^3 \theta - \cos \theta - 2\cos \theta + 2\cos^3 \theta \\ &= 4\cos^3 \theta - 3\cos \theta \end{aligned}$	$\begin{aligned} & \checkmark \cos(2\theta + \theta) \\ & \checkmark \cos 2\theta \cdot \cos \theta - \sin 2\theta \cdot \sin \theta \\ & \checkmark 2\cos^2 \theta - 1 \\ & \checkmark 2\sin \theta \cos \theta \\ & \checkmark 1 - \cos^2 \theta \\ & \checkmark 4\cos^3 \theta - 3\cos \theta \end{aligned}$ <p style="text-align: right;"><b>[6]</b></p>



### SECTION C: HOMEWORK

#### QUESTION 1

Determine the value of the following without using a calculator.

(a) 
$$\frac{\sin 34^\circ \cos 10^\circ - \cos 34^\circ \sin 10^\circ}{\sin 12^\circ \cos 12^\circ} \quad (3)$$

(b)  $\sin(-285^\circ) \quad (5)$

(c) 
$$\frac{\cos^2 15^\circ - \sin 15^\circ \cos 75^\circ}{\cos^2 15^\circ + \sin 15^\circ \cos 15^\circ \tan 15^\circ} \quad (6)$$

[14]

**QUESTION 2**

(a) Prove that  $\sin(45^\circ + \theta) \cdot \sin(45^\circ - \theta) = \frac{1}{2} \cos 2\theta$  (5)

(b) Hence determine the value of  $\sin 75^\circ \cdot \sin 15^\circ$  (3)  
 [8]

**QUESTION 3**

Prove that:  $\sin 4\theta = 4\sin \theta \cdot \cos \theta - 8\sin^3 \theta \cdot \cos \theta$  [4]

**SOLUTIONS TO HOMEWORK: TRIGONOMETRY (A)**

1(a)	$\frac{\sin 34^\circ \cos 10^\circ - \cos 34^\circ \sin 10^\circ}{\sin 12^\circ \cos 12^\circ}$ $= \frac{\sin(34^\circ - 10^\circ)}{\sin 12^\circ \cos 12^\circ}$ $= \frac{\sin 24^\circ}{\sin 12^\circ \cos 12^\circ}$ $= \frac{2 \sin 12^\circ \cos 12^\circ}{\sin 12^\circ \cos 12^\circ}$ $= 2$	<ul style="list-style-type: none"> <li>✓ <math>\sin 24^\circ</math></li> <li>✓ <math>2 \sin 12^\circ \cos 12^\circ</math></li> <li>✓ 2</li> </ul> <p style="text-align: right;">(3)</p>
1(b)	$\sin(-285^\circ)$ $= -\sin 285^\circ$ $= -\sin(360^\circ - 75^\circ)$ $= -(-\sin 75^\circ)$ $= \sin 75^\circ$ $= \sin(45^\circ + 30^\circ)$ $= \sin 45^\circ \cos 30^\circ + \cos 45^\circ \sin 30^\circ$ $= \left(\frac{\sqrt{2}}{2}\right)\left(\frac{\sqrt{3}}{2}\right) + \left(\frac{\sqrt{2}}{2}\right)\left(\frac{1}{2}\right)$ $= \frac{\sqrt{6} + \sqrt{2}}{4}$	<ul style="list-style-type: none"> <li>✓ <math>-\sin 285^\circ</math></li> <li>✓ <math>\sin 75^\circ</math></li> <li>✓ <math>\sin 45^\circ \cos 30^\circ + \cos 45^\circ \sin 30^\circ</math></li> <li>✓ <math>\left(\frac{\sqrt{2}}{2}\right)\left(\frac{\sqrt{3}}{2}\right) + \left(\frac{\sqrt{2}}{2}\right)\left(\frac{1}{2}\right)</math></li> <li>✓ <math>\frac{\sqrt{6} + \sqrt{2}}{4}</math></li> </ul> <p style="text-align: right;">(5)</p>

## MATHEMATICS

## GRADE 12

## SESSION 6

## (TEACHER NOTES)

1(c)	$\frac{\cos^2 15^\circ - \sin 15^\circ \cos 75^\circ}{\cos^2 15^\circ + \sin 15^\circ \cos 15^\circ \tan 15^\circ}$ $= \frac{\cos^2 15^\circ - \sin 15^\circ \cos(90^\circ - 15^\circ)}{\cos^2 15^\circ + \sin 15^\circ \cos 15^\circ \left(\frac{\sin 15^\circ}{\cos 15^\circ}\right)}$ $= \frac{\cos^2 15^\circ - \sin 15^\circ \cdot \sin 15^\circ}{\cos^2 15^\circ + \sin^2 15^\circ}$ $= \frac{\cos^2 15^\circ - \sin^2 15^\circ}{1}$ $= \cos^2 15^\circ - \sin^2 15^\circ$ $= \cos 2(15^\circ) = \cos 30^\circ = \frac{\sqrt{3}}{2}$	<ul style="list-style-type: none"> <li>✓ <math>\cos^2 15^\circ - \sin 15^\circ \cdot \sin 15^\circ</math></li> <li>✓ <math>\frac{\sin 15^\circ}{\cos 15^\circ}</math></li> <li>✓ <math>\cos^2 15^\circ + \sin^2 15^\circ</math></li> <li>✓ 1</li> <li>✓ <math>\cos 30^\circ</math></li> <li>✓ <math>\frac{\sqrt{3}}{2}</math></li> </ul> <p style="text-align: right;">(6)</p> <p style="text-align: right;"><b>[14]</b></p>
2(a)	$\sin(45^\circ + \theta) \cdot \sin(45^\circ - \theta)$ $= [\sin 45^\circ \cos \theta + \cos 45^\circ \sin \theta][\sin 45^\circ \cos \theta - \cos 45^\circ \sin \theta]$ $= \left[ \frac{\sqrt{2}}{2} \cos \theta + \frac{\sqrt{2}}{2} \sin \theta \right] \left[ \frac{\sqrt{2}}{2} \cos \theta - \frac{\sqrt{2}}{2} \sin \theta \right]$ $= \left[ \frac{\sqrt{2}}{2} (\cos \theta + \sin \theta) \right] \left[ \frac{\sqrt{2}}{2} (\cos \theta - \sin \theta) \right]$ $= \frac{2}{4} (\cos \theta + \sin \theta)(\cos \theta - \sin \theta)$ $= \frac{1}{2} (\cos^2 \theta - \sin^2 \theta) = \frac{1}{2} \cos 2\theta$	<ul style="list-style-type: none"> <li>✓ expansion of <math>\sin(45^\circ + \theta)</math></li> <li>✓ expansion of <math>\sin(45^\circ - \theta)</math></li> <li>✓ <math>\sin 45^\circ = \cos 45^\circ = \frac{\sqrt{2}}{2}</math></li> <li>✓ <math>(\cos^2 \theta - \sin^2 \theta)</math></li> <li>✓ <math>\frac{1}{2} \cos 2\theta</math></li> </ul> <p style="text-align: right;">(5)</p>
2(b)	$\sin 75^\circ \cdot \sin 15^\circ$ $= \sin(45^\circ + 30^\circ) \cdot \sin(45^\circ - 30^\circ)$ $= \frac{1}{2} \cos 2(30^\circ) = \frac{1}{2} \cos 60^\circ = \frac{1}{2} \left( \frac{1}{2} \right) = \frac{1}{4}$	<ul style="list-style-type: none"> <li>✓ <math>45^\circ + 30^\circ</math>; <math>45^\circ - 30^\circ</math></li> <li>✓ <math>\frac{1}{2} \cos 60^\circ</math></li> <li>✓ <math>\frac{1}{4}</math></li> </ul> <p style="text-align: right;">(3)</p> <p style="text-align: right;"><b>[8]</b></p>
3	$\sin 4\theta$ $= \sin 2(2\theta)$ $= 2 \sin 2\theta \cdot \cos 2\theta$ $= 2(2 \sin \theta \cdot \cos \theta)(1 - 2 \sin^2 \theta)$ $= 4 \sin \theta \cdot \cos \theta - 8 \sin^3 \theta \cdot \cos \theta$	<ul style="list-style-type: none"> <li>✓ <math>2 \sin 2\theta \cdot \cos 2\theta</math></li> <li>✓ <math>2 \sin \theta \cdot \cos \theta</math></li> <li>✓ <math>1 - 2 \sin^2 \theta</math></li> <li>✓ <math>4 \sin \theta \cdot \cos \theta - 8 \sin^3 \theta \cdot \cos \theta</math></li> </ul> <p style="text-align: right;"><b>[4]</b></p>

**TRIGONOMETRY**

**! Teacher Note:** Trigonometry is an extremely important and large part of Paper 2. Ensure that learners master all the basic rules and definitions and be able to apply these rules in many different types of questions. In this session learners will concentrate on Grade 12 Trigonometry which involves compound and double angles. These Grade 12 concepts will be integrated with the trigonometry studied in Grade 11. Before attempting the typical exam questions, learners must familiarise themselves with the basics in Section B of the learner notes.

**LESSON OVERVIEW**

1. Introduction session: 5 minutes  
 Homework Discussion 10 minutes
2. Typical exam questions:  
 Question 1: 15 minutes  
 Question 2: 10 minutes  
 Question 3: 10 minutes
3. Discussion of solutions: 35 minutes
4. Discuss Homework 5 minutes



**SECTION A: TYPICAL EXAM QUESTIONS**

**QUESTION 1**

Prove that:

(a) 
$$\frac{1 - \cos 2x - \sin x}{\sin 2x - \cos x} = \tan x \quad (6)$$

(b) 
$$\frac{\sin \theta + \sin 2\theta}{1 + \cos \theta + \cos 2\theta} = \tan \theta \quad (6)$$

(c) 
$$\frac{\cos 2A}{1 + \sin 2A} = \frac{\cos A - \sin A}{\cos A + \sin A} \quad (5)$$

[17]

**QUESTION 2**

It is known that  $13\sin \alpha - 5 = 0$  and  $\tan \beta = -\frac{3}{4}$  where  $\alpha \in [90^\circ ; 270^\circ]$  and  $\beta \in [90^\circ ; 270^\circ]$ .

Determine, without using a calculator, the values of the following:

(a)  $\cos \alpha \quad (5)$

(b)  $\cos(\alpha + \beta) \quad (6)$

[11]

## QUESTION 3



If  $\sin 18^\circ = t$  determine the following in terms of  $t$ .



(a)  $\cos 18^\circ$  (4)

(b)  $\sin 78^\circ$  (5)

## SECTION B: SOLUTIONS AND HINTS

1(a)	$\frac{1 - \cos 2x - \sin x}{\sin 2x - \cos x}$ $= \frac{1 - (1 - 2\sin^2 x) - \sin x}{2\sin x \cos x - \cos x}$ $= \frac{1 - 1 + 2\sin^2 x - \sin x}{2\sin x \cos x - \cos x}$ $= \frac{2\sin^2 x - \sin x}{2\sin x \cos x - \cos x}$ $= \frac{\sin x(2\sin x - 1)}{\cos x(2\sin x - 1)}$ $= \frac{\sin x}{\cos x} = \tan x$	<ul style="list-style-type: none"> <li>✓ <math>1 - 2\sin^2 x</math></li> <li>✓ <math>2\sin x \cos x</math></li> <li>✓ <math>\sin x(2\sin x - 1)</math></li> <li>✓ <math>\cos x(2\sin x - 1)</math></li> <li>✓ <math>\frac{\sin x}{\cos x}</math></li> <li>✓ <math>\tan x</math></li> </ul> <p style="text-align: right;">(6)</p>
1(b)	$\frac{\sin \theta + \sin 2\theta}{1 + \cos \theta + \cos 2\theta}$ $= \frac{\sin \theta + 2\sin \theta \cos \theta}{1 + \cos \theta + (2\cos^2 \theta - 1)}$ $= \frac{\sin \theta(1 + 2\cos \theta)}{\cos \theta + 2\cos^2 \theta}$ $= \frac{\sin \theta(1 + 2\cos \theta)}{\cos \theta(1 + 2\cos \theta)}$ $= \frac{\sin \theta}{\cos \theta}$ $= \tan \theta$	<ul style="list-style-type: none"> <li>✓ <math>2\sin \theta \cos \theta</math></li> <li>✓ <math>2\cos^2 \theta - 1</math></li> <li>✓ <math>\sin \theta(1 + 2\cos \theta)</math></li> <li>✓ <math>\cos \theta(1 + 2\cos \theta)</math></li> <li>✓ <math>\frac{\sin \theta}{\cos \theta}</math></li> <li>✓ <math>\tan \theta</math></li> </ul> <p style="text-align: right;">(6)</p>

<p>1(c)</p>	$\frac{\cos 2A}{1 + \sin 2A}$ $= \frac{\cos^2 A - \sin^2 A}{1 + 2 \sin A \cos A}$ $= \frac{(\cos A + \sin A)(\cos A - \sin A)}{\sin^2 A + \cos^2 A + 2 \sin A \cos A}$ $= \frac{(\cos A + \sin A)(\cos A - \sin A)}{\sin^2 A + 2 \sin A \cos A + \cos^2 A}$ $= \frac{(\cos A + \sin A)(\cos A - \sin A)}{(\sin A + \cos A)(\sin A + \cos A)}$ $= \frac{\cos A - \sin A}{\cos A + \sin A}$	<ul style="list-style-type: none"> <li>✓ <math>\cos^2 A - \sin^2 A</math></li> <li>✓ <math>2 \sin A \cos A</math></li> <li>✓ <math>(\cos A + \sin A)(\cos A - \sin A)</math></li> <li>✓ <math>(\sin A + \cos A)(\sin A + \cos A)</math></li> <li>✓ <math>\frac{\cos A - \sin A}{\cos A + \sin A}</math></li> </ul> <p style="text-align: right;">(5)</p> <p style="text-align: right;">[17]</p>
<p>2(a)</p>	<div style="text-align: center;">  </div> <p style="text-align: center;">  </p> $y_\alpha = 5 \quad r_\alpha = 13$ $x^2 + (5)^2 = (13)^2$ $\therefore x^2 = 144$ $\therefore x_\alpha = -12$ $\therefore \cos \alpha = -\frac{12}{13}$	<ul style="list-style-type: none"> <li>✓ <math>\sin \alpha = \frac{5}{13}</math></li> <li>✓ diagram</li> <li>✓ Pythagoras</li> <li>✓ <math>x_\alpha = -12</math></li> <li>✓ <math>\cos \alpha = -\frac{12}{13}</math></li> </ul> <p style="text-align: right;">(5)</p>

<p>2(b)</p>	$\tan \beta = -\frac{3}{4}$ $y_{\beta} = 3 \quad x_{\beta} = -4$ $r = 5$  $\cos(\alpha + \beta)$ $= \cos \alpha \cdot \cos \beta - \sin \alpha \cdot \sin \beta$ $= \left(-\frac{12}{13}\right) \cdot \left(-\frac{4}{5}\right) - \left(\frac{5}{13}\right) \cdot \left(\frac{3}{5}\right)$ $= \frac{48 - 15}{65} = \frac{33}{65}$	<ul style="list-style-type: none"> <li>✓ diagram</li> <li>✓ Pythagoras</li> <li>✓ <math>r = 5</math></li> <li>✓ <math>\cos \alpha \cdot \cos \beta - \sin \alpha \cdot \sin \beta</math></li> <li>✓ <math>\left(-\frac{12}{13}\right) \cdot \left(-\frac{4}{5}\right) - \left(\frac{5}{13}\right) \cdot \left(\frac{3}{5}\right)</math></li> <li>✓ <math>\frac{33}{65}</math></li> </ul> <p style="text-align: right;">(6)</p> <p style="text-align: right;"><b>[11]</b></p>
<p>3(a)</p>	 $\sin 18^{\circ} = t = \frac{t}{1}$ $x^2 = r^2 - y^2$ $\therefore x^2 = 1^2 - t^2$ $\therefore x^2 = 1 - t^2$ $\therefore x = \sqrt{1 - t^2}$ $\therefore \cos 18^{\circ} = \frac{\sqrt{1 - t^2}}{1} = \sqrt{1 - t^2}$	<ul style="list-style-type: none"> <li>✓ diagram</li> <li>✓ Pythagoras</li> <li>✓ <math>x = \sqrt{1 - t^2}</math></li> <li>✓ <math>\cos 18^{\circ} = \frac{\sqrt{1 - t^2}}{1} = \sqrt{1 - t^2}</math></li> </ul> <p style="text-align: right;">(4)</p>
<p>3(b)</p>	$\sin 78^{\circ}$ $= \sin (60^{\circ} + 18^{\circ})$ $= \sin 60^{\circ} \cdot \cos 18^{\circ} + \cos 60^{\circ} \cdot \sin 18^{\circ}$ $= \left(\frac{\sqrt{3}}{2}\right) \cdot \sqrt{1 - t^2} + \left(\frac{1}{2}\right) \cdot t$ $= \frac{\sqrt{3} \sqrt{1 - t^2}}{2} + \frac{t}{2} = \frac{\sqrt{3(1 - t^2)} + t}{2}$	<ul style="list-style-type: none"> <li>✓ <math>\sin (60^{\circ} + 18^{\circ})</math></li> <li>✓ <math>\sin 60^{\circ} \cdot \cos 18^{\circ} + \cos 60^{\circ} \cdot \sin 18^{\circ}</math></li> <li>✓ <math>\left(\frac{\sqrt{3}}{2}\right)</math> and <math>\left(\frac{1}{2}\right)</math></li> <li>✓ <math>\sqrt{1 - t^2}</math> and <math>t</math></li> <li>✓ <math>\frac{\sqrt{3(1 - t^2)} + t}{2}</math></li> </ul> <p style="text-align: right;">(5)</p> <p style="text-align: right;"><b>[9]</b></p>

**SECTION C: HOMEWORK**

**QUESTION 1**

Prove that:

(a)  $(\tan x - 1)(\sin 2x - 2\cos^2 x) = 2(1 - 2\sin x \cos x)$  (6)



$$(b) \quad \frac{\cos 2x}{\cos x - \sin x} = \cos x + \sin x \quad (3)$$

[9]

**QUESTION 2**

$$(a) \quad \text{Show that } \sin(45^\circ - \alpha) = \frac{\sqrt{2}(\cos \alpha - \sin \alpha)}{2} \quad (3)$$

$$(b) \quad \text{Hence prove that } \sin 2\alpha + 2\sin^2(45^\circ - \alpha) = 1 \quad (6)$$

[9]

**QUESTION 3**

If  $\cos \beta = \frac{p}{\sqrt{5}}$  where  $p < 0$  and  $\beta \in [180^\circ; 360^\circ]$ , determine, using a diagram, an expression in terms of  $p$  for:

$$(a) \quad \tan \beta \quad (4)$$

$$(b) \quad \cos 2\beta \quad (3)$$

[7]

**QUESTION 4**

If  $\sin 61^\circ = \sqrt{a}$ , determine the value of the following in terms of  $a$ :


$$\cos 73^\circ \cos 15^\circ + \sin 73^\circ \sin 15^\circ \quad (6)$$

**SECTION D: SOLUTIONS TO HOMEWORK: TOPIC 2: TRIGONOMETRY (2)**


1(a)	$  \begin{aligned}  & (\tan x - 1)(\sin 2x - 2\cos^2 x) \\  &= \left( \frac{\sin x}{\cos x} - 1 \right) (2\sin x \cdot \cos x - 2\cos^2 x) \\  &= \left( \frac{\sin x - \cos x}{\cos x} \right) 2\cos x(\sin x - \cos x) \\  &= 2(\sin x - \cos x)^2 \\  &= 2(\sin^2 x - 2\sin x \cdot \cos x + \cos^2 x) \\  &= 2(1 - 2\sin x \cdot \cos x)  \end{aligned}  $	$  \begin{aligned}  & \checkmark \frac{\sin x}{\cos x} \\  & \checkmark 2\sin x \cos x \\  & \checkmark \frac{\sin x - \cos x}{\cos x} \\  & \checkmark 2\cos x(\sin x - \cos x) \\  & \checkmark 2(\sin^2 x - 2\sin x \cdot \cos x + \cos^2 x) \\  & \checkmark 2(1 - 2\sin x \cdot \cos x)  \end{aligned}  $ <p style="text-align: right;">(6)</p>
1(b)	$  \begin{aligned}  & \frac{\cos 2x}{\cos x - \sin x} \\  &= \frac{\cos^2 x - \sin^2 x}{\cos x - \sin x} \\  &= \frac{(\cos x - \sin x)(\cos x + \sin x)}{\cos x - \sin x} \\  &= \cos x + \sin x  \end{aligned}  $	$  \begin{aligned}  & \checkmark \cos^2 x - \sin^2 x \\  & \checkmark (\cos x - \sin x)(\cos x + \sin x) \\  & \checkmark \cos x + \sin x  \end{aligned}  $ <p style="text-align: right;">(3)</p> <p style="text-align: right;">[9]</p>

<p>2(a)</p>	$\begin{aligned} & \sin(45^\circ - \alpha) \\ &= \sin 45^\circ \cdot \cos \alpha - \cos 45^\circ \cdot \sin \alpha \\ &= \frac{\sqrt{2}}{2} \cdot \cos \alpha - \frac{\sqrt{2}}{2} \cdot \sin \alpha \\ &= \frac{\sqrt{2}}{2} (\cos \alpha - \sin \alpha) \\ &= \frac{\sqrt{2}(\cos \alpha - \sin \alpha)}{2} \end{aligned}$	$\begin{aligned} & \checkmark \sin 45^\circ \cdot \cos \alpha - \cos 45^\circ \cdot \sin \alpha \\ & \checkmark \frac{\sqrt{2}}{2} \cdot \cos \alpha - \frac{\sqrt{2}}{2} \cdot \sin \alpha \\ & \checkmark \frac{\sqrt{2}}{2} (\cos \alpha - \sin \alpha) \end{aligned} \quad (3)$
<p>2(b)</p>	$\begin{aligned} & \sin 2\alpha + 2\sin^2(45^\circ - \alpha) \\ &= 2\sin \alpha \cdot \cos \alpha + 2[\sin(45^\circ - \alpha)]^2 \\ &= 2\sin \alpha \cdot \cos \alpha + 2\left[\frac{\sqrt{2}(\cos \alpha - \sin \alpha)}{2}\right]^2 \\ &= 2\sin \alpha \cdot \cos \alpha + 2\left(\frac{2(\cos \alpha - \sin \alpha)^2}{4}\right) \\ &= 2\sin \alpha \cdot \cos \alpha + (\cos \alpha - \sin \alpha)^2 \\ &= 2\sin \alpha \cdot \cos \alpha + \cos^2 \alpha - 2\sin \alpha \cos \alpha + \sin^2 \alpha \\ &= \cos^2 \alpha + \sin^2 \alpha = 1 \end{aligned}$	$\begin{aligned} & \checkmark 2\sin \alpha \cdot \cos \alpha \\ & \checkmark 2\left[\frac{\sqrt{2}(\cos \alpha - \sin \alpha)}{2}\right]^2 \\ & \checkmark (\cos \alpha - \sin \alpha)^2 \\ & \checkmark \cos^2 \alpha - 2\sin \alpha \cos \alpha + \sin^2 \alpha \\ & \checkmark \cos^2 \alpha + \sin^2 \alpha \\ & \checkmark 1 \end{aligned} \quad (6)$



<p>3(a)</p>	$\begin{aligned} \cos \beta &= \frac{p}{\sqrt{5}} \\ x &= p \\ r &= \sqrt{5} \\ p^2 + y^2 &= (\sqrt{5})^2 \\ \therefore y^2 &= 5 - p^2 \\ \therefore y &= -\sqrt{5 - p^2} \\ \therefore \tan \beta &= \frac{-\sqrt{5 - p^2}}{p} \end{aligned}$ 	$\begin{aligned} & \checkmark \text{diagram} \\ & \checkmark \text{Pythagoras} \\ & \checkmark y = -\sqrt{5 - p^2} \\ & \checkmark \tan \beta = \frac{-\sqrt{5 - p^2}}{p} \end{aligned} \quad (4)$
<p>3(b)</p>	$\begin{aligned} \cos 2\beta &= 2\cos^2 \beta - 1 \\ &= 2\left(\frac{p}{\sqrt{5}}\right)^2 - 1 \\ &= \frac{2p^2}{5} - 1 \end{aligned}$	$\begin{aligned} & \checkmark \cos 2\beta = 2\cos^2 \beta - 1 \\ & \checkmark 2\left(\frac{p}{\sqrt{5}}\right)^2 - 1 \\ & \checkmark \frac{2p^2}{5} - 1 \end{aligned} \quad (3)$

[7]

4	$\sin 61^\circ = \frac{\sqrt{a}}{1}$ $x^2 + (\sqrt{a})^2 = (1)^2$ $\therefore x^2 = 1 - a$ $\therefore x = \sqrt{1 - a}$ $\cos 73^\circ \cos 15^\circ + \sin 73^\circ \sin 15^\circ$ $= \cos(73^\circ - 15^\circ)$ $= \cos 58^\circ$ $= 2 \cos^2 29^\circ - 1$ $= 2 \sin^2 61^\circ - 1$ $= 2(\sqrt{a})^2 - 1$ $= 2a - 1$		<ul style="list-style-type: none"> <li>✓ diagram</li> <li>✓ <math>x = \sqrt{1 - a}</math></li> <li>✓ <math>\cos 58^\circ</math></li> <li>✓ <math>2 \cos^2 29^\circ - 1</math></li> <li>✓ <math>2 \sin^2 61^\circ - 1</math></li> <li>✓ <math>2a - 1</math></li> </ul>
---	--	---	---

[6]



**CONSOLIDATION****Notes to Teachers**

Learners should attempt the questions under exam conditions. It is suggested that learners complete each section in time set and then to discuss the solutions before doing the second topic. Learners can find the solutions with Learner Homework Solutions. Encourage learners to re-do questions they did not get right for Homework.

**TOPIC 1: NUMBER PATTERNS, SEQUENCES AND SERIES****SECTION A: TYPICAL EXAM QUESTION****QUESTION 1***(DoE Nov 2008)**Time: 25 minutes*

1.1 Consider the sequence:  $\frac{1}{2}; 4; \frac{1}{4}; 7; \frac{1}{8}; 10; \dots$

1.1.1 If the pattern continues in the same way, write down the next TWO terms in the sequence. (1)

1.1.2 Calculate the sum of the first 50 terms of the sequence. (7)

1.2 Consider the sequence:  $8; 18; 30; 44; \dots$

1.2.1 Write down the next TWO terms of the sequence, if the pattern continues in the same way. (2)

1.2.2 Calculate the  $n^{\text{th}}$  term of the sequence. (6)


1.2.3 Which term of the sequence is 330? (6)

[22]

**SECTION B: SOLUTIONS TO SECTION A: TOPIC 1****QUESTION 1****Time for discussion of solution: 15 minutes**

1.1.1	$\frac{1}{16}; 13$	✓ answers (1)
-------	--------------------	---------------

MATHEMATICS GRADE 12 SESSION 8 (TEACHER NOTES)

1.1.2	$S_{50} = 25 \text{ terms of } 1^{\text{st}} \text{ sequence} + 25 \text{ terms of } 2^{\text{nd}} \text{ sequence}$ $S_{50} = \left( \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots \text{ to } 25 \text{ terms} \right) + (4 + 7 + 10 + 13 + \dots \text{ to } 25 \text{ terms})$ $S_{50} = \frac{\frac{1}{2} \left( \left( \frac{1}{2} \right)^{25} - 1 \right)}{\frac{1}{2} - 1} + \frac{25}{2} [2(4) + 24(3)]$ $S_{50} = 0,999999\dots + 1000$ $S_{50} = 1001,00$	<ul style="list-style-type: none"> <li>✓ separating the sequences</li> <li>✓✓ sum of geometric series</li> <li>✓✓ sum of arithmetic series</li> <li>✓✓ final answer (7)</li> </ul>
1.2.1	60 ; 78	✓✓ answers (2)
1.2.2	 <p style="text-align: center;"> <math>\therefore 2a = 2</math>      <math>3a + b = 10</math>      <math>a + b + c = 8</math>  <math>\therefore a = 1</math>      <math>\therefore 3(1) + b = 10</math>      <math>\therefore 1 + 7 + c = 8</math>  <math>\therefore b = 7</math>      <math>\therefore c = 0</math>  <math>\therefore T_n = n^2 + 7n</math> </p>	<ul style="list-style-type: none"> <li>✓ first difference row</li> <li>✓ second difference</li> <li>✓ <math>a = 1</math></li> <li>✓ <math>b = 7</math></li> <li>✓ <math>c = 0</math></li> <li>✓ <math>T_n = n^2 + 7n</math> (6)</li> </ul>
1.2.3	$330 = n^2 + 7n$ $\therefore -n^2 - 7n + 330 = 0$ $\therefore n^2 + 7n - 330 = 0$ $\therefore n = \frac{-7 \pm \sqrt{(7)^2 - 4(1)(-330)}}{2}$ $\therefore n = \frac{-7 \pm 37}{2}$ $\therefore n = 15 \text{ or } n = -22$ <p style="text-align: center;">(not valid)</p> <p>Alternatively, factorise:</p> $(n - 15)(n + 22) = 0$	<ul style="list-style-type: none"> <li>✓ <math>330 = n^2 + 7n</math></li> <li>✓ <math>n^2 + 7n - 330 = 0</math></li> <li>✓✓ Substitution into formula or factorising</li> <li>✓ <math>n = 15</math> or <math>n = -22</math></li> <li>✓ <math>n \neq -22</math> (6)</li> </ul> <p style="text-align: right;"><b>[22]</b></p>

**CONSOLIDATION: TOPIC 2: FINANCIAL MATHS AND TRIGONOMETRY**

Time: 25 minutes

**SECTION A: TYPICAL EXAM QUESTIONS****QUESTION 1**


- 1.1 Nicholas decides to invest money into the share market in order to save up R800 000 in ten years time. He believes that he can average a return of 18% per annum compounded monthly. Starting immediately, he starts making monthly payments into a share market account.
- 1.1.1 How much must Nicholas invest per month in order to obtain his R800 000? (5)
- 1.1.2 At the end of the ten-year period, Nicholas decides not to spend the R800 000 but to rather invest it at an interest rate of 18% per annum compounded half-yearly. How much money will he have then saved four years later? (4)
- 1.2 Bonolo borrows money from the bank in order to finance a new home. She takes out a twenty year loan and begins to make monthly payments of R6000 per month starting in one month's time. The current interest rate is 15% per annum compounded monthly. Calculate how much the new home originally cost. (4)  
[9]

**QUESTION 2**


- 2.1 If  $13\sin\theta - 5 = 0$  where  $\theta \in [90^\circ; 360^\circ]$  and  $5\cos\beta + 4 = 0$  where  $\tan\beta > 0$ , calculate without the use of a calculator and with the aid of diagrams the value of the following:
- 2.1.1  $\sin(\theta - \beta)$  (9)
- 2.1.2  $\tan(90^\circ + \theta)$  (5)
- 2.2 Simplify without using a calculator:
- 2.2.1  $\cos(50^\circ + x)\cos(20^\circ + x) + \sin(50^\circ + x)\sin(20^\circ + x)$  (3)
- 2.2.2  $\cos(-140^\circ)\cos 740^\circ - \sin 140^\circ\sin(-20^\circ)$  (6)  
[23]

**SECTION B: SOLUTIONS TO SECTION A: TOPIC 2**

Time for discussion of solution: 20 minutes



<p>1.1.1</p>	$800\,000 = \frac{x \left[ \left( 1 + \frac{0,18}{12} \right)^{121} - 1 \right]}{\frac{0,18}{12}}$ <p><math>\therefore x = R2372,07</math></p>	<ul style="list-style-type: none"> <li>✓ formula</li> <li>✓ <math>n = 121</math></li> <li>✓ <math>\frac{0,18}{12}</math></li> <li>✓ <math>F = 800\,000</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(5)</p>
<p>1.1.2</p>	$A = 800\,000 \left( 1 + \frac{0,18}{2} \right)^8$ <p><math>\therefore A = R1\,594\,050,11</math></p>	<ul style="list-style-type: none"> <li>✓ formula</li> <li>✓ <math>n = 8</math></li> <li>✓ <math>\frac{0,18}{2}</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p>



<p>1.2</p>	$P = \frac{6000 \left[ 1 - \left( 1 + \frac{0,15}{12} \right)^{-240} \right]}{\frac{0,15}{12}}$ <p><math>\therefore x = R455\,653,67</math></p>	<ul style="list-style-type: none"> <li>✓ formula</li> <li>✓ <math>n = 240</math></li> <li>✓ <math>\frac{0,15}{12}</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p>
------------	---	---

2.1.1

$$13\sin\theta - 5 = 0$$

$$\therefore \sin\theta = \frac{5}{13} \text{ (positive in Quad 1 or 2)}$$


$$\theta \in [90^\circ; 360^\circ] \text{ (Quad 2, 3 or 4)}$$

Common quad is 2

$$x^2 + (5)^2 = (13)^2$$

$$\therefore x^2 = 144$$

$$\therefore x = -12$$



$$5\cos\beta + 4 = 0$$

$$\therefore \cos\beta = \frac{-4}{5} \text{ (negative in Quad 2 or 3)}$$


$$\tan\beta > 0 \text{ (Quad 1 or 3)}$$

Common quad is 3

$$(-4)^2 + y^2 = (5)^2$$

$$\therefore y^2 = 9$$

$$\therefore y = -3$$



$$\sin(\theta - \beta)$$

$$= \sin\theta\cos\beta - \cos\theta\sin\beta$$

$$= \left(\frac{5}{13}\right)\left(\frac{-4}{5}\right) - \left(\frac{-12}{13}\right)\left(\frac{-3}{5}\right)$$

$$= \frac{-20}{65} - \frac{36}{65}$$

$$= -\frac{56}{65}$$

- ✓  $\sin\theta = \frac{5}{13}$
- ✓ diagram for  $\theta$
- ✓  $x = -12$
- ✓  $\cos\beta = \frac{-4}{5}$
- ✓ diagram for  $\beta$
- ✓  $y = -3$
- ✓  $\sin\theta\cos\beta - \cos\theta\sin\beta$
- ✓ correct substitution
- ✓ answer

(9)



<p>2.1.2</p>	$\begin{aligned} & \tan(90^\circ + \theta) \\ &= \frac{\sin(90^\circ + \theta)}{\cos(90^\circ + \theta)} \\ &= \frac{\cos \theta}{-\sin \theta} \\ &= \frac{-12}{-\frac{5}{13}} \\ &= \frac{12}{5} \end{aligned}$	<ul style="list-style-type: none"> <li>✓ <math>\frac{\sin(90^\circ + \theta)}{\cos(90^\circ + \theta)}</math></li> <li>✓ <math>\cos \theta</math></li> <li>✓ <math>-\sin \theta</math></li> <li>✓ correct substitution</li> <li>✓ answer</li> </ul> <p style="text-align: right;">(5)</p>
<p>2.2.1</p>	$\begin{aligned} & \cos(50^\circ + x) \cos(20^\circ + x) + \sin(50^\circ + x) \sin(20^\circ + x) \\ &= \cos[(50^\circ + x) - (20^\circ + x)] \\ &= \cos 30^\circ \\ &= \frac{\sqrt{3}}{2} \end{aligned}$	<ul style="list-style-type: none"> <li>✓ <math>\cos[(50^\circ + x) - (20^\circ + x)]</math></li> <li>✓ <math>\cos 30^\circ</math></li> <li>✓ <math>\frac{\sqrt{3}}{2}</math></li> </ul> <p style="text-align: right;">(3)</p>
<p>2.2.2</p>	$\begin{aligned} & \cos(-140^\circ) \cos 740^\circ - \sin 140^\circ \sin(-20^\circ) \\ &= \cos 140^\circ \cos 20^\circ - (\sin 40^\circ)(-\sin 20^\circ) \\ &= (-\cos 40^\circ)(\cos 20^\circ) + \sin 40^\circ \sin 20^\circ \\ &= -\cos 40^\circ \cos 20^\circ + \sin 40^\circ \sin 20^\circ \\ &= -(\cos 40^\circ \cos 20^\circ - \sin 40^\circ \sin 20^\circ) \\ &= -\cos 60^\circ \\ &= -\frac{1}{2} \end{aligned}$	<ul style="list-style-type: none"> <li>✓ <math>-\cos 40^\circ</math></li> <li>✓ <math>\cos 20^\circ</math></li> <li>✓ <math>\sin 40^\circ</math></li> <li>✓ <math>-\sin 20^\circ</math></li> <li>✓ <math>-\cos 60^\circ</math></li> <li>✓ <math>-\frac{1}{2}</math></li> </ul> <p style="text-align: right;">(6)</p>

[23]

## CONSOLIDATION

**Note to Teachers:** Encourage learners to complete each section in the time allocated. 80 minutes is allocated to doing questions. Learners need to check solutions on their own.

## TOPIC 1: NUMBER PATTERNS, SEQUENCES AND SERIES

## SECTION A: TYPICAL EXAM QUESTIONS

## QUESTION 1: 25 minutes

1.1 Consider the following sequence:  $\frac{3}{2}; \frac{5}{4}; \frac{7}{8}; \frac{9}{16}; \dots$

1.1.1 Write down the next two terms of the sequence. (2)


1.1.2 Determine the  $n$ th term of the sequence in simplified form. (5)

1.2 By using appropriate formulae and without using a calculator, calculate the value of the following:

1.2.1  $\sum_{k=2}^8 \left(\frac{1}{2}\right)^{k-1}$  (4)

1.2.2  $\sum_{k=2}^{\infty} \left(\frac{1}{2}\right)^{k-1}$  (2)

1.3 In the figure below, a stack of cans is shown. There are 30 cans in the first layer, 29 cans in the second layer (lying on top of the first layer), 28 cans in the third layer. This pattern of stacking continues.



Determine the maximum number of cans that can be stacked in this way. (4)

[17]

## QUESTION 2: 15 minutes

Consider the sequence:  $3; a; 10; b; 21; \dots$

The sequence has a constant second difference of 1.

2.1 Determine the value of  $a$  and  $b$ . (4)

2.2 Determine the  $n$ th term of the sequence. (5)

2.3 Hence, prove that the sum of any two consecutive numbers in the sequence equals a square number. (4)

[13]

TOPIC 2: FINANCIAL MATHEMATICS

**SECTION A: TYPICAL EXAM QUESTIONS**

**QUESTION 1: 40 minutes**

- 1.1 A motor car costing R200 000 depreciated at a rate of 8% per annum on the reducing balance method. Calculate how long it took for the car to depreciate to R90 000 under these conditions. (4)
- 1.2 Mpho starts a five year savings plan. At the beginning of the month he deposits R2000 into the account and makes a further deposit of R2000 at the end of that month. He then continues to make month end payments of R2000 into the account for the five year period (starting from his first deposit). The interest rate is 6% per annum compounded monthly.
- 1.2.1 Calculate the future value of his investment at the end of the five year period. (4)
- 1.2.2 Due to financial difficulty, Mpho misses the last two payments of R2000. What will the value of his investment now be at the end of the five year period? (4)
- 1.3 Lucy takes out a twenty year loan of R400 000. She repays the loan by means of equal monthly payments starting one month after the granting of the loan. The interest rate is 18% per annum compounded monthly.
- 1.3.1 Calculate the monthly repayments. (3)
- 1.3.2 Calculate the amount owed after the 3<sup>rd</sup> payment was made. (2)
- 1.3.3 Due to financial difficulty, Brenda misses the 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> payments. Calculate her increased monthly payment which comes into effect from the 7<sup>th</sup> payment onwards. (4)
- [21]

**SECTION B: SOLUTIONS AND HINTS TO SECTION A**

**1. TOPIC 1: NUMBER PATTERNS, SEQUENCES AND SERIES**

**QUESTION 1**

1.1.1	$\frac{11}{32}; \frac{13}{64}$	✓✓ answers (2)
-------	--------------------------------	----------------

## MATHEMATICS

## GRADE 12

## SESSION 9

## (TEACHER NOTES)

1.1.2	<p>The numerators represent an arithmetic sequence:  <math>T_n = 3 + (n-1)(2)</math>  <math>\therefore T_n = 3 + 2n - 2</math>  <math>\therefore T_n = 2n + 1</math></p> <p>The denominators represent a geometric sequence:  <math>T_n = (2)(2)^{n-1}</math>  <math>\therefore T_n = 2^n</math></p> <p>The general term for the given sequence is:  <math>T_n = \frac{2n+1}{2^n}</math></p>	<p>✓ <math>a=3</math> and <math>d=2</math>  ✓ <math>T_n = 2n+1</math>  ✓ <math>a=2</math> and <math>r=2</math>  ✓ <math>T_n = 2^n</math>  ✓ <math>T_n = \frac{2n+1}{2^n}</math></p> <p>(5)</p>
1.2.1	$\sum_{k=2}^8 \left(\frac{1}{2}\right)^{k-1} = \left(\frac{1}{2}\right)^1 + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^3 + \dots + \left(\frac{1}{2}\right)^7$ $\therefore S_7 = \frac{\left(\frac{1}{2}\right)\left(1 - \left(\frac{1}{2}\right)^7\right)}{1 - \frac{1}{2}} = 1 - \left(\frac{1}{2}\right)^7 = 1 - \frac{1}{128} = \frac{127}{128}$	<p>✓ expansion  ✓ correct formula  ✓ correct substitution  ✓ <math>\frac{127}{128}</math></p> <p>(4)</p>
1.2.2	$\sum_{k=2}^{\infty} \left(\frac{1}{2}\right)^{k-1} = \left(\frac{1}{2}\right)^1 + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^3 + \dots$ $\therefore S_{\infty} = \frac{\frac{1}{2}}{1 - \frac{1}{2}} = 1$	<p>✓ correct formula  ✓ answer</p> <p>(2)</p>
1.3	<p>The layers are stacked as follows:  30; 29; 28; 27; .....; 1</p> <p>This forms an arithmetic sequence.  <math>\therefore S_{30} = \frac{30}{2} [2(30) + (30-1)(-1)]</math>  <math>\therefore S_{30} = 465</math></p> <p>There are a maximum of 465 cans that can be stacked in this way.</p>	<p>✓ 30; 29; 28; 27; .....; 1  ✓ correct formula  ✓ correct substitution  ✓ 465 cans</p> <p>(4)</p> <p>[17]</p>

QUESTION 2

<p>2.1</p>	<p>3; a; 10; b; 21; .....</p> $(10 - a) - (a - 3) = 1$ $\therefore 10 - a - a + 3 = 1$ $\therefore -2a = -12$ $\therefore a = 6$ $(21 - b) - (b - 10) = 1$ $\therefore 21 - b - b + 10 = 1$ $\therefore -2b = -30$ $\therefore b = 15$	<p>✓ <math>(10 - a) - (a - 3) = 1</math>                  ✓ <math>a = 6</math>                  ✓ <math>(21 - b) - (b - 10) = 1</math>                  ✓ <math>b = 15</math></p> <p>(4)</p>
<p>2.2</p>	<p>3; 6; 10; 15; 21; .....</p> $2a = 1$ $\therefore a = \frac{1}{2}$ $3a + b = 3$ $\therefore 3\left(\frac{1}{2}\right) + b = 3$ $\therefore b = \frac{3}{2}$ $a + b + c = 3$ $\therefore \frac{1}{2} + \frac{3}{2} + c = 3$ $\therefore c = 1$ $\therefore T_n = \frac{1}{2}n^2 + \frac{3}{2}n + 1$	<p>✓ 3; 6; 10; 15; 21; .....</p> <p>✓ <math>a = \frac{1}{2}</math>                  ✓ <math>b = \frac{3}{2}</math>                  ✓ <math>c = 1</math>                  ✓ <math>T_n = \frac{1}{2}n^2 + \frac{3}{2}n + 1</math></p> <p>(5)</p>
<p>2.3</p>	$\left(\frac{1}{2}n^2 + \frac{3}{2}n + 1\right) + \left(\frac{1}{2}(n+1)^2 + \frac{3}{2}(n+1) + 1\right)$ $= \frac{1}{2}n^2 + \frac{3}{2}n + 1 + \frac{1}{2}(n^2 + 2n + 1) + \frac{3}{2}n + \frac{3}{2} + 1$ $= \frac{1}{2}n^2 + \frac{3}{2}n + 1 + \frac{1}{2}n^2 + n + \frac{1}{2} + \frac{3}{2}n + \frac{3}{2} + 1$ $= n^2 + 4n + 4$ $= (n + 2)^2$	<p>✓ <math>T_n + T_{n+1}</math>                  ✓ expanding                  ✓ <math>n^2 + 4n + 4</math>                  ✓ <math>(n + 2)^2</math></p> <p>(4)</p>

[13]

MATHEMATICS



GRADE 12


SESSION 9

(TEACHER NOTES)

## TOPIC 2: FINANCIAL MATHEMATICS

## QUESTION 1

1.1	$A = P(1 - i)^n$ $\therefore 90\,000 = 200\,000(1 - 0,08)^n$ $\therefore 90\,000 = 200\,000(0,92)^n$ $\therefore \frac{90\,000}{200\,000} = (0,92)^n$ $\therefore 0,45 = (0,92)^n$ $\therefore \log_{0,92} 0,45 = n$ $\therefore n = 9,576544593$ <p>It will take approximately 9 years 7 months to depreciate to R90 000.</p>	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ correct substitution</li> <li>✓ use of logs</li> <li>✓ value of <math>n</math></li> </ul> <p style="text-align: right;">(4)</p>
1.2.1	 $F = \frac{2000 \left[ \left( 1 + \frac{0,06}{12} \right)^{61} - 1 \right]}{\frac{0,06}{12}}$ $\therefore F = R142\,237,76$	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>\frac{0,06}{12}</math></li> <li>✓ value of <math>n</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p>
1.2.2	 $F = \frac{2000 \left[ \left( 1 + \frac{0,06}{12} \right)^{59} - 1 \right]}{\frac{0,06}{12}} \cdot \left( 1 + \frac{0,06}{12} \right)^2$ $\therefore F = R138\,227,76$	<ul style="list-style-type: none"> <li>✓ correct annuity formula</li> <li>✓ <math>n = 59</math></li> <li>✓ <math>\left( 1 + \frac{0,06}{12} \right)^2</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p>

<p>1.3.1</p>	 $400\,000 = \frac{x \left[ 1 - \left( 1 + \frac{0,18}{12} \right)^{-240} \right]}{\frac{0,18}{12}}$ $\therefore \frac{400\,000 \times \frac{0,18}{12}}{\left[ 1 - \left( 1 + \frac{0,18}{12} \right)^{-240} \right]} = x$ $\therefore x = R6173,25$	<ul style="list-style-type: none"> <li>✓ formula</li> <li>✓ correct substitution</li> <li>✓ answer</li> </ul> <p style="text-align: right;">(3)</p>
<p>1.3.2</p>	$B = \frac{6173,25 \left[ 1 - (1,015)^{-237} \right]}{0,015}$ $\therefore B = R399\,472,68$	<ul style="list-style-type: none"> <li>✓ correct substitution</li> <li>✓ answer</li> </ul> <p style="text-align: right;">(2)</p>
<p>1.3.3</p>	$399\,472,68 \left( 1 + \frac{0,18}{12} \right)^3 = \frac{x \left[ 1 - \left( 1 + \frac{0,18}{12} \right)^{-234} \right]}{\frac{0,18}{12}}$ $\therefore \frac{399\,472,68 \left( 1 + \frac{0,18}{12} \right)^3 \times \frac{0,18}{12}}{\left[ 1 - \left( 1 + \frac{0,18}{12} \right)^{-234} \right]} = x$ $\therefore x = R6464,16$	<ul style="list-style-type: none"> <li>✓ <math>399\,472,68 \left( 1 + \frac{0,18}{12} \right)^3</math></li> <li>✓ present value formula</li> <li>✓ <math>n = 234</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(4)</p>

[21]

## SECTION C: HOMEWORK

## TOPIC 1: NUMBER PATTERNS, SEQUENCES AND SERIES

## QUESTION 1


- 1.1 Calculate the value of  $\sum_{r=3}^{17} 10$  (1)
- 1.2 An auditorium has 30 rows of seats. There are 20 seats in the first row and 136 seats in the last row. The number of seats in the first row, second row, third row and so forth forms an arithmetic sequence.
- 1.2.1 Determine the number of seats in the second row. (4)
- 1.2.2 Determine the total number of seats in the auditorium. (1)
- 1.3 The sequence  $12; x; \dots$  is a convergent geometric sequence in which the sum to infinity is equal to 24.
- 1.3.1 Determine the value of  $x$ . (3)
- 1.3.2 Hence determine which term of the sequence is equal to  $\frac{3}{128}$ . (3)

[11]



## QUESTION 2

A sequence of isosceles triangles is drawn. The first triangle has a base of 2cm and height of 2cm. The second triangle has a base that is 2cm longer than the base of the first triangle. The height of the second triangle is 1cm longer than the height of the first triangle. This pattern of enlargement will continue with each triangle that follows.



the 100<sup>th</sup> triangle.

- 2.1 Determine the area of (4)
- 2.2 Which triangle will have an area of  $240\text{cm}^2$ ? (4)

[8]



TOPIC 2: FINANCIAL MATHEMATICS

QUESTION 1

- 1.1 Joshua takes out a retirement annuity that will supplement his pension when he retires in thirty years' time. He estimates that he will need R3000 000 in this retirement fund at that stage. The interest rate he earns is 12% per annum compounded monthly. Calculate his monthly payment into this fund if he starts paying immediately and makes his final payment in 20 years' time. (3)
- 1.2 Mpho takes out a bank loan for R250 000. The interest rate charged by the bank is 18,5% per annum compounded monthly.
- 1.2.1 What will his monthly repayment be if he pays the loan back over five years, starting **FOUR** months after the granting of the loan? (5)
- 1.2.2 Calculate the balance outstanding after the 25<sup>th</sup> repayment. (5)
- [13]

SECTION D: SOLUTIONS AND HINTS TO HOMEWORK

TOPIC 1: NUMBER PATTERNS, SEQUENCES AND SERIES

QUESTION 1



1.1	$\sum_{r=3}^{17} 10 = (\text{number of terms}) \times 10$ $= (17 - 3 + 1) \times 10$ $= 15 \times 10$ $= 150$	✓ 150 (1)
1.2.1	$a = 20$ $a + (n - 1)d = 136$ $\therefore 20 + (30 - 1)d = 136$ $\therefore 20 + 29d = 136$ $\therefore 29d = 116$ $\therefore d = 4$ <p>There are 24 seats in the second row.</p>	✓ $a = 20$ ✓ $20 + 29d = 136$ ✓ $d = 4$ ✓ 24 seats in second row ✓ $S_{30} = 2340$ (4)
1.2.2	$S_{30} = \frac{30}{2} [20 + 136] = 2340 \text{ seats}$	✓ $S_{30} = 2340$ (1)

## MATHEMATICS

## GRADE 12

## SESSION 9

## (TEACHER NOTES)

1.3.1	$r = \frac{x}{12}$ $\frac{12}{1 - \frac{x}{12}} = 24$ $\therefore 12 = 24 \left( 1 - \frac{x}{12} \right)$ $\therefore 12 = 24 - 2x$ $\therefore 2x = 12$ $\therefore x = 6$	$\checkmark r = \frac{x}{12}$ $\checkmark \frac{12}{1 - \frac{x}{12}} = 24$ $\checkmark x = 6$ <p style="text-align: right;">(3)</p>
1.3.2	$12 \left( \frac{1}{2} \right)^{n-1} = \frac{3}{128}$ $\therefore \left( \frac{1}{2} \right)^{n-1} = \frac{1}{512}$ $\therefore \left( \frac{1}{2} \right)^{n-1} = \left( \frac{1}{2} \right)^9$ $\therefore n - 1 = 9$ $\therefore n = 10$	$\checkmark 12 \left( \frac{1}{2} \right)^{n-1} = \frac{3}{128}$ $\checkmark \left( \frac{1}{2} \right)^{n-1} = \frac{1}{512}$ $\checkmark n = 10$ <p style="text-align: right;">(3)</p> <p style="text-align: right;"><b>[12]</b></p>

## QUESTION 2



2.1	<p>Area of triangle 1: <math>\frac{1}{2}(2cm)(2cm) = (1)(2)cm^2</math></p> <p>Area of triangle 2: <math>\frac{1}{2}(4cm)(3cm) = (2)(3)cm^2</math></p> <p>Area of triangle 3: <math>\frac{1}{2}(6cm)(4cm) = (3)(4)cm^2</math></p> <p>Area of triangle 4: <math>\frac{1}{2}(8cm)(5cm) = (4)(5)cm^2</math></p> <p>The areas form the following pattern:  <math>(1)(2); (2)(3); (3)(4); (4)(5); \dots</math></p> <p>Area of triangle <math>n</math>: <math>(n)(n+1)cm^2</math></p> <p>Area of triangle 100: <math>(100)(100+1)cm^2 = 10100cm^2</math></p>	<ul style="list-style-type: none"> <li><math>\checkmark</math> determining areas</li> <li><math>\checkmark</math> establishing pattern</li> <li><math>\checkmark</math> obtaining general term</li> <li><math>\checkmark</math> area of 100<sup>th</sup> triangle</li> </ul> <p style="text-align: right;">(4)</p>
-----	---	--

2.2	$n(n+1) = 240$ $\therefore n^2 + n - 240 = 0$ $\therefore (n+16)(n-15) = 0$ $\therefore n = -16 \text{ or } n = 15$ <p>But <math>n \neq -16</math></p> $\therefore n = 15$ <p>The 15th triangle will have an area of <math>240 \text{ cm}^2</math></p>	<ul style="list-style-type: none"> <li>✓ correct <math>a</math> and <math>d</math></li> <li>✓ correct <math>n</math></li> <li>✓ <math>S_n</math> formula</li> <li>✓ correct answer</li> </ul> <p style="text-align: right;">(4)</p> <p style="text-align: right;"><b>[8]</b></p>
-----	--	--

**TOPIC 2: FINANCIAL MATHEMATICS**

**QUESTION 1**

1.1	$3000000 = \frac{x \left[ \left( 1 + \frac{0,12}{12} \right)^{241} - 1 \right]}{\frac{0,12}{12}}$ $\therefore x = \text{R}2999,56$	<ul style="list-style-type: none"> <li>✓ formula</li> <li>✓ correct substitution</li> <li>✓ answer</li> </ul> <p style="text-align: right;">(3)</p>
1.2.1	$250\,000 \left( 1 + \frac{0,185}{12} \right)^3 = \frac{x \left[ 1 - \left( 1 + \frac{0,185}{12} \right)^{-57} \right]}{\frac{0,185}{12}}$ $\therefore x = \text{R } 6\,934,45$	<ul style="list-style-type: none"> <li>✓ correct formula</li> <li>✓ <math>250\,000 \left( 1 + \frac{0,185}{12} \right)^3</math></li> <li>✓ <math>n = 57</math></li> <li>✓ <math>\frac{0,185}{12}</math> ✓ answer</li> </ul> <p style="text-align: right;">(5)</p>
1.2.2	$B = 250\,000 \left( 1 + \frac{0,185}{12} \right)^{28} - \frac{6\,934,45 \left[ \left( 1 + \frac{0,185}{12} \right)^{25} - 1 \right]}{\frac{0,185}{12}}$ $\therefore B = \text{R}174\,122,48$ <p>Alternatively:</p> $B = \frac{6\,934,45 \left[ 1 - \left( 1 + \frac{0,185}{12} \right)^{-32} \right]}{\frac{0,185}{12}}$ $\therefore B = \text{R}174\,122,75$	<ul style="list-style-type: none"> <li>✓ growing 250 000</li> <li>✓ <math>n = 28</math></li> <li>✓ future value formula</li> <li>✓ <math>n = 25</math></li> <li>✓ answer</li> </ul> <p>OR</p> <ul style="list-style-type: none"> <li>✓✓ present value formula</li> <li>✓ 6934,45</li> <li>✓ <math>n = 32</math></li> <li>✓ answer</li> </ul> <p style="text-align: right;">(5)</p>

**[13]**

## TOPIC : TRANSFORMATIONS

**Teacher note:** Transformations are easy to master and learners should score well in questions involving this topic. Ensure that they know the different algebraic transformation rules.


## LESSON OVERVIEW

1. Introduction session: 5 minutes
2. Typical exam questions:  
Question 1: 30 minutes  
Question 2: 10 minutes
3. Discussion of solutions 45 minutes

## SECTION A: TYPICAL EXAM QUESTIONS

## QUESTION 1 30 minutes 16 marks

- 1.1 On the diagram below, represent the point  $A(-3;2)$ .



- 1.2 Now represent the following points on the diagram provided above:  
Point B, the rotation of point A,  $90^\circ$  anticlockwise about the origin.  
Point C, the rotation of point A,  $180^\circ$  about the origin.  
Point D, the rotation of point A,  $90^\circ$  clockwise about the origin.


(3)

- 1.3 Write down algebraic rules to describe the above transformations. (3)
- 1.4 What type of quadrilateral is figure ABCD? Explain by referring to the properties of quadrilaterals. (2)
- 1.5 Figure **ABCD** is enlarged by a scale factor of 2 units through the origin to form its image **A'B'C'D'**. On the diagram provided on the previous page, sketch the image **A'B'C'D'** and indicate the coordinates of the vertices. (2)
- 1.6 Determine the ratio:  $\frac{\text{Area ABCD}}{\text{Area A'B'C'D'}}$  (1)
- 1.7 **ABCD** is reflected about the  $y$ -axis to form its image **EFGH**.  
 1.7.1 Write down the coordinates of E. (1)  
 1.7.2 Determine the ratio:  $\frac{\text{Perimeter ABCD}}{\text{Perimeter EFGH}}$  (1)
- 1.8 Describe, using words and algebraic notation, the single transformations involved if figure **ABCD** is transformed by the rule:  
 $(x; y) \rightarrow \left(\frac{1}{2}x; -\frac{1}{2}y - 1\right)$  (3)

**QUESTION 2      10 minutes      6 marks**

- 2.1 Show that the coordinates of **P'**, the image of **P(x; y)** rotated about the origin through an angle of  $135^\circ$ , in the anti-clockwise direction, is given by:

$$\left(-\frac{\sqrt{2}}{2}x - \frac{\sqrt{2}}{2}y; -\frac{\sqrt{2}}{2}y + \frac{\sqrt{2}}{2}x\right) \quad (4)$$




- 2.2 **M'** is the image of **M(2; 4)** under a rotation about the origin through  $135^\circ$ , in the anti-clockwise direction.

Determine the coordinates of  $M'$ , using the results in (a) (2)

**SECTION B: SOLUTIONS AND HINTS TO SECTION A**

**QUESTION 1**

1.1 ; 1.2



1.2	B(-2;-3) C(3;-2) D(2;3)	<ul style="list-style-type: none"> <li>✓ B(-2;-3)</li> <li>✓ C(3;-2)</li> <li>✓ D(2;3)</li> </ul>	(3)
1.3	90° anticlockwise: $(x; y) \rightarrow (-y; x)$ 180° anti-clockwise or clockwise: $(x; y) \rightarrow (-x; -y)$ 90° clockwise: $(x; y) \rightarrow (y; -x)$	<ul style="list-style-type: none"> <li>✓ <math>(x; y) \rightarrow (-y; x)</math></li> <li>✓ <math>(x; y) \rightarrow (-x; -y)</math></li> <li>✓ <math>(x; y) \rightarrow (y; -x)</math></li> </ul>	(3)
1.4	ABCD is a square since: Diagonals are equal in length Diagonals bisect each other at right angles	<ul style="list-style-type: none"> <li>✓ square</li> <li>✓ properties</li> </ul>	(2)

1.5	$A'(-6;4)$ $B'(-4;-6)$ $C'(6;-4)$ $D'(4;6)$ See diagram	✓ correct coordinates indicated ✓ joining points to form enlarged square (2)
1.6	$\frac{\text{Area ABCD}}{\text{Area A'B'C'D'}} = \frac{1}{2^2} = \frac{1}{4}$	✓ $\frac{1}{4}$ (1)
1.7.1	E(3;2)	✓ answer (1)
1.7.2	$\frac{\text{Perimeter ABCD}}{\text{Perimeter EFGH}} = \frac{4 \times \text{side AB}}{4 \times \text{side EF}} = 1$ (since AB = EF)	✓ answer (1)
1.8	$(x; y) \rightarrow \left(\frac{1}{2}x; \frac{1}{2}y\right)$ reduction by a factor of $\frac{1}{2}$ $\left(\frac{1}{2}x; \frac{1}{2}y\right) \rightarrow \left(\frac{1}{2}x; -\frac{1}{2}y\right)$ reflection about $x$ -axis $\left(\frac{1}{2}x; -\frac{1}{2}y\right) \rightarrow \left(\frac{1}{2}x; -\frac{1}{2}y - 1\right)$ translation 1 unit downwards $\therefore (x; y) \rightarrow \left(\frac{1}{2}x; -\frac{1}{2}y - 1\right)$	✓ reduction ✓ reflection ✓ translation (3)

[16]

QUESTION 2


2.1	$x' = x \cos(135^\circ) - y \sin(135^\circ)$ $x' = -x \cos 45^\circ - y \sin 45^\circ$ $x' = x \left(\frac{-\sqrt{2}}{2}\right) - y \left(\frac{\sqrt{2}}{2}\right)$ $x' = -\frac{\sqrt{2}}{2}x - \frac{\sqrt{2}}{2}y$ $y' = y \cos(135^\circ) + x \sin(135^\circ)$ $y' = -y \cos 45^\circ + x \sin 45^\circ$ $y' = y \left(-\frac{\sqrt{2}}{2}\right) + x \left(\frac{\sqrt{2}}{2}\right)$ $y' = -\frac{\sqrt{2}}{2}y + \frac{\sqrt{2}}{2}x$	✓ $x' = x \cos(135^\circ) - y \sin(135^\circ)$ ✓ $x' = -\frac{\sqrt{2}}{2}x - \frac{\sqrt{2}}{2}y$ ✓ $y' = y \cos(135^\circ) + x \sin(135^\circ)$ ✓ $y' = -\frac{\sqrt{2}}{2}y + \frac{\sqrt{2}}{2}x$ (4)
-----	--	---

2.2	$x' = -\frac{\sqrt{2}}{2}(2) - \frac{\sqrt{2}}{2}(4)$ $x' = -\sqrt{2} - 2\sqrt{2}$ $x' = -3\sqrt{2}$ $y' = -\frac{\sqrt{2}}{2}(4) + \frac{\sqrt{2}}{2}(2)$ $y' = -\sqrt{2}$ $\therefore M(-3\sqrt{2}; -\sqrt{2})$	$\checkmark x' = -3\sqrt{2}$ $\checkmark y' = -\sqrt{2}$	(2)
			[6]

### SECTION C: HOMEWORK

#### QUESTION 1

- 1.1 The point  $P(-2; 5)$  lies in a Cartesian plane. Determine the coordinates of  $P'$ , the image of  $P$  if:
- 1.1.1  $P$  is reflected about the line  $y = x$  (1)
- 1.1.2  $P$  has been rotated about the origin through  $90^\circ$  in a clockwise direction. (2)
- 1.2 KUHLE has undergone two transformations to obtain  $K''U''H''L''E''$ .  
 $K''(2; 7)$ ,  $U''(3; 9)$ ,  $H''(4; 8)$ ,  $L''(5; 9)$  and  $E''(6; 7)$  are the coordinates of the vertices of  $K''U''H''L''E''$ .





MATHEMATICS GRADE 12 SESSION 10 (TEACHER NOTES)


1.2.1 Describe, in words, two transformations of KUHLE (in the order which they occurred), to obtain  $K''U''H''L''E''$  (4)

1.2.2 Write down TWO possible sets of coordinates for  $H'$ , the image of H after the first transformation. (2)

1.2.3 Determine:  $\frac{\text{Area KUHLE}}{\text{Area } K''U''H''L''E''}$  (2)

[11]

QUESTION 2



2.1 The point  $P(3; 2)$  is rotated about the origin through an angle of  $120^\circ$  in an anti-clockwise direction. Determine  $x'$  and  $y'$ , the coordinates of  $P'$ . (6)

2.2 The same rotation sends a point Q into  $(-2; 0)$ . Determine the coordinates of Q, (4)  
 [10]

SECTION D: SOLUTIONS TO HOMEWORK SESSION 6

QUESTION 1

1.1.1	$P'(5; -2)$	✓ answer (1)
1.1.2	$P'(5; 2)$	✓ x-coordinate ✓ y-coordinate (2)
1.2.1	Reduction by a scale factor of $\frac{1}{2}$ : $(x; y) \rightarrow \left(\frac{1}{2}x; \frac{1}{2}y\right)$ Reflection about the line $y = x$ $\left(\frac{1}{2}x; \frac{1}{2}y\right) \rightarrow \left(\frac{1}{2}y; \frac{1}{2}x\right)$ $\therefore (x; y) \rightarrow \left(\frac{1}{2}y; \frac{1}{2}x\right)$	✓✓ reduction ✓✓ reflection (4)

MATHEMATICS

GRADE 12

SESSION 10

(TEACHER NOTES)

1.2.2	<p>If the first transformation is the reflection, then:  <math>H'(8;16)</math></p> <p>If the first transformation is the reduction, then:  <math>H'(8;4)</math></p>	<p>✓ <math>H'(8;16)</math></p> <p>✓ <math>H'(8;4)</math></p> <p>(2)</p>
1.2.3	<p><math>\frac{\text{Area of original}}{\text{Area of image}} = \frac{1}{k^2}</math></p> <p><math>\therefore \frac{\text{Area KUHLE}}{\text{Area } K''U''H''L''E''} = \frac{1}{\left(\frac{1}{2}\right)^2} = 4</math></p> <p><math>\therefore \text{Area KUHLE}'' : \text{Area } K''U''H''L''E'' = 4:1</math></p>	<p>✓✓ answer</p> <p>(2)</p> <p>[11]</p>

## QUESTION 2

2.1	<p><math>A'(x \cos \theta - y \sin \theta; y \cos \theta + x \sin \theta)</math></p> <p><math>x' = x \cos \theta - y \sin \theta</math></p> <p><math>\therefore x' = 3 \cos 120^\circ - 2 \sin 120^\circ</math></p> <p><math>\therefore x' = 3(-\cos 60^\circ) - 2 \sin 60^\circ</math></p> <p><math>\therefore x' = -3\left(\frac{1}{2}\right) - 2\left(\frac{\sqrt{3}}{2}\right)</math></p> <p><math>\therefore x' = \frac{-3 - 2\sqrt{3}}{2}</math></p> <p><math>y' = y \cos \theta + x \sin \theta</math></p> <p><math>\therefore y' = 2 \cos 120^\circ + 3 \sin 120^\circ</math></p> <p><math>\therefore y' = 2(-\cos 60^\circ) + 3 \sin 60^\circ</math></p> <p><math>\therefore y' = -2\left(\frac{1}{2}\right) + 3\left(\frac{\sqrt{3}}{2}\right)</math></p> <p><math>\therefore y' = \frac{-2 + 3\sqrt{3}}{2}</math></p>	<p>✓ formula</p> <p>✓ simplification</p> <p>✓ substitution</p> <p>✓ answer</p> <p>✓ simplification</p> <p>✓ answer</p> <p>(6)</p>
-----	--	---

<p>2.2</p> <p><math>Q(x; y) \rightarrow Q'(-2; 0)</math></p> <p><math>\therefore Q(x; y) \rightarrow Q'(x \cos 120^\circ - y \sin 120^\circ; y \cos 120^\circ + x \sin 120^\circ)</math></p> <p><math>-2 = x \cos 120^\circ - y \sin 120^\circ</math> and <math>0 = y \cos 120^\circ + x \sin 120^\circ</math></p> <p><math>-2 = x \left(-\frac{1}{2}\right) - y \left(\frac{\sqrt{3}}{2}\right)</math> and <math>0 = y \left(-\frac{1}{2}\right) + x \left(\frac{\sqrt{3}}{2}\right)</math></p> <p><math>\therefore -4 = -x - \sqrt{3}y</math> and <math>0 = -y + \sqrt{3}x</math></p> <p><math>\therefore 4 = x + \sqrt{3}y</math> and <math>y = \sqrt{3}x</math></p> <p><math>\therefore 4 = x + \sqrt{3}(\sqrt{3}x)</math></p> <p><math>\therefore 4 = x + 3x</math></p> <p><math>\therefore 4 = 4x</math></p> <p><math>\therefore x = 1</math></p> <p><math>\therefore y = \sqrt{3}(1) = \sqrt{3}</math></p> <p><math>\therefore Q(1; \sqrt{3})</math></p>	<p><math>\checkmark -2 = x \left(-\frac{1}{2}\right) - y \left(\frac{\sqrt{3}}{2}\right)</math></p> <p><math>\checkmark 0 = y \left(-\frac{1}{2}\right) + x \left(\frac{\sqrt{3}}{2}\right)</math></p> <p><math>\checkmark</math> x-coordinate</p> <p><math>\checkmark</math> y-coordinate</p> <p style="text-align: right;">(4)</p>
---	--

[10]



## TOPIC : FUNCTIONS AND GRAPHS

**Teacher Note:** Functions form a large part of Paper 1 and learners should score well in questions involving this topic. Ensure that they know how to link transformation rules to these graphs.

## LESSON OVERVIEW

1. Introduction session: 5 minutes
2. Typical exam questions:
  - Question 1: 20 minutes
  - Question 2: 20 minutes
3. Discussion of solutions 45 minutes


## SECTION A: TYPICAL EXAM QUESTIONS

## QUESTION 1: 20 minutes

In the diagram, the graphs of the following functions have been sketched:

$$f(x) = a(x + p)^2 + q \quad \text{and} \quad g(x) = \frac{a}{x + p} + q$$

The two graphs intersect at  $A(2; 4)$  and the turning point of the parabola lies at the point of intersection of the asymptotes of the hyperbola. The line  $x = 1$  is the axis of symmetry of the parabola.



- 1.1 Determine the equation of  $f(x)$  in the form  $y = a(x + p)^2 + q$  (3)

- 1.2 Determine the equation of  $g(x)$  in the form  $y = \frac{a}{x+p} + q$  (3)
- 1.3 Write down the range for the graph of  $f$ . (1)
- 1.4 If the graph of  $f$  is shifted 1 unit left and 2 units downwards, write down the equation of the new graph formed. (2)
- 1.5 Write down the values of  $x$  for which  $g(x) \leq 0$  (2)
- [11]

**QUESTION 2: 20 minutes**

Given:  $f(x) = 2(x-1)^2 - 8$  and  $h(x) = 4^x$

- 2.1 Sketch the graphs of  $h$  and  $f$  on the diagram sheet provided. Indicate ALL intercepts with the axes and any turning points. (9)
- 2.2 The graph of  $f$  is shifted 2 units to the LEFT. Write down the equation of the new graph. (1)
- 2.3 Show, algebraically, that  $h\left(x + \frac{1}{2}\right) = 2h(x)$ . (3)
- [13]

**SECTION B: SOLUTIONS AND HINTS TO SECTION A**


**QUESTION 1**

1.1	For the graph of $f(x) = a(x+p)^2 + q$ Substitute the turning point T(1; 2): $y = a(x-1)^2 + 2$ Substitute the point A(2; 4): $4 = a(2-1)^2 + 2$ $\therefore 4 = a + 2$ $\therefore a = 2$ The equation of the parabola is therefore: $y = 2(x-1)^2 + 2$	✓ $y = a(x-1)^2 + 2$ ✓ $a = 2$ ✓ $y = 2(x-1)^2 + 2$ (3)
-----	--	--

1.2	<p>For the graph of <math>g(x) = \frac{a}{x+p} + q</math></p> <p>The vertical asymptote is <math>x=1</math> and the horizontal asymptote is <math>y=2</math>.</p> <p><math>\therefore y = \frac{a}{x-1} + 2</math></p> <p>Substitute the point A(2; 4):</p> <p><math>\therefore 4 = \frac{a}{2-1} + 2</math></p> <p><math>\therefore 4 = a + 2</math></p> <p><math>\therefore a = 2</math></p> <p>The equation of the hyperbola is therefore:</p> <p><math>y = \frac{2}{x-1} + 2</math></p>	<p>✓ <math>y = \frac{a}{x-1} + 2</math></p> <p>✓ <math>a = 2</math></p> <p>✓ <math>y = \frac{2}{x-1} + 2</math></p> <p>(3)</p>
1.3	Range: $y \in [2; \infty)$	<p>✓ <math>y \in [2; \infty)</math></p> <p>(1)</p>
1.4	<p><math>y = 2((x+1)-1)^2 + 2 - 2</math></p> <p><math>\therefore y = 2x^2</math></p>	<p>✓ adding 1 to x and subtracting 2</p> <p>✓ <math>y = 2x^2</math></p> <p>(2)</p>
1.5	<p><math>g(x) \leq 0</math></p> <p><math>\therefore 0 \leq x &lt; 1</math></p>	<p>✓✓ answers</p> <p>(2)</p>

[11]

QUESTION 2

<p>2.1</p>	<p> <math>f(x) = 2(x-1)^2 - 8</math>                      Turning point: <math>(1; -8)</math>                      x-intercepts of parabola:  <math>0 = 2(x-1)^2 - 8</math>  <math>8 = 2(x-1)^2</math>  <math>4 = (x-1)^2</math>  <math>2 = x-1</math> or <math>-2 = x-1</math>  <math>x = 3</math> or <math>x = -1</math>                      y-intercept of parabola: <math>y = 2(0-1)^2 - 8 = -6</math> </p> 	<p>For <math>f(x) = 2(x-1)^2 - 8</math></p> <ul style="list-style-type: none"> <li>✓ turning point</li> <li>✓ shape</li> <li>✓ axis of symmetry</li> <li>✓ y-intercept</li> <li>✓ x-intercepts</li> </ul> <p>For <math>h(x) = 4^x</math></p> <ul style="list-style-type: none"> <li>✓ y-intercept</li> <li>✓ shape</li> <li>✓ coordinates</li> </ul> <p>(9)</p>
<p>2.2</p>	<p> <math>y = 2(x-1+2)^2 - 8</math>  <math>\therefore y = 2(x+1)^2 - 8</math> </p>	<p>✓ <math>y = 2(x+1)^2 - 8</math></p> <p>(1)</p>
<p>2.3</p>	<p> <math>h\left(x + \frac{1}{2}\right)</math>  <math>= 4^{x + \frac{1}{2}} = 4^x \cdot 4^{\frac{1}{2}} = (4^x) \cdot 2 = 2h(x)</math> </p>	<ul style="list-style-type: none"> <li>✓ <math>4^{x + \frac{1}{2}}</math></li> <li>✓ <math>4^x \cdot 4^{\frac{1}{2}}</math></li> <li>✓ <math>(4^x) \cdot 2 = 2h(x)</math></li> </ul> <p>(3)</p>

## SECTION C: HOMEWORK

## QUESTION 1

Given:  $f(x) = \frac{2}{x+1}$

- 1.1 Write down the equations of the asymptotes. (2)
- 1.2 Sketch the graph of  $f$  indicating the coordinates of the  $y$ -intercept as well as the asymptotes. (4)
- 1.3 Write down the equation of the graph formed if the graph of  $f$  is shifted 3 units right and 2 units upwards. (2)
- 1.4 Determine graphically the values of  $x$  for which  $\frac{2}{x+1} \geq 1$  (4)


[12]

## QUESTION 2

In the diagram below, the graphs of the following functions are represented:

$$f(x) = a(x+p)^2 + q \quad \text{and} \quad g(x) = \frac{a}{x+p} + q$$

The turning point of  $f$  is  $(1; 4)$  and the asymptotes of  $g$  intersect at the turning point of  $f$ . Both graphs cut the  $y$ -axis at 3.





- 2.1 Determine the equation of  $f$ . (4)
- 2.2 Determine the equation of  $g$ . (4)
- 2.3 Determine the coordinates of the  $x$ -intercept of  $g$ . (3)
- 2.4 For which values of  $x$  will  $g(x) \leq 0$ ? (2)

[13]



**SECTION D: SOLUTIONS TO HOMEWORK**

**QUESTION 1**

1.1	vertical: $x = -1$ horizontal: $y = 0$	<ul style="list-style-type: none"> <li>✓ vertical</li> <li>✓ horizontal</li> </ul> (2)
1.2		<ul style="list-style-type: none"> <li>✓ left branch</li> <li>✓ right branch</li> <li>✓ coordinates</li> <li>✓ asymptotes</li> </ul> (4)
1.3	$y = \frac{2}{x+1-3} + 2$ $\therefore y = \frac{2}{x-2} + 2$	<ul style="list-style-type: none"> <li>✓ <math>\frac{2}{x+1-3}</math></li> <li>✓ <math>y = \frac{2}{x-2} + 2</math></li> </ul> (2)
1.4	 <p>Therefore <math>\frac{2}{x+1} \geq 1</math> for <math>-1 &lt; x \leq 1</math></p>	<ul style="list-style-type: none"> <li>✓ drawing the line <math>y = 1</math></li> <li>✓ (1; 1)</li> <li>✓✓ <math>-1 &lt; x \leq 1</math></li> </ul> (4)

[12]

## QUESTION 2

2.1	$y = a(x-1)^2 + 4$ Substitute (0 ; 3) $3 = a(0-1)^2 + 4$ $\therefore 3 = a + 4$ $\therefore a = -1$ $\therefore f(x) = -(x-1)^2 + 4$	$\checkmark y = a(x-1)^2 + 4$ $\checkmark 3 = a(0-1)^2 + 4$ $\checkmark a = -1$ $\checkmark f(x) = -(x-1)^2 + 4$ <p style="text-align: right;">(4)</p>
2.2	$y = \frac{a}{x-1} + 4$ Substitute (0 ; 3) $\therefore 3 = \frac{a}{0-1} + 4$ $\therefore 3 = -a + 4$ $\therefore a = 1$ $\therefore g(x) = \frac{1}{x-1} + 4$	$\checkmark \checkmark y = \frac{a}{x-1} + 4$ $\checkmark a = 1$ $\checkmark g(x) = \frac{1}{x-1} + 4$ <p style="text-align: right;">(4)</p>
2.3	$0 = \frac{1}{x-1} + 4$ $\therefore 0 = 1 + 4(x-1)$ $\therefore 0 = 1 + 4x - 4$ $\therefore 0 = 4x - 3$ $\therefore -4x = -3$ $\therefore x = \frac{3}{4}$ $\left(\frac{3}{4}; 0\right)$	$\checkmark 0 = \frac{1}{x-1} + 4$ $\checkmark x = \frac{3}{4}$ $\checkmark \left(\frac{3}{4}; 0\right)$ <p style="text-align: right;">(3)</p>
2.4	$g(x) \leq 0 \text{ for } 0 \leq x < \frac{3}{4}$	$\checkmark x \geq 0$ $\checkmark x < \frac{3}{4}$ <p style="text-align: right;">(2)</p>

[13]

**TOPIC : INVERSE GRAPHS**

**Teacher Note:** Functions form a large part of Paper 1 and learners should score well in questions involving this topic. Ensure that they know how to link transformation rules to these graphs.


**LESSON OVERVIEW**

1. Introduction session            5 minutes
2. Typical exam questions:  
     Question 1:                    40 minutes
3. Discussion of solutions        45 minutes

**SECTION A: TYPICAL EXAM QUESTIONS**

**QUESTION 1:        40 minutes**

Sketched below are the graphs of  $f(x) = 3^x$  and  $g(x) = -x^2$



- 1.1 Write down the equation of the inverse of the graph of  $f(x) = 3^x$  in the form  $f^{-1}(x) = \dots$  (2)
- 1.2 On a set of axes, draw the graph of the inverse of  $f(x) = 3^x$  (2)
- 1.3 Write down the domain of the graph of  $f^{-1}(x)$  (1)
- 1.4 Explain why the inverse of the graph of  $g(x) = -x^2$  is not a function. (1)
- 1.5 Consider the graph of  $g(x) = -x^2$ 
  - 1.5.1 Write down a possible restriction for the domain of  $g(x) = -x^2$  so that the inverse of the graph of  $g$  will now be a function. (1)
  - 1.5.2 Hence draw the graph of the inverse function in 2.5.1 (2)

MATHEMATICS

GRADE 12

SESSION 12

(TEACHER NOTES)

1.6 Explain how, using the transformation of the graph of  $f$ , you would sketch the graphs of:

1.6.1  $h(x) = -\log_3 x$  (2)


1.6.2  $p(x) = \left(\frac{1}{3}\right)^x + 1$  (2)



1.7 Sketch the graph of  $p(x) = \left(\frac{1}{3}\right)^x + 1$  on a set of axes. (3)

[16]

<b>SECTION B: SOLUTIONS AND HINTS TO SECTION A</b>
--

**QUESTION 1**

1.1	$y = 3^x$ $\therefore x = 3^y$ $\therefore \log_3 x = y$ $\therefore f^{-1}(x) = \log_3 x$	✓ $x = 3^y$ ✓ $f^{-1}(x) = \log_3 x$ (2)
1.2		✓ shape ✓ (1; 0) (2)
1.3	Domain: $x \in (0; \infty)$	✓ $x \in (0; \infty)$ (1)
1.4	The inverse is a one-to-many relation, which is not a function.	✓ one-to-many (1)

1.5.1	$x \geq 0$ OR $x \leq 0$	✓ answer (1)
1.5.2		✓✓ shape (2)
1.6.1	$y = 3^x$ $\therefore x = 3^y$ $\therefore \log_3 x = y$ (reflection about the line $y = x$ ) $y = \log_3 x$ $\therefore -y = \log_3 x$ $\therefore y = -\log_3 x$ (reflection about the $x$ -axis)	✓ reflection about $y = x$ ✓ reflection about $x$ -axis (2)
1.6.2	$y = 3^x$ $\therefore y = 3^{-x}$ (Reflection about the $y$ -axis) $\therefore y = \left(\frac{1}{3}\right)^x$ $y = \left(\frac{1}{3}\right)^x + 1$ (translation of 1 unit upwards)	✓ reflection about $y$ -axis ✓ translation (2)
1.7		✓✓ decreasing shape ✓ $y$ -intercept (3)

## SECTION C: HOMEWORK

## QUESTION 1


Consider the functions:  $f(x) = 2x^2$  and  $g(x) = \left(\frac{1}{2}\right)^x$

- 1.1 Restrict the domain of  $f$  in one specific way so that the inverse of  $f$  will also be a function. (1)
  - 1.2 Hence draw the graph of your new function  $f$  and its inverse function  $f^{-1}$  on the same set of axes. (2)
  - 1.3 Write the inverse of  $g$  in the form  $g^{-1}(x) = \dots\dots\dots$  (2)
  - 1.4 Sketch the graph of  $g^{-1}$ . (2)
  - 1.5 Determine graphically the values of  $x$  for which  $\log_{\frac{1}{2}} x < 0$  (1)
- [8]

## QUESTION 2

In the diagram below (not drawn to scale),  
the graph of  $f(x) = 2a^x$ .


The graph of  $f$  passes through the point  $(-1; 4)$  and cuts the  $y$ -axis at A.




- 2.1 Show that  $a = \frac{1}{2}$  and hence write down the equation of  $f$ . (3)
  - 2.2 Determine the coordinates of A. (2)
  - 2.3 Show that the equation of  $g$ , the reflection of  $f$  about the  $y$ -axis, can be written as  $g(x) = 2^{x+1}$  (3)
  - 2.4 Draw a neat sketch graph of  $y = f(x-1) - 2$  indicating the intercepts with the axes as well as the equation of the asymptote. (4)
- [12]

**SECTION D: SOLUTIONS TO HOMEWORK**

**QUESTION 1**

<p>1.1</p>	<p><math>f(x) = 2x^2</math> where <math>x \geq 0</math>                  OR  <math>f(x) = 2x^2</math> where <math>x \leq 0</math></p>	<p>✓ <math>x \geq 0</math> OR <math>x \leq 0</math>                  (1)</p>
<p>1.2</p>	<p>OR</p> 	<p>✓ <math>f</math>                  ✓ <math>f^{-1}</math>                  (2)</p>
<p>1.3</p>	<p><math>y = \left(\frac{1}{2}\right)^x</math>  <math>x = \left(\frac{1}{2}\right)^y</math>  <math>\therefore \log_{\frac{1}{2}} x = y</math>  <math>\therefore g^{-1}(x) = \log_{\frac{1}{2}} x</math></p>	<p>✓ <math>x = \left(\frac{1}{2}\right)^y</math>                  ✓ <math>g^{-1}(x) = \log_{\frac{1}{2}} x</math>                  (2)</p>

1.4		✓ shape ✓ (1;0) (2)
1.5	$\log_{\frac{1}{2}} x < 0$ for $x > 1$	✓ $x > 1$ (1)


[8]

QUESTION 2



2.1	$y = 2a^x$ Substitute $(-1; 4)$ $\therefore 4 = 2a^{-1}$ $\therefore 4 = \frac{2}{a}$ $\therefore 4a = 2$ $\therefore a = \frac{1}{2}$ $\therefore f(x) = 2\left(\frac{1}{2}\right)^x$	✓ $4 = 2a^{-1}$ ✓ $a = \frac{1}{2}$ ✓ $f(x) = 2\left(\frac{1}{2}\right)^x$ (3)
2.2	$y = 2\left(\frac{1}{2}\right)^0 = 2$ $(0; 2)$	✓ $2\left(\frac{1}{2}\right)^0$ ✓ $(0; 2)$ (2)



<p>2.3</p>	$g(x) = 2\left(\frac{1}{2}\right)^{-x}$ $\therefore g(x) = 2(2^x)$ $\therefore g(x) = 2^{1+x}$ $\therefore g(x) = 2^{x+1}$	<ul style="list-style-type: none"> <li>✓ <math>g(x) = 2\left(\frac{1}{2}\right)^{-x}</math></li> <li>✓ <math>g(x) = 2(2^x)</math></li> <li>✓ <math>g(x) = 2^{x+1}</math></li> </ul> <p style="text-align: right;">(3)</p>
<p>2.4</p>		<ul style="list-style-type: none"> <li>✓ y-intercept</li> <li>✓ x-intercept</li> <li>✓ asymptote</li> <li>✓ shape</li> </ul> <p style="text-align: right;">(4)</p>

[12]



## TOPIC 1 : CALCULUS – LIMITS AND FIRST PRINCIPLES

## LESSON OVERVIEW FOR EACH TOPIC

1. Introduction session: 10 minutes
2. Typical exam questions:
  - Question 1: 10 minutes
  - Question 2: 10 minutes
3. Discussion of solutions: 15 minutes

**Teacher Note:** Calculus forms a large part of Paper 1 and learners should score well in questions involving this topic. Ensure that they don't lose marks by using faulty notation.

## SECTION A: TYPICAL EXAM QUESTIONS

## TOPIC 1 : CALCULUS – LIMITS AND FIRST PRINCIPLES

## QUESTION 1: 10 minutes

- 1.1 Given:  $f(x) = -2x^2 + 1$ 
    - 1.1.1 Determine  $f'(x)$  from first principles. (5)
    - 1.1.2 Determine the gradient of the graph at  $x = -2$ , i.e.  $f'(-2)$  (1)
    - 1.1.3 Determine  $f(-2)$ . What does your answer represent? (2)
    - 1.1.4 Determine the average gradient of  $f$  between  $x = -2$  and  $x = 4$  (4)
  - 1.2 Use first principles to determine the derivative of  $f(x) = \frac{1}{x}$  (5)
- [17]

## QUESTION 2: 10 minutes

- 2.1 Differentiate  $f$  by first principles where  $f(x) = x^2 - 2x$ . (5)
  - 2.2 Determine the gradient of the tangent to the graph of  $g(x) = x^3$  at  $x = 3$  (6)
- [11]

TOPIC 2 : CALCULUS – RULES OF DIFFERENTIATION AND TANGENTS

QUESTION 1: 10 minutes

Determine the following and leave your answer with positive exponents:

1.1  $D_x [(2x-3)(x+4)]$  (2)

1.2  $f'(x)$  if  $f(x) = \frac{1}{2\sqrt[4]{x^3}}$  (3)

1.3  $\frac{dy}{dx}$  if  $y = \left(2\sqrt{x} - \frac{1}{3x}\right)^2$  (5)

1.4  $D_x \left[ \frac{1}{\sqrt{x}} (x^3 - 2x^2 + 3x) \right]$  (4)

[14]

QUESTION 2: 10 minutes

2.1 Determine the equation of the tangent to  $f(x) = x^2 - 6x + 5$  at  $x = 2$ . (5)

2.2 Find the equation of the tangent to  $f(x) = 3x^2 - 5x + 1$  which is parallel to the line  $y - 7x + 4 = 0$ . (6)

[11]



## SECTION B: SOLUTIONS AND HINTS TO SECTION A - TOPICS 1&amp;2

## TOPIC 1 : CALCULUS – LIMITS AND FIRST PRINCIPLES

## QUESTION 1

1.1.1	$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{-2(x+h)^2 + 1 - (-2x^2 + 1)}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{-2(x^2 + 2xh + h^2) + 1 + 2x^2 - 1}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{-2x^2 - 4xh - 2h^2 + 1 + 2x^2 - 1}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{h(-4x - 2h)}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} (-4x - 2h)$ $\therefore f'(x) = -4x - 2(0)$ $\therefore f'(x) = -4x$	$\checkmark -2(x+h)^2 + 1$ $\checkmark -(-2x^2 + 1)$ $\checkmark -2x^2 - 4xh - 2h^2$ $\checkmark \frac{h(-4x - 2h)}{h}$ $\checkmark -4x$ <p style="text-align: right;">(5)</p>
1.1.2	$f'(x) = -4x$ $\therefore f'(-2) = -4(-2) = 8$	$\checkmark$ answer (1)
1.1.3	$f(x) = -2x^2 + 1$ $\therefore f(-2) = -2(-2)^2 + 1$ $\therefore f(-2) = -7$ The answer represents the y-value corresponding to $x = -2$	$\checkmark f(-2) = -7$ $\checkmark$ interpretation (2)

<p>1.1.4</p>	$f(x) = -2x^2 + 1$ $f(-2) = -2(-2)^2 + 1$ $\therefore f(-2) = -7$ $f(4) = -2(4)^2 + 1$ $\therefore f(4) = -31$ $(-2; -7) \text{ and } (4; -31)$ $\text{Average gradient} = \frac{-31 - (-7)}{4 - (-2)} = \frac{-24}{6} = -4$	$\checkmark f(-2) = -7$ $\checkmark f(4) = -31$ $\checkmark \frac{-31 - (-7)}{4 - (-2)}$ $\checkmark -4$ <p style="text-align: right;">(4)</p>
<p>1.2</p>	$\therefore f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{\frac{x - (x+h)}{x(x+h)}}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{\frac{x - x - h}{x(x+h)}}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{\frac{-h}{x(x+h)}}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{-h}{x(x+h)} \times \frac{1}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{-1}{x(x+h)}$ $\therefore f'(x) = \frac{-1}{x(x+0)}$ $\therefore f'(x) = -\frac{1}{x^2}$	$\checkmark \frac{\frac{1}{x+h} - \frac{1}{x}}{h}$ $\checkmark \frac{x - (x+h)}{x(x+h)}$ $\checkmark \frac{-h}{x(x+h)}$ $\checkmark \frac{-1}{x(x+h)}$ $\checkmark -\frac{1}{x^2}$ <p style="text-align: right;">(5)</p>

[17]

## QUESTION 2

2.1	$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{(x+h)^2 - 2(x+h) - x^2 + 2x}{h}$ $= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - 2x - 2h - x^2 + 2x}{h}$ $= \lim_{h \rightarrow 0} \frac{2xh + h^2 - 2h}{h}$ $= \lim_{h \rightarrow 0} \frac{h(2x + h - 2)}{h}$ $= \lim_{h \rightarrow 0} (2x - 2 + h)$ $= 2x - 2$	$\checkmark (x+h)^2 - 2(x+h)$ $\checkmark -x^2 + 2x$ $\checkmark \frac{2xh + h^2 - 2h}{h}$ $\checkmark (2x - 2 + h)$ $\checkmark 2x - 2$ <p style="text-align: right;">(5)</p>
2.2	$g'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^3 - x^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{(x+h)(x+h)^2 - x^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{(x+h)(x^2 + 2xh + h^2) - x^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{x^3 + 2x^2h + xh^2 + x^2h + 2xh^2 + h^3 - x^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{3x^2h + 3xh^2 + h^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{h(3x^2 + 3xh + h^2)}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} (3x^2 + 3xh + h^2)$ $\therefore g'(x) = 3x^2$ $\therefore g'(3) = 3(3)^2 = 27$	$\checkmark (x+h)^3 - x^3$ $\checkmark \checkmark 3x^2h + 3xh^2 + h^3$ $\checkmark (3x^2 + 3xh + h^2)$ $\checkmark 3x^2$ $\checkmark 27$ <p style="text-align: right;">(6)</p>

[11]

TOPIC 2 : CALCULUS – RULES OF DIFFERENTIATION AND TANGENTS

QUESTION 1

1.1	$D_x [(2x-3)(x+4)]$ $= D_x [2x^2 + 5x - 12]$ $= 4x + 5$	$\checkmark 2x^2 + 5x - 12$ $\checkmark 4x + 5$ <p style="text-align: right;">(2)</p>
1.2	$f(x) = \frac{1}{2\sqrt[4]{x^3}}$ $\therefore f(x) = \frac{1}{2x^{\frac{3}{4}}}$ $\therefore f(x) = \frac{1}{2}x^{-\frac{3}{4}}$ $\therefore f'(x) = \frac{1}{2} \times -\frac{3}{4}x^{-\frac{3}{4}-1}$ $\therefore f'(x) = -\frac{3}{8}x^{-\frac{7}{4}}$ $\therefore f'(x) = -\frac{3}{8x^{\frac{7}{4}}}$	$\checkmark \frac{1}{2}x^{-\frac{3}{4}}$ $\checkmark -\frac{3}{8}x^{-\frac{7}{4}}$ $\checkmark -\frac{3}{8x^{\frac{7}{4}}}$ <p style="text-align: right;">(3)</p>
1.3	$y = \left(2\sqrt{x} - \frac{1}{3x}\right)^2$ $\therefore y = 4x - \frac{4\sqrt{x}}{3x} + \frac{1}{9x^2}$ $\therefore y = 4x - \frac{4x^{\frac{1}{2}}}{3x} + \frac{1}{9}x^{-2}$ $\therefore y = 4x - \frac{4}{3}x^{-\frac{1}{2}} + \frac{1}{9}x^{-2}$ $\therefore \frac{dy}{dx} = 4 - \frac{4}{3} \times -\frac{1}{2}x^{-\frac{3}{2}} + \frac{1}{9} \times -2x^{-3}$ $\therefore \frac{dy}{dx} = 4 + \frac{2}{3}x^{-\frac{3}{2}} - \frac{2}{9}x^{-3}$ $\therefore \frac{dy}{dx} = 4 + \frac{2}{3x^{\frac{3}{2}}} - \frac{2}{9x^3}$	$\checkmark \text{squaring}$ $\checkmark 4x - \frac{4}{3}x^{-\frac{1}{2}} + \frac{1}{9}x^{-2}$ $\checkmark \checkmark \checkmark 4 + \frac{2}{3x^{\frac{3}{2}}} - \frac{2}{9x^3}$ <p style="text-align: right;">(5)</p>

1.4	$D_x \left[ \frac{1}{\sqrt{x}} (x^3 - 2x^2 + 3x) \right]$ $= D_x \left[ \frac{x^3}{x^{\frac{1}{2}}} - \frac{2x^2}{x^{\frac{1}{2}}} + \frac{3x}{x^{\frac{1}{2}}} \right]$ $= D_x \left[ x^{\frac{5}{2}} - 2x^{\frac{3}{2}} + 3x^{\frac{1}{2}} \right]$ $= \frac{5}{2} x^{\frac{3}{2}} - 3x^{\frac{1}{2}} + \frac{3}{2} x^{-\frac{1}{2}}$ $= \frac{5}{2} x^{\frac{3}{2}} - 3x^{\frac{1}{2}} + \frac{3}{2x^{\frac{1}{2}}}$	$\checkmark x^{\frac{1}{2}}$ $\checkmark D_x \left[ x^{\frac{5}{2}} - 2x^{\frac{3}{2}} + 3x^{\frac{1}{2}} \right]$ $\checkmark \frac{5}{2} x^{\frac{3}{2}} - 3x^{\frac{1}{2}} + \frac{3}{2} x^{-\frac{1}{2}}$ $\checkmark \frac{5}{2} x^{\frac{3}{2}} - 3x^{\frac{1}{2}} + \frac{3}{2x^{\frac{1}{2}}}$ <p style="text-align: right;">(4)</p> <p style="text-align: right;"><b>[14]</b></p>
-----	---	--

## QUESTION 2

2.1	$f(x) = x^2 - 6x + 5$ $x_T = 2$ $y_T = f(2) = (2)^2 - 6(2) + 5$ $\therefore y_T = -3$ $m_T = f'(x) = 2x - 6$ $\therefore f'(2) = 2(2) - 6 = -2$ $y - y_T = m_T(x - x_T)$ $\therefore y - (-3) = -2(x - 2)$ $\therefore y + 3 = -2x + 4$ $\therefore y = -2x + 1$	$\checkmark f(2) = -2$ $\checkmark f'(x) = 2x - 6$ $\checkmark f'(2) = 2(2) - 6 = -2$ $\checkmark y - (-3) = -2(x - 2)$ $\checkmark y = -2x + 1$ <p style="text-align: right;">(5)</p>
2.2	$f(x) = 3x^2 - 5x + 1$ $\therefore f'(x) = 6x - 5$ $y = 7x - 4$ $\therefore 6x - 5 = 7$ $\therefore 6x = 12$ $\therefore x = 2$ $f(2) = 3(2)^2 - 5(2) + 1$ $\therefore f(2) = 3$ $\therefore y - 3 = 7(x - 2)$ $\therefore y - 3 = 7x - 14$ $\therefore y = 7x - 11$	$\checkmark f'(x) = 6x - 5$ $\checkmark 6x - 5 = 7$ $\checkmark x = 2$ $\checkmark f(2) = 3$ $\checkmark y - 3 = 7(x - 2)$ $\checkmark y = 7x - 11$ <p style="text-align: right;">(6)</p> <p style="text-align: right;"><b>[11]</b></p>



**SECTION C: HOMEWORK**

**TOPIC 1 : CALCULUS – LIMITS AND FIRST PRINCIPLES**

**QUESTION 1**

- 1.1 Given:  $f(x) = 1 - \frac{1}{4}x^2$
- 1.1.1 Determine  $f'(x)$  from first principles. (6)
- 1.1.2 Determine the gradient of the graph at  $x = -4$ , i.e.  $f'(-4)$  (1)
- 1.1.3 Determine  $f(-2)$ . What does your answer represent? (2)
- 1.1.4 Determine the average gradient of  $f$  between  $x = -2$  and  $x = 4$  (4)
- 1.2 Use first principles to determine the derivative of  $f(x) = -\frac{3}{x}$  (5)
- [18]

**QUESTION 2**

- 2.1 Differentiate  $f$  by first principles where  $f(x) = -2x$ . (4)
- 2.2 Determine the gradient of the tangent to the graph of  $g(x) = -2x^3$  at  $x = 2$  (6)
- [10]



**TOPIC 2 : CALCULUS – RULES OF DIFFERENTIATION AND TANGENTS**

**QUESTION 1**


Determine the following and leave your answer with positive exponents:

- 1.1  $f'(x)$  if  $f(x) = (4x - 3)^2$  (2)
- 1.2  $D_x \left[ \sqrt[3]{x} + \frac{1}{\sqrt{x}} \right]$  (3)
- 1.3  $D_x \left[ (x^2 - \sqrt{x})^2 \right]$  (5)
- 1.4  $\frac{dy}{dx}$  if  $y = \frac{2x^2 - \sqrt{x} + 5}{\sqrt{x}}$  (4)
- [14]

**QUESTION 2 10 minutes**

- 2.1 Determine the equation of the tangent to the curve  $y = 3x^2 - 2x + 2$  at  $x = -4$ . (5)
- 2.2 The graph of  $f(x) = ax^2 + bx$  passes through the point  $P(3; 6)$  and cuts the  $x$ -axis

at  $(4; 0)$ . Determine the equation of the tangent to  $f$  at  $P$ . (6)



[11]



**SECTION D: SOLUTIONS TO HOMEWORK**

**2. TOPIC 1 : CALCULUS – LIMITS AND FIRST PRINCIPLES**

**QUESTION 1**

<p>1.1.1</p>	$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{1 - \frac{1}{4}(x+h)^2 - \left(1 - \frac{1}{4}x^2\right)}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{1 - \frac{1}{4}(x^2 + 2xh + h^2) - 1 + \frac{1}{4}x^2}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{1 - \frac{1}{4}x^2 - \frac{1}{2}xh - \frac{1}{4}h^2 - 1 + \frac{1}{4}x^2}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{-\frac{1}{2}xh - \frac{1}{4}h^2}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{h\left(-\frac{1}{2}x - \frac{1}{4}h\right)}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \left(-\frac{1}{2}x - \frac{1}{4}h\right)$ $\therefore f'(x) = -\frac{1}{2}x$	$\checkmark 1 - \frac{1}{4}(x+h)^2$ $\checkmark -1 + \frac{1}{4}x^2$ $\checkmark -\frac{1}{4}x^2 - \frac{1}{2}xh - \frac{1}{4}h^2$ $\checkmark -\frac{1}{2}xh - \frac{1}{4}h^2$ $\checkmark \left(-\frac{1}{2}x - \frac{1}{4}h\right)$ $\checkmark -\frac{1}{2}x$ <p style="text-align: right;">(6)</p>
<p>1.1.2</p>	$f'(-4) = -\frac{1}{2}(-4) = 2$	$\checkmark \text{ answer}$ <p>(1)</p>

1.1.3	$f(x) = 1 - \frac{1}{4}x^2$ $\therefore f(-2) = 1 - \frac{1}{4}(-2)^2$ $\therefore f(-2) = 0$ <p>The answer represents the y-value corresponding to <math>x = -2</math></p>	$\checkmark f(-2) = 0$ $\checkmark$ interpretation <p style="text-align: right;">(2)</p>
1.1.4	$f(x) = 1 - \frac{1}{4}x^2$ $f(-2) = 0$ $f(4) = 1 - \frac{1}{4}(4)^2$ $\therefore f(4) = -3$ <p><math>(-2; 0)</math> and <math>(4; -3)</math></p> $\text{Average gradient} = \frac{-3 - 0}{4 - (-2)} = \frac{-3}{6} = -\frac{1}{2}$	$\checkmark f(-2) = -7$ $\checkmark f(4) = -31$ $\checkmark \frac{-31 - (-7)}{4 - (-2)}$ $\checkmark -4$ <p style="text-align: right;">(4)</p>
1.2	$\therefore f'(x) = \lim_{h \rightarrow 0} \frac{-\frac{3}{x+h} - \left(-\frac{3}{x}\right)}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{-\frac{3}{x+h} + \frac{3}{x}}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{\frac{-3x+3(x+h)}{x(x+h)}}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{\frac{-3x+3x+3h}{x(x+h)}}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{\frac{3h}{x(x+h)}}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{3h}{x(x+h)} \times \frac{1}{h}$ $\therefore f'(x) = \lim_{h \rightarrow 0} \frac{3}{x(x+h)}$ $\therefore f'(x) = \frac{3}{x(x+0)}$ $\therefore f'(x) = \frac{3}{x^2}$	$\checkmark -\frac{3}{x+h} - \left(-\frac{3}{x}\right)$ $\checkmark -\frac{3}{x+h} + \frac{3}{x}$ $\checkmark \frac{3h}{x(x+h)}$ $\checkmark \frac{3}{x(x+h)}$ $\checkmark \frac{3}{x^2}$ <p style="text-align: right;">(5)</p>

[18]

QUESTION 2

<p>2.1</p>	$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{-2(x+h) - (-2x)}{h}$ $= \lim_{h \rightarrow 0} \frac{-2x - 2h + 2x}{h}$ $= \lim_{h \rightarrow 0} \frac{-2h}{h}$ $= \lim_{h \rightarrow 0} (-2)$ $= -2$	<p>✓ <math>-2(x+h) - (-2x)</math>                  ✓ <math>-x^2 + 2x</math>                  ✓ <math>\frac{-2h}{h}</math>                  ✓ <math>-2</math></p> <p>(4)</p>
<p>2.2</p>	$g'(x) = \lim_{h \rightarrow 0} \frac{-2(x+h)^3 - (-2x^3)}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{-2(x+h)(x+h)^2 + 2x^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{-2(x+h)(x^2 + 2xh + h^2) + 2x^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{-2(x^3 + 2x^2h + xh^2 + x^2h + 2xh^2 + h^3) + 2x^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{-2(x^3 + 3x^2h + 3xh^2 + h^3) + 2x^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{-2x^3 - 6x^2h - 6xh^2 - 2h^3 + 2x^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{-6x^2h - 6xh^2 - 2h^3}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} \frac{h(-6x^2 - 6xh - 2h)}{h}$ $\therefore g'(x) = \lim_{h \rightarrow 0} (-6x^2 - 6xh - 2h)$ $\therefore g'(x) = -6x^2$ $\therefore g'(2) = -6(2)^2 = -24$	<p>✓ <math>-2(x+h)^3 - (-2x^3)</math>                  ✓✓ <math>-6x^2h - 6xh^2 - 2h^3</math>                  ✓✓ <math>(-6x^2 - 6xh - 2h)</math>                  ✓ <math>-6x^2</math></p> <p>(6)</p>

[10]

## TOPIC 2 : CALCULUS – RULES OF DIFFERENTIATION AND TANGENTS

## QUESTION 1

1.1	$f(x) = (4x - 3)^2$ $\therefore f(x) = 16x^2 - 24x + 9$ $\therefore f'(x) = 16 \times 2x^{2-1} - 24 + 0$ $\therefore f'(x) = 32x - 24$	$\checkmark 16x^2 - 24x + 9$ $\checkmark 32x - 24$ <p style="text-align: right;">(2)</p>
1.2	$D_x \left[ \sqrt[3]{x} + \frac{1}{\sqrt{x}} \right]$ $= D_x \left[ x^{\frac{1}{3}} + x^{-\frac{1}{2}} \right]$ $= \frac{1}{3} x^{-\frac{2}{3}} - \frac{1}{2} x^{-\frac{3}{2}}$ $= \frac{1}{3x^{\frac{2}{3}}} - \frac{1}{2x^{\frac{3}{2}}}$	$\checkmark x^{\frac{1}{3}} + x^{-\frac{1}{2}}$ $\checkmark \frac{1}{3} x^{-\frac{2}{3}} - \frac{1}{2} x^{-\frac{3}{2}}$ $\checkmark \frac{1}{3x^{\frac{2}{3}}} - \frac{1}{2x^{\frac{3}{2}}}$ <p style="text-align: right;">(3)</p>
1.3	$D_x \left[ (x^2 - \sqrt{x})^2 \right]$ $= D_x \left[ x^4 - 2x^2\sqrt{x} + x \right]$ $= D_x \left[ x^4 - 2x^2x^{\frac{1}{2}} + x \right]$ $= D_x \left[ x^4 - 2x^{\frac{5}{2}} + x \right]$ $= 4x^3 - 5x^{\frac{3}{2}} + 1$	$\checkmark \checkmark x^4 - 2x^{\frac{5}{2}} + x$ $\checkmark \checkmark \checkmark 4x^3 - 5x^{\frac{3}{2}} + 1$ <p style="text-align: right;">(5)</p>
1.4	$y = \frac{2x^2 - \sqrt{x} + 5}{\sqrt{x}}$ $\therefore y = \frac{2x^2}{x^{\frac{1}{2}}} - 1 + \frac{5}{x^{\frac{1}{2}}}$ $\therefore y = 2x^{\frac{3}{2}} - 1 + 5x^{-\frac{1}{2}}$ $\therefore \frac{dy}{dx} = 3x^{\frac{1}{2}} - \frac{5}{2}x^{-\frac{3}{2}}$ $\therefore \frac{dy}{dx} = 3x^{\frac{1}{2}} - \frac{5}{2x^{\frac{3}{2}}}$	$\checkmark x^{\frac{1}{2}}$ $\checkmark 2x^{\frac{3}{2}} - 1 + 5x^{-\frac{1}{2}}$ $\checkmark 3x^{\frac{1}{2}} - \frac{5}{2}x^{-\frac{3}{2}}$ $\checkmark 3x^{\frac{1}{2}} - \frac{5}{2x^{\frac{3}{2}}}$ <p style="text-align: right;">(4)</p>

[14]

QUESTION 2

<p>2.1</p>	$y = f(x) = 3x^2 - 2x + 2$ $x_T = -4$ $\therefore f(-4) = 3(-4)^2 - 2(-4) + 2$ $\therefore f(-4) = 58$ $\therefore y_T = 58$ $m_T = f'(x) = 6x - 2$ $\therefore f'(-4) = 6(-4) - 2$ $\therefore f'(5) = -26$ $y - y_T = m_t(x - x_T)$ $\therefore y - 58 = -26(x - (-4))$ $\therefore y - 58 = -26x - 104$ $\therefore y = -26x - 46$	$\checkmark f(-4) = 58$ $\checkmark f'(x) = 6x - 2$ $\checkmark f'(5) = -26$ $\checkmark y - 58 = -26(x - (-4))$ $\checkmark y = -26x - 46$ <p style="text-align: right;">(5)</p>
<p>2.2</p>	$y = a(x - 0)(x - 4)$ <p>Substitute the point (3; 6)</p> $\therefore 6 = a(3 - 0)(3 - 4)$ $\therefore 6 = -3a$ $\therefore a = -2$ $\therefore y = -2(x - 0)(x - 4)$ $\therefore y = -2x(x - 4)$ $\therefore y = -2x^2 + 8x$ $f(x) = -2x^2 + 8x$ $\therefore f'(x) = -4x + 8$ $\therefore f'(3) = -4(3) + 8 = -4$ $y - y_T = m_t(x - x_T)$ $\therefore y - 6 = -4(x - 3)$ $\therefore y - 6 = -4x + 12$ $\therefore y = -4x + 18$	$\checkmark y = a(x - 0)(x - 4)$ $\checkmark a = -2$ $\checkmark f(x) = -2x^2 + 8x$ $\checkmark f'(x) = -4x + 8$ $\checkmark f'(3) = -4(3) + 8 = -4$ $\checkmark y = -4x + 18$ <p style="text-align: right;">(6)</p>

[11]

**TOPIC 1 : CALCULUS – GRAPHICAL APPLICATIONS**

**Teacher Note: TOPIC 1:** Cubic graphs are easy to sketch and are worth a lot of marks in the matric exam. Make sure the learners learn how to sketch these graphs.

**TOPIC 2:** Linear Programming requires learners to be able to translate from words into maths, which is quite a challenge. However, if the words “at least”, “at most”, “may not exceed” and “must not be more than” are fully explained and understood, then this topic can be well answered. Revise the linear function in detail before approaching this topic.

**LESSON OVERVIEW FOR EACH TOPIC**

1. Introduction session: 10 minutes
2. Typical exam questions:
  - Question 1: 10 minutes
  - Question 2: 10 minutes
3. Discussion of solutions: 15 minutes

**SECTION A: TYPICAL EXAM QUESTIONS****TOPIC 1 : CALCULUS – GRAPHICAL APPLICATIONS****QUESTION 1: 10 minutes**

Sketch the graph of  $f(x) = 2x^3 - 6x - 4$

[17]

**QUESTION 2: 10 minutes**

$(2; 9)$  is a turning point on the graph of  $f(x) = ax^3 + 5x^2 + 4x + b$ . Determine the value of  $a$  and  $b$  and hence the equation of the cubic function.

[7]

**3. TOPIC 2: LINEAR PROGRAMMING****QUESTION 1: 20 minutes**

A clothing company manufactures white shirts and grey trousers for schools.


- A minimum of 200 shirts must be manufactured daily.
- In total, not more than 600 pieces of clothing can be manufactured daily.
- It takes 50 machine minutes to manufacture a shirt and 100 machine minutes to manufacture a pair of trousers.
- There are at most 45 000 machine minutes available per day.



Let the number of white shirts manufactured in a day be  $x$ .

Let the number of pairs of grey trousers manufactured in a day be  $y$ .

- 1.1 Write down the constraints, in terms of  $x$  and  $y$ , to represent the above information.  
(You may assume:  $x \geq 0$ ,  $y \geq 0$ ) (3)
- 1.2 Use the diagram provided on the next page to represent the constraints graphically. (5)
- 1.3 Clearly indicate the feasible region by shading it. (1)
- 1.4 If the profit is R30 for a shirt and R40 for a pair of trousers, write down the equation indicating the profit in terms of  $x$  and  $y$ . (2)
- 1.5 Using a search line and your graph, determine the number of shirts and pairs of trousers that will yield a maximum daily profit. (3)




[14]

## SECTION B: SOLUTIONS AND HINTS TO SECTION A

## TOPIC 1 : CALCULUS – GRAPHICAL APPLICATIONS

## QUESTION 1

<p><b>y-intercept:</b> <math>(0; -4)</math></p> <p><b>x-intercepts:</b></p> $0 = 2x^3 - 6x - 4$ $\therefore 0 = x^3 - 3x - 2$ $\therefore 0 = (x+1)(x^2 - x - 2) \quad (\text{using the factor theorem})$ $\therefore 0 = (x+1)(x-2)(x+1)$ $\therefore x = -1 \text{ or } x = 2$ $(-1; 0) \quad (2; 0)$ <p><b>Stationary points:</b></p> $f(x) = 2x^3 - 6x - 4$ $\therefore f'(x) = 6x^2 - 6$ $\therefore 0 = 6x^2 - 6 \quad (\text{At a turning point, } f'(x) = 0)$ $\therefore 0 = x^2 - 1$ $\therefore x = \pm 1$ $f(1) = -8$ $f(-1) = 0$ <p>Turning points are <math>(1; -8)</math> and <math>(-1; 0)</math></p> <p><b>Point of inflection:</b></p> $f'(x) = 6x^2 - 6$ $\therefore f''(x) = 12x$ $\therefore 0 = 12x$ $\therefore x = 0$ $f(0) = -4$ <p>Point of inflection at <math>(0; -4)</math></p> <p>Alternatively: The x-coordinate of the point of inflection can be determined by adding the x-coordinates of the turning points and then dividing the result by 2.</p> $x = \frac{(1) + (-1)}{2} = 0$	<p>✓ <math>(0; -4)</math></p> <p>✓ <math>0 = 2x^3 - 6x - 4</math></p> <p>✓ <math>0 = (x+1)(x^2 - x - 2)</math></p> <p>✓ <math>0 = (x+1)(x-2)(x+1)</math></p> <p>✓ <math>(-1; 0) \quad (2; 0)</math></p> <p>✓ <math>f'(x) = 6x^2 - 6</math></p> <p>✓ <math>0 = 6x^2 - 6</math></p> <p>✓ <math>x = \pm 1</math></p> <p>✓ <math>(1; -8)</math> and <math>(-1; 0)</math></p> <p>✓ <math>f''(x) = 12x</math></p> <p>✓ <math>(0; -4)</math></p> <p>✓ <math>\frac{(1) + (-1)}{2}</math></p> <p>✓ <math>x = 0</math></p>
--	---



	The graph is represented above	<ul style="list-style-type: none"> <li>✓ intercepts with the axes</li> <li>✓ turning points</li> <li>✓ shape</li> <li>✓ point of inflection</li> </ul> <p style="text-align: right;"><b>[17]</b></p>
--	--------------------------------	--


**QUESTION 2**

	<p>At the turning point (2; 9) , we know that</p> $f'(2) = 0$ $f(x) = ax^3 + 5x^2 + 4x + b$ $\therefore f'(x) = 3ax^2 + 10x + 4$ $\therefore f'(2) = 3a(2)^2 + 10(2) + 4$	<ul style="list-style-type: none"> <li>✓ <math>f'(x) = 3ax^2 + 10x + 4</math></li> <li>✓ <math>f'(2) = 12a + 24</math></li> <li>✓ <math>a = -2</math></li> <li>✓ <math>y = -2x^3 + 5x^2 + 4x + b</math></li> <li>✓ <math>9 = -2(2)^3 + 5(2)^2 + 4(2) + b</math></li> <li>✓ <math>b = -3</math></li> <li>✓ <math>f(x) = -x^3 + 5x^2 + 4x - 3</math></li> </ul> <p style="text-align: right;"><b>[7]</b></p>
--	---	--

$\therefore f'(2) = 12a + 24$ $\therefore 0 = 12a + 24$ $\therefore -12a = 24$ $\therefore a = -2$ <p>We can now substitute <math>a = -2</math> into the original equation:</p> $y = -2x^3 + 5x^2 + 4x + b$ <p>In order to get the value of <math>b</math>, substitute the point <math>(2; 9)</math> into this equation:</p> $\therefore 9 = -2(2)^3 + 5(2)^2 + 4(2) + b$ $\therefore 9 = -16 + 20 + 8 + b$ $\therefore 9 = 12 + b$ $\therefore -b = 3$ $\therefore b = -3$ <p>The equation of the cubic function is therefore</p> $f(x) = -x^3 + 5x^2 + 4x - 3$	
--	--

**TOPIC 2 : LINEAR PROGRAMMING****QUESTION 1**

1.1	$x \geq 200$ $x + y \leq 600$ $50x + 100y \leq 45000$	$\checkmark x \geq 200$ $\checkmark x + y \leq 600$ $\checkmark 50x + 100y \leq 45000$ <div style="text-align: right;">(3)</div>
1.2	See diagram	See diagram for mark allocation <div style="text-align: right;">(5)</div>
1.3	See diagram	See diagram for mark allocation <div style="text-align: right;">(1)</div>



1.4	$P = 30x + 40y$	✓ $30x$ ✓ $40y$ (2)
1.5	$30x + 40y = P$ $\therefore 40y = -30x + P$ $\therefore y = -\frac{3}{4}x + \frac{P}{40}$ <b>3 cuts on y-axis</b> <b>4 cuts on x-axis</b> <b>Maximum at (300; 300)</b>	✓ $y = -\frac{3}{4}x + \frac{P}{40}$ ✓ search line ✓ (300; 300) (3)

[14]

## SECTION C: HOMEWORK

## TOPIC 1 : CALCULUS – GRAPHICAL APPLICATIONS

## QUESTION 1


Sketch the graph of  $f(x) = x^3 - 3x^2 + 4$

Indicate the coordinates of the stationary points, intercepts with the axes and any points of inflection. [15]

## QUESTION 2

(DoE Feb. 2009 Paper 1)

The graph of  $h(x) = -x^3 + ax^2 + bx$  is shown below.  $A(-1; -3,5)$  and  $B(2; 10)$  are the turning points of  $h$ . The graph passes through the origin and further cuts the  $x$ -axis at  $C$  and  $D$ .



- 2.1 Show that  $a = \frac{3}{2}$  and  $b = 6$  (7)
- 2.2 Calculate the average gradient between A and B. (2)
- 2.3 Determine the equation of the tangent to  $h$  at  $x = -2$ . (5)
- 2.4 Determine the  $x$ -value of the point of inflection of  $h$ . (3)

[17]

**TOPIC 2 : LINEAR PROGRAMMING**

**QUESTION 1**

A chemical solution is made by mixing two chemicals, P and Q into a solution called S. The solution S requires at least 900kg but not more than 1400kg of the chemicals. The solution must contain at least 2kg of P to every kg of Q. Write down the constraints and then sketch the feasible region. Find the mixture that will be the most cost effective if P costs R5 per kg and Q costs R3 per kg. [13]

**QUESTION 2**

In order to paint the walls of his home, Joseph will require at least 10 litres of purple paint. Purple paint is obtained by mixing quantities of red and blue paint. To obtain a suitable shade of purple paint, the volume of blue paint used must be at least half the volume of red paint used. The hardware store where Joseph intends buying the paint, has only 8 litres of blue paint in stock. Let the number of litres of red paint be  $x$  and the number of litres of blue paint be  $y$ .


- 2.1 Write down the inequalities in terms of  $x$  and  $y$  which represent the constraints of this situation. (3)
  - 2.2 On the attached diagram provided, represent the constraints graphically and clearly indicate the feasible region. (4)
  - 2.3 The cost of both red and blue paint is R40 per litre, but the paint is only sold in 2-litre tins. Determine the number of litres of red and blue paint which can be bought maintaining a minimum cost. Show all possible combinations. (5)
- [12]

**SECTION D: SOLUTIONS TO HOMEWORK**

**4. TOPIC 1 : CALCULUS – GRAPHICAL APPLICATIONS**

**QUESTION 1**

<p>x-intercepts:  <math>0 = x^3 - 3x^2 + 4</math>  <math>\therefore (x+1)(x^2 - 4x + 4) = 0</math>  <math>\therefore (x+1)(x-2)(x-2) = 0</math>  <math>\therefore x = -1</math> or <math>x = 2</math>  <math>f'(x) = 3x^2 - 6x</math>  <math>\therefore 0 = 3x^2 - 6x</math>  <math>\therefore 0 = x^2 - 2x</math>  <math>\therefore 0 = x(x-2)</math>  <math>\therefore x = 0</math> or <math>x = 2</math></p>	<p>y-intercept: 4</p> <p>✓ <math>0 = x^3 - 3x^2 + 4</math>                  ✓ <math>(x+1)(x^2 - 4x + 4) = 0</math>                  ✓ <math>x = -1</math> or <math>x = 2</math></p> <p>✓ <math>f'(x) = 3x^2 - 6x</math>                  ✓ <math>0 = 3x^2 - 6x</math>                  ✓ <math>x = 0</math> or <math>x = 2</math></p>	
---	---	--

	<p>For <math>x=0</math> <math>f(0) = (0)^3 - 3(0)^2 + 4 = 4</math>  Max turning point at <math>(0; 4)</math></p> <p>For <math>x=2</math> <math>f(2) = (2)^3 - 3(2)^2 + 4 = 0</math>  Min turning point at <math>(2; 0)</math></p> <p><math>f'(x) = 3x^2 - 6x</math>  <math>\therefore f''(x) = 6x - 6</math>  <math>\therefore 0 = 6x - 6</math>  <math>\therefore -6x = -6</math>  <math>\therefore x = 1</math></p> <p><math>f(1) = (1)^3 - 3(1)^2 + 4</math>  <math>f(1) = 2</math></p> <p>The point of inflection is <math>(1; 2)</math></p> 	<ul style="list-style-type: none"> <li>✓ <math>(0; 4)</math></li> <li>✓ <math>(2; 0)</math></li> <li>✓ <math>f''(x) = 6x - 6</math></li> <li>✓ <math>x = 1</math></li> <li>✓ <math>(1; 2)</math></li> <li>✓ intercepts with the axes</li> <li>✓ turning points</li> <li>✓ shape</li> <li>✓ point of inflection</li> </ul> <p style="text-align: right;">(15)</p>
--	--	--

## QUESTION 2

2.1	<p><math>h'(x) = -3x^2 + 2ax + b</math></p> <p><math>h'(-1) = -3(-1)^2 + 2a(-1) + b</math></p> <p><math>0 = -3 - 2a + b</math></p> <p><math>2a - b = -3 \quad \dots (i)</math></p> <p><math>h'(2) = -3(2)^2 + 2a(2) + b</math></p> <p><math>0 = -12 + 4a + b</math></p> <p><math>4a + b = 12 \quad \dots (ii)</math></p>	<ul style="list-style-type: none"> <li>✓ <math>h'(x) = -3x^2 + 2ax + b</math></li> <li>✓ <math>h'(-1) = -3(-1)^2 + 2a(-1) + b</math></li> <li>✓ <math>2a - b = -3</math></li> <li>✓ <math>h'(2) = -3(2)^2 + 2a(2) + b</math></li> <li>✓ <math>4a + b = 12</math></li> <li>✓ <math>a = \frac{3}{2}</math></li> <li>✓ <math>b = 6</math></li> </ul> <p style="text-align: right;">(7)</p>
-----	--	---



	$6a = 9$ (i) + (ii) $\therefore a = \frac{3}{2}$ $\therefore 2\left(\frac{3}{2}\right) - b = -3$ $b = 6$	
2.2	Average gradient $= \frac{10 - (-3,5)}{2 - (-1)}$ $= \frac{13,5}{3}$ $= \frac{9}{2}$	$\checkmark \frac{10 - (-3,5)}{2 - (-1)}$ $\checkmark \frac{9}{2}$ (2)
2.3	$h(x) = -x^3 + \frac{3}{2}x^2 + 6x$ $\therefore h'(x) = -3x^2 + 3x + 6$ $h'(-2) = -3(-2)^2 + 3(-2) + 6$ $h'(-2) = -12$ Point of contact $(-2; 2)$ $y - 2 = -12(x + 2)$ $y = -12x - 22$	$\checkmark h(x) = -x^3 + \frac{3}{2}x^2 + 6x$ $\checkmark h'(x) = -3x^2 + 3x + 6$ $\checkmark h'(-2) = -12$ $\checkmark y = -12x - 22$ $\checkmark h'(-2) = -12$ (5)
2.4	$h'(x) = -3x^2 + 3x + 6$ $h''(x) = -6x + 3$ $-6x + 3 = 0$ $x = \frac{1}{2}$	$\checkmark h''(x) = -6x + 3$ $\checkmark -6x + 3 = 0$ $\checkmark x = \frac{1}{2}$ (3)

[17]

TOPIC 2 : LINEAR PROGRAMMING

QUESTION 1

Let  $x$  = chemical P and  $y$  = chemical Q

$$x + y \geq 900$$

$$x + y \leq 1400$$

$$x \geq 2y$$

$$\therefore -2y \geq -x$$

$$\therefore y \leq \frac{1}{2}x$$

The objective function is  $C = 5x + 3y$

$$\therefore 5x + 3y = C$$

$$\therefore 3y = -5x + C$$

$$\therefore y = -\frac{5}{3}x + \frac{C}{3}$$

Cut on  $y$ -axis is 5    Cut on  $x$ -axis is 3

The most cost effective mixture is the point at which the cost is a minimum. The minimum cost is at point B (600 ; 300).

Therefore, the most cost effective mixture is:  
600kg of P and 300kg of Q.

$$\checkmark x + y \geq 900$$

$$\checkmark x + y \leq 1400$$

$$\checkmark x \geq 2y$$

$$\checkmark y \leq \frac{1}{2}x$$

$$\checkmark C = 5x + 3y$$


$$\checkmark y = -\frac{5}{3}x + \frac{C}{3}$$

$$\checkmark 600\text{kg of P}$$

$$\checkmark 300\text{kg of Q}$$


See graph for other marks

[13]



QUESTION 2

2.1	$x + y \geq 10$ $y \geq \frac{1}{2}x$ $y \leq 8$	$\checkmark x + y \geq 10$ $\checkmark y \geq \frac{1}{2}x$ $\checkmark y \leq 8$	(3)
2.2	see below		
2.3	$C = 40x + 40y$ $\therefore 40x + 40y = C$ $\therefore 40y = -40x + C$ $\therefore y = -1x + \frac{C}{40}$	$\checkmark C = 40x + 40y$ $\checkmark$ search line on diagram $\checkmark (2;8)$ $\checkmark (4;6)$ $\checkmark (6;4)$	(5)



(4)  
[12]