

# SENIOR SECONDARY IMPROVEMENT PROGRAMME 2013



**education**

Department: Education

**GAUTENG PROVINCE**

## GRADE 12



## MATHEMATICS

## TEACHER NOTES

The SSIP is supported by



# TABLE OF CONTENTS

## TEACHER NOTES

SESSION	TOPIC	PAGE
16	Data Handling	3 - 21
	Transformations	22 - 32
17	Functions	33 - 44
	Calculus	45 - 53
18	Linear Programming	54 - 67
	Trigonometry	
19	2D Trigonometry	68 - 74
	3D Trigonometry	75 - 86



## SESSION 16.1

### TOPIC: DATA HANDLING

**Teacher Note:** Data Handling makes up approximately 20% of Paper 2. Based on the analysis of the November 2011 exam, this is a section that learners generally perform well in. However, there are still key concepts that need attention. The following should be emphasised during this session: language (terminology used in data questions), use of the STAT mode on the calculator and how to interpret results and communicate conclusions. This session is designed in particular to help learners understand how to apply what they have learnt in grade 11 to answer questions regarding best fit and distribution of data. It is important that learners understand that it is crucial that they are able to interpret a set of data and communicate that.

### LESSON OVERVIEW

**Teacher Note:** Remind learners of the concepts that they need to know from grade 11. The following concepts need to be understood: mean, mode, quartiles, range, interquartile range, five number summary, ogives, Box and Whisker plots, variance and standard deviation.

It is also important to remind learners of the formulae given on the formula sheet. Learners must familiarise themselves with that sheet.

Provide learners with tips regarding neatness, layout, notation and mark allocations.

- |                            |        |
|----------------------------|--------|
| 1. Introduction session:   | 5 min  |
| 2. Typical exam questions: |        |
| Question 1:                | 15 min |
| Question 2:                | 10 min |
| Question 3:                | 10 min |
| Question 4:                | 5 min  |
| 3 Discussion:              | 5 min  |

### SECTION A: TYPICAL EXAM QUESTIONS

#### QUESTION 1

The ages of the final 23 players selected by coach Carlos Perreira to play for Bafana Bafana in the 2010 FIFA World Cup are provided on the following page.

Position	Player	Age
1	Shu-Aib Walters	28
2	Siboniso Gaxa	26
3	Tshepo Masilela	25
4	Aaron Moekoena	29
5	Lucas Thwala	28
6	Macbeth Sibaya	32
7	Lance Davids	25
8	Siphiwe Tshabalala	25
9	Katlego Mphela	25
10	Steven Pienaar	28
11	Teko Modise	27
12	Reneilwe Letsholonyane	28
13	Kagisho Dikgacoi	25
14	Matthew Booth	33
15	Bernard Parker	24
16	Itumeleng Khune	22
17	Surprise Moriri	30
18	Siyabonga Nomvetho	32
19	Anele Ngcongca	22
20	Bongani Khumalo	23
21	Siyabonga Sangweni	28
22	Moeneeb Josephs	30
23	Thanduyise Khuboni	24



Source: [www.2010FifaWorldCup.com](http://www.2010FifaWorldCup.com):final squads – [MediaClubSouthAfrica.com](http://MediaClubSouthAfrica.com)


The ages of the players are to be grouped into class intervals.

(a) Complete the following table:

(2)

Class intervals (ages)	Frequency	Cumulative frequency
$16 \leq x < 20$		
$20 \leq x < 24$		
$24 \leq x < 28$		
$28 \leq x < 32$		
$32 \leq x < 36$		

- (b) On the diagram provided below, draw a cumulative frequency curve for this data. (6)



- (c) Use your graph to read off approximate values for the quartiles. (3)  
 [11]

**QUESTION 2**

- (a) Complete the table and then use the table to calculate the standard deviation. (5)

Class intervals	Frequency ( $f$ )	Midpoint ( $m$ )	$f \times m$	$m - \bar{x}$	$(m - \bar{x})^2$	$f \times (m - \bar{x})^2$
$20 \leq x < 24$	3	22				
$24 \leq x < 28$	9	26				
$28 \leq x < 32$	8	30				
$32 \leq x < 36$	3	34				
			$\bar{x} =$			

- (b) Hence calculate the standard deviation using the table. (2)
- (c) Now use your calculator to verify your answer. (2)


[9]

**QUESTION 3**

The table below represents the number of people infected with malaria in a certain area from 2001 to 2006:

YEAR	NUMBER OF PEOPLE INFECTED
2001	117
2002	122
2003	130
2004	133
2005	135
2006	137

- (a) Draw a scatter plot to represent the above data. Use the diagram provided below. (2)




- (b) Explain whether a linear, quadratic or exponential curve would be a line of best fit for the above-mentioned data. (1)
- (c) If the same trend continued, estimate, by using your graph, the number of people who will be infected with malaria in 2008. (1)  
[4]

**QUESTION 4**

A medical researcher recorded the growth in the number of bacteria over a period of 10 hours. The results are recorded in the following table:

Time in hours	0	1	2	3	4	5	6	7	8	9	10
Number of bacteria	5	10	7	13	10	20	30	35	45	65	80

- (a) On the diagram provided below, draw a scatter plot to represent this data. (2)
- (b) State the type of relationship (linear, quadratic or exponential) that exists between the number of hours and the growth in the number of bacteria. (1)




(3)  
[6]

### QUESTION 5

The duration of telephone calls made by a receptionist was monitored for a week. The data obtained is represented by the normal distribution curve on the following page. The mean time was 176 seconds with a standard deviation of 30 seconds.

- (a) What percentage of calls made was between 146 seconds and 206 seconds in duration? Fill in the necessary information on the graph provided below. (2)
- (b) Determine the time interval for the duration of calls for the middle 95% of the data. (2)
- (c) What percentage of calls made were in excess of 146 seconds? (2)



[6]

**SECTION D: SOLUTIONS AND HINTS TO SECTION A**

**QUESTION 1**


(a)

Class intervals (ages)	Frequency ✓	Cumulative frequency ✓
$16 \leq x < 20$	0	0
$20 \leq x < 24$	3	3
$24 \leq x < 28$	9	12
$28 \leq x < 32$	8	20
$32 \leq x < 36$	3	23

(2)

(b)

Class intervals (ages)	Frequency	Cumulative frequency	Graph points
$16 \leq x < 20$	0	0	(20 ; 0)
$20 \leq x < 24$	3	3	(24 ; 3)
$24 \leq x < 28$	9	12	(28 ; 12)
$28 \leq x < 32$	8	20	(32 ; 20)
$32 \leq x < 36$	3	23	(36 ; 23)



**Teacher Note:** recommend that learners write down the co-ordinates. X coordinate is the last number in the interval and y value is the cumulative frequency value.

**Teacher Note:** highlight that a cumulative frequency graph can also be referred to as an ogive.

(6)



(c)

<p><b>Lower quartile</b></p> $23 \times \frac{1}{4} = 5.75$ <p>(5,75;25) Therefore <math>Q_1 = 25</math> ✓</p> <p><b>Median</b></p> $23 \times \frac{1}{2} = 11.5$ <p>(11.5;28) Therefore Median = 28 ✓</p> <p><b>Upper quartile</b></p> $23 \times \frac{3}{4} = 17.25$ <p>(17.25;31) Therefore <math>Q_3 = 31</math> ✓</p>	<p><b>Teacher Note:</b> remind learners that the lower quartile is the 25<sup>th</sup> percentile. Therefore, multiply the cumulative frequency by a quarter and read off the graph to determine the y-value. Similarly the median is the 50<sup>th</sup> percentile. Therefore, multiply the cumulative frequency by a half. The upper quartile is determined by multiplying the cumulative frequency by three quarters (the 75<sup>th</sup> percentile)</p> <p style="text-align: right;">(3)</p>
--	---

[11]

**QUESTION 2**

(a)



Class intervals	Frequency (f)	Midpoint (m)	$f \times m$ ✓	$m - \bar{x}$ ✓	$(m - \bar{x})^2$ ✓	$f \times (m - \bar{x})^2$ ✓
$20 \leq x < 24$	3	22	66	-5,9	34,81	104,43
$24 \leq x < 28$	9	26	234	-1,9	3,61	32,49
$28 \leq x < 32$	8	30	240	2,1	4,41	35,28
$32 \leq x < 36$	3	34	102	6,1	37,21	111,63
			$\bar{x} = \frac{642}{23} = 27,9$ ✓			$\sum f \times (m - \bar{x})^2 = 283,83$

(5)

**Teacher Note:** emphasise that mean can be calculated by multiplying the midpoint of the interval and relative frequency.


<p>(b) <math>SD = \sqrt{\frac{\sum f \cdot (x - \bar{x})^2}{23}} = \sqrt{\frac{283,83}{23}} = 3,5</math></p>	<p>✓✓</p> <p style="text-align: right;">(2)</p>
--	---

<p><b>CASIO fx-82ES PLUS:</b></p> <p>MODE</p> <p>2 : STAT</p> <p>1 : 1 – VAR</p> <p>SHIFT SETUP</p> <p>3: STAT (you need to scroll down to get this function)</p> <p>1: ON</p> <p>Enter the midpoints: 22= 26= 30= 34=</p> <p>Enter the frequencies: 3= 9= 8= 3=</p> <p>AC SHIFT 1</p> <p>4: VAR</p> <p>3 : <math>x\sigma n =</math></p> <p>The answer will read:3,5</p> <p><b>SHARP DAL:</b></p> <p>MODE 1=</p> <p>Enter data:</p> <p>22 STO 3 M+</p> <p>26 STO 9 M+</p> <p>30 STO 8 M+</p> <p>34 STO 3 M+</p> <p>RCL 6 to get 3,5</p>	<p>✓✓</p> <p>(2)</p>
---	----------------------

[9]

QUESTION 3

(a)




(2)

b. Linear	✓
c. Approx. 145 people	✓

[4]


QUESTION 4



b. Quadratic or exponential	✓ <b>Teacher Note:</b> remind learners that if ever asked to describe or state the relationship, they should answer by identifying the function, e.g. linear, quadratic, exponential.
-----------------------------	--

[3]

QUESTION 5



One standard deviation interval:

$$\begin{aligned}
 &(\bar{x} - s; \bar{x} + s) \\
 &= (176 - 30; 176 + 30) \\
 &= (146; 206)
 \end{aligned}$$

Two standard deviation intervals:

$$\begin{aligned}
 &(\bar{x} - s; \bar{x} + s) \\
 &= (176 - 2(30); 176 + 2(30)) \\
 &= (116; 236)
 \end{aligned}$$

Three standard deviation intervals:

$$\begin{aligned}
 &(\bar{x} - s; \bar{x} + s) \\
 &= (176 - 3(30); 176 + 3(30)) \\
 &= (86; 266)
 \end{aligned}$$



<p>a. The interval between 146 seconds and 206 seconds lies between one standard deviation of the mean. For the normal distribution, approximately 68% of the data lies between one standard deviation of the mean.</p>	<p>✓✓</p>
<p>b. The middle 95% of the data for a normal distribution lies between two standard deviations on either side of the mean. The middle 95% of the calls will be between 116 and 236 seconds.</p>	<p>✓✓</p>
<p>c. Approximately 34% of the calls are between 146 and 176 seconds. Another 49,85% of the calls are in excess of 176 seconds. Therefore, in total, approximately 84% of the calls are in excess of 146 seconds.</p>	<p>✓✓</p>

[6]

**SECTION C: HOMEWORK**

**QUESTION 1**

The ages of the final 23 players selected by coach Oscar Tabarez to play for Uruguay in the 2010 FIFA World Cup are provided below.

Position	Player	Age
1	Fernando Musiera	23
2	Diego Lugano (captain)	29
3	Diego Godin	24
4	Jorge Fucile	25
5	Walter Gargano	25
6	Andres Scotti	35
7	Edinson Cavani	23
8	Sebastian Eguren	29
9	Luis Suarez	23
10	Diego Forlan	31
11	Alvaro Pereira	25
12	Juan Castillo	32
13	Sebastian Abreu	33
14	Nicolas Lodeira	23
15	Diago Perez	30
16	Maxi Perrreira	26
17	Ignacio Gonzales	28
18	Egidio Arevalo Rios	27
19	Sebastian Fernandes	25
20	Mauricio Victorino	27
21	Alvaro Fernandez	24
22	Martin Caceres	23
23	Martin Silva	27




Source: [www.2010FifaWorldCup.finalsquads – MediaClubSouthAfrica.com](http://www.2010FifaWorldCup.finalsquads-MediaClubSouthAfrica.com)

(a) Complete the following table: (2)

Class intervals (ages)	Frequency	Cumulative frequency
$16 \leq x < 20$		
$20 \leq x < 24$		
$24 \leq x < 28$		
$28 \leq x < 32$		
$32 \leq x < 36$		

(b) On the diagram provided below, draw an ogive representing the above data.

(6)



(c) Use your graph to read off approximate values for the quartiles.

(3)

[11]

**QUESTION 2**

(a) Complete the table and then use the table to calculate the standard deviation.

(5)

Class intervals	Frequency ( $f$ )	Midpoint ( $m$ )	$f \times m$	$m - \bar{x}$	$(m - \bar{x})^2$	$f \times (m - \bar{x})^2$
$20 \leq x < 24$	3	22				
$24 \leq x < 28$	9	26				
$28 \leq x < 32$	8	30				
$32 \leq x < 36$	3	34				
			$\bar{x} =$			

(b) Hence calculate the standard deviation using the table.

(2)


(c) Now use your calculator to verify your answer.

(2)

[9]

**QUESTION 3**

After protracted union protests, a company analysed its salary structure for employees. They found that the salaries were symmetrically distributed with a mean of R8 850 per month and a standard deviation of R2 950 per month. Research indicated that if the monthly salary was below R3 000, the employee would not maintain an acceptable quality of life.



- (a) Estimate the percentage of employees who will struggle to maintain an acceptable quality of life. (2)
  - (b) Estimate the percentage of employees who earn more than R11 800 per month. (1)
  - (c) Do you think that the company has a fair salary structure? Use the given data to motivate your answer. (1)
- [4]




**QUESTION 4**

A motor company did research on how the speed of a car affects the fuel consumption of the vehicle. The following data was obtained:

Speed in km/h	60	75	115	85	110	95	120	100	70
Fuel consumption in $\ell/100$ km	11,5	10	8,4	9,2	7,8	8,9	8,8	8,6	10,2

- (a) Represent the data as a scatter plot on the diagram provided. (3)



- (b) Suggest whether a linear, quadratic or exponential function would best fit the data. (1)
- (c) What advice can the company give about the driving speed in order to keep the cost of fuel to a minimum? (2)
- [6]



**SECTION D: HOMEWORK SOLUTIONS**

**QUESTION 1**


(a)

Class intervals (ages)	Frequency ✓	Cumulative frequency ✓
$16 \leq x < 20$	0	0
$20 \leq x < 24$	5	5
$24 \leq x < 28$	10	15
$28 \leq x < 32$	5	20
$32 \leq x < 36$	3	23

(2)

(b)

Class intervals (ages)	Frequency	Cumulative frequency	Graph points
$16 \leq x < 20$	0	0	(20; 0)
$20 \leq x < 24$	5	5	(24; 5)
$24 \leq x < 28$	10	15	(28; 15)
$28 \leq x < 32$	5	20	(32; 20)
$32 \leq x < 36$	3	23	(36; 23)



**Teacher Note:**  
 Explain to learners that the above table is not required as marks are not given for the column with the coordinates. However, it is a good idea to determine the coordinates carefully before plotting. A mark is given to each point plotted correctly, and one mark given for correct shape of the ogive.

(6)

(c)

<p><b>Lower quartile</b></p> $23 \times \frac{1}{4} = 5,75$ <p>Therefore <math>Q_1 = 24</math></p> <p><b>Median</b></p> $23 \times \frac{1}{2} = 11,5$ <p>Therefore Median = 26</p> <p><b>Upper quartile</b></p> $23 \times \frac{3}{4} = 17,25$ <p>Therefore <math>Q_3 = 30</math></p>	<p style="text-align: center;">✓</p> <p style="text-align: center;">✓</p> <p style="text-align: right;">(3)</p>
---	---

[11]

**QUESTION 2**

(a)



Class intervals	Frequency (f)	Midpoint (m)	$f \times m$ ✓	$m - \bar{x}$ ✓	$(m - \bar{x})^2$ ✓	$f \times (m - \bar{x})^2$ ✓
$20 \leq x < 24$	5	22	110	-5	25	125
$24 \leq x < 28$	10	26	260	-1	1	10
$28 \leq x < 32$	5	30	150	3	9	45
$32 \leq x < 36$	3	34	102	7	49	147
			$\bar{x} = \frac{622}{23} = 27$ ✓			$\sum f \times (m - \bar{x})^2 = 327$

(5)


<p>(b)</p> $SD = \sqrt{\frac{\sum f \cdot (x - \bar{x})^2}{23}} = \sqrt{\frac{327}{23}} = 3,8$	<p style="text-align: center;">✓✓</p> <p style="text-align: right;">(2)</p>
--	---

<p>(c) <b>CASIO fx-82ES PLUS:</b>                  MODE                  2 : STAT                  1 : 1 – VAR                  SHIFT SETUP                  3: STAT                  1: ON                  Enter the midpoints:                  22= 26= 30= 34=                  Enter the frequencies:                  5= 10= 5= 3=                  AC SHIFT 1                  4: VAR                  3 : <math>x\sigma n</math> =                  The answer will read: 3,8</p> <p><b>SHARP DAL:</b>                  MODE 1=                  Enter data:                  22 STO 3 M+                  26 STO 9 M+                  30 STO 8 M+                  34 STO 3 M+                  RCL 6 to get 3,8</p>	<p>✓✓</p> <p>(2)</p>
--	----------------------



[9]

QUESTION 3



One standard deviation interval:

$$(\bar{x} - s ; \bar{x} + s)$$

$$= (8850 - 2950 ; 8850 + 2950)$$

$$= (5900 ; 11800)$$

Two standard deviation intervals:

$$(\bar{x} - 2s ; \bar{x} + 2s)$$

$$= (8850 - 2 \times 2950 ; 8850 + 2 \times 2950)$$

$$= (2950 ; 14750)$$

Three standard deviation intervals:

$$(\bar{x} - 3s ; \bar{x} + 3s)$$


$$= (8850 - 3 \times 2950 ; 8850 + 3 \times 2950)$$

$$= (0 ; 17700)$$

2%	✓✓	(2)
16%	✓	(1)
No, since there are some employees (less than 2%) earn below R3000,00. These employees will not live an acceptable lifestyle economically. <b>OR</b> Yes, there is a fair distribution of salaries since the majority of the employees, i.e. 68% earn a salary between R5 900 and R11 800 per month. Some employees will have more responsibilities or work longer hours and thus must be compensated accordingly. Less than 2% earn below R3000,00.	✓	(1)
		[4]

QUESTION 4

a.



✓  
✓  
✓

(3)

(b) Quadratic	✓ (1)
(c) Based on the quadratic trend the best fuel consumption occurs when the car is driven at 110 km/h. To keep its fuel bill to a minimum, drivers should drive at 110km/h	✓✓ (2)

[6]



## SESSION 16.2

## TOPIC: TRANSFORMATIONS



**Teacher Note:** Learners must revise all previous grades work on Transformations. The focus in grade 12 will be on rotations through any angle  $\theta$  however you may be examined on all the work learned thus far on Transformations.

## SECTION A: TYPICAL EXAM QUESTIONS

## QUESTION 1

Determine the coordinates of the image of point  $A(-2; -\sqrt{3})$  after a clockwise rotation, about the origin through an angle of  $210^\circ$ . [6]

**Hint:** Be careful with your signs, as this is a clockwise rotation.


## QUESTION 2

Consider a square with coordinates  $P(0; -5)$ ,  $Q(5;0)$ ,  $R(0;5)$  and  $S(-5;0)$  and answer the following questions:

- 2.1 If  $P'Q'R'S'$  is the reflection of  $PQRS$  about the line  $y = -x$  give the coordinates of  $P'$  and  $R'$ . (4)
  - 2.2 Calculate, correct to two decimal places, the coordinates of  $P''$  and  $R''$  if  $P'Q'R'S'$  is rotated about the origin through an angle of  $45^\circ$  anticlockwise to give  $P''Q''R''S''$ . (4)
  - 2.3 If  $S$  is rotated anticlockwise about the origin through  $60^\circ$  to a new position  $S'''$ , find the coordinates of  $S'''$  without a calculator – leave your answer on surd form. (4)
- [24]

## QUESTION 3

- 3.1 On the diagram provided below, draw figure  $ABCD$  with the coordinates of the vertices as follows:  $A(-6;6)$ ,  $B(-4;2)$ ,  $C(-2;6)$  and  $D(-4;8)$ . (1)
- 3.2 On the same diagram, draw the image  $A'B'C'D'$  if  $ABCD$  is rotated  $90^\circ$  anticlockwise. Indicate the coordinates of  $B'$  and  $C'$ . (2)



- 3.3 State the general rule in terms of  $x$  and  $y$  of the rotation  $90^\circ$  anti-clockwise, using the notation  $(x; y) \rightarrow$  (1)
- 3.4 Now draw the image  $A'B'C'D'$  if  $ABCD$  is transformed under the rule:  
 $(x; y) \rightarrow (\frac{1}{2}x; \frac{1}{2}y)$ . (2)
- 3.5 Write down the value of the following ratio:  $\frac{\text{Area } ABCD}{\text{Area } A'B'C'D'}$  (2)
- 3.6 Draw image  $A''B''C''D''$  if  $A'B'C'D'$  is rotated  $180^\circ$  clockwise. Indicate the coordinates of  $A''$ . (2)
- 3.7 Draw the image  $EFGH$  if  $A'B'C'D'$  is transformed under the rule  $(x, y) \rightarrow (-x; y)$ . Indicate the coordinates of  $E$ . (2)
- 3.8 Write down the single algebraic rule if  $ABCD$  is reflected about the  $x$ -axis, followed by a translation of 7 units right, followed by an enlargement by a scale factor of 2 units. (4)
- [16]

**QUESTION 4**

A circle with equation  $x^2 + y^2 - 2x - 4y = 4$  is rotated  $90^\circ$  anticlockwise about the origin and then enlarged by a scale factor of 2, find the new equation. [6]

**Hint:** Rule for rotation through  $90^\circ$  anticlockwise about the origin  $(x, y) \rightarrow (-y; x)$ .


**QUESTION 5**

(DOE Nov. 2010 P2)

A transformation T is described as follows:

- A reflection in the x-axis, followed by
- A translation of 4 units left and 2 units down, followed by
- An enlargement through the origin by a factor of 2

In the diagram  $\Delta ABC$  is given with vertices  $A(2 ; -2)$ ,  $B(4 ; -3)$  and  $C(1 ; -4)$ .



- 5.1 If  $\Delta ABC$  is transformed by T to  $A'B'C'$  (in that order), on the same system of axes sketch  $\Delta A'B'C'$ . Show ALL the steps. (6)
- 5.2 Write down the general rule for  $(x ; y)$  under transformation T in the form  $(x ; y) \rightarrow$  (4)
- 5.3 Calculate the area of  $\Delta A'B'C'$ . (4)  
[14]

**QUESTION 6**

(DOE March 2011)

Consider the point A  $(-12 ; 6)$ . The point is reflected about the x-axis to  $A'$

- 6.1 Write down the coordinates of  $A'$ . (1)
- 6.2 An alternative transformation from A to  $A'$  is a rotation about the origin through  $\alpha^\circ$ , where  $\alpha \in [0; 90]$  Calculate  $\alpha$ . (6)  
[7]



**SECTION B: SOLUTIONS TO SECTION A**

**QUESTION 1**

$$\begin{aligned}
 x' &= x_A \cos \theta - y_A \sin \theta = -2 \cos(-210) - (-\sqrt{3}) \sin(-210) \\
 &= -2 \cos(-210+360) + 3 \sin(-210+360) \quad (\text{Add } 360 \text{ to make angles positive}) \\
 &= -2 \cos 150 + \sqrt{3} \sin 150 \\
 &= -2 \cos(180-30) + 3 \sin(180-30) \\
 &= 2 \cos 30 + \sqrt{3} \sin 30 \\
 &= 2\left(\frac{\sqrt{3}}{2}\right) + 3\left(\frac{1}{2}\right) = \frac{3\sqrt{3}}{2}
 \end{aligned}$$

$$\begin{aligned}
 y' &= y_A \cos \theta + x_A \sin \theta = -\sqrt{3} \cos(-210) + (-2) \sin(-210) \\
 &= -\sqrt{3} \cos 150 - 2 \sin 150 \\
 &= 3 \cos 30 - 2 \sin 30 = \sqrt{3} \left(\frac{\sqrt{3}}{2}\right) - 2 \left(\frac{1}{2}\right) = \frac{1}{2}
 \end{aligned}$$

The image is  $A'\left(\frac{3\sqrt{3}}{2}; \frac{1}{2}\right)$  **[6]**

**QUESTION 2**

2.1 Reflection in  $y=-x$ :  $(x; y) \rightarrow (-y; x)$   
 $P'(5; 0)$  and  $R'(-5; 0)$  (4)


2.2  $P''(5 \cos 45; 5 \sin 45) \therefore P''(3.54; 3.54)$  (4)

2.3  $S'''(-5 \cos 60; -5 \sin 60)$   
 $S'''(-5\left(\frac{1}{2}\right); -5\left(\frac{\sqrt{3}}{2}\right)) \therefore S'''(\frac{-5}{2}; \frac{-5\sqrt{3}}{2})$  (4)

**[12]**

**QUESTION 3**

3.1 & 3.2 See diagram



(1)

(2)

3.3  $(x; y) \rightarrow (-y; x)$



(1)

3.4 See diagram

(2)

3.5  $\frac{\text{Area ABCD}}{\text{Area A''B''C''D''}} = \frac{1}{k^2} = \frac{1}{(\frac{1}{2})^2} = 4$

(2)

3.6 See diagram

(2)

3.7 See diagram

(2)

3.8

$(x; y) \rightarrow (x; -y)$

$(x; -y) \rightarrow (x+7; -y)$

$(x+7; -y) \rightarrow (2(x+7); -2y)$

$\therefore (x; y) \rightarrow (2x+14; -2y)$

(4)

[16]

**QUESTION 4**

$x^2 + y^2 - 2x - 4y = 4$

$(x - 1)^2 + (y - 2)^2 = 9$

$\therefore$  the centre is (1;2) and the radius = 3

Rule for rotation through 90° anti-clockwise about the origin;  $(x; y) \rightarrow (-y; x)$

$\therefore$  The centre of the image is (-2;1) The image is then enlarged by a factor of 2


$\therefore$  the centre is (-4;2) and the radius = 6.

New Eqn:  $(x + 4)^2 + (y - 2)^2 = 36$

[6]

QUESTION 5

5.1



**Note:**

- If the candidate only draws the correct triangle with labels, **full marks**
- If they plot the points correctly and do not draw the triangle, **max 5 / 6 marks**
- In the 3 sketches, if one vertex of the three is wrong, then 1 / 2 marks for the incorrect sketch, then CA applies.
- If they write down the points and do not plot the points and draw the triangle **max 3 / 6 marks**
- If the vertices are correct but not labelled and the points are joined **max 5 / 6 marks**
- If the vertices are correct, not labelled and not joined **max 4 / 6 marks**
- If a candidate finds a formula first and gets it wrong

Max 1 mark for the formula

Max 2 marks for the calculation of A'B'C' coordinates (CA)

1 mark for plotting 3 vertices

1 mark for completing the triangle and labelling (6)

5.2  $(x; y) \rightarrow (x; -y) \rightarrow (x-4; -y-2) \rightarrow (2x-8; -2y-4)$  (If the candidate gives the answer only-award full marks) (4)

5.3 (Please note: there are several different ways to do this question)

$$m_{AC}=2 \text{ and } m_{AB}=-0.5$$

$$\text{Therefore } \hat{C}AB=90 \text{ (} m_{AC} \times m_{AB}=-1 \text{); } AB=\sqrt{5} \text{ and } AC=\sqrt{5}$$

$$\text{Area of } \triangle ABC = 0.5 (\sqrt{5})(\sqrt{5}) = 5/2$$

$$\therefore \text{Area of } \triangle A'B'C' = 4 \times 5/2 = 10 \text{ square units}$$

(4)

**[14]****QUESTION 6**

6.1  $A'(-12; -6)$

(1)

6.2  $x' = x_A \cos \alpha - y_A \sin \alpha$   
 $-12 \cos \alpha - 6 \sin \alpha = -12$   
 $-2 \cos \alpha - \sin \alpha = -2 \quad (1)$

$$y' = y_A \cos \alpha + x_A \sin \alpha = 6 \cos \alpha - 12 \sin \alpha = -6$$

$$\cos \alpha = 2 \sin \alpha - 1 \quad (2)$$

substitute (2) into (1)

$$-2(2 \sin \alpha - 1) - \sin \alpha = -2$$

$$-4 \sin \alpha + 2 - \sin \alpha = -2$$

$$-5 \sin \alpha = -4$$

$$\sin \alpha = 4/5 \dots\dots \alpha = 53,13^\circ$$

(6)

**[7]**
**SECTION C: HOMEWORK**
**QUESTION 1**

Point P(2; 4) is rotated about the origin through an obtuse angle  $\theta$ , in an anti-clockwise

direction. The image is X(-3 2; y),  $y < 0$ .


Find:

a) The value of y

b) The angle  $\theta$

**QUESTION 2**


In the diagram below, the circle with centre the origin is rotated anti-clockwise about the origin through an angle of  $\theta$  degrees. Point  $A(4;3)$  lies on the circle and the image of point A is point B (coordinates are indicated on the diagram).



- 2.1 Determine the length of the radius of the circle
- 2.2 Calculate the size of angle  $\theta$
- 2.3 Hence show that  $AB = 5\sqrt{2 - \sqrt{3}}$
- 2.4 Calculate the area of  $\triangle OAB$


**QUESTION 3**

In the diagram below  $\triangle PQR$  has been enlarged through the origin by a scale factor of 3 to give its image  $\triangle XYZ$ .



- 3.1 Write down the coordinates of  $\triangle XYZ$ .
- 3.2 Calculate  $\hat{Y}$  (round off your answer to one decimal place)

**SECTION D: HOMEWORK SOLUTIONS**



**QUESTION 1**

a)  $OP=OX$

$$(2 - 0)^2 + (4 - 0)^2 = (-3\sqrt{2} - 0)^2 + (y)^2$$

$$20=18 + y^2$$

$$\therefore y^2 = 2$$

$$\therefore y = \pm\sqrt{2} \quad \text{but } y < 0 \quad \therefore y = -\sqrt{2}$$

$$X(-3\sqrt{2}; -\sqrt{2})$$

b)  $x' = x_A \cos\theta - y_A \sin\theta$  and  $y' = y_A \cos\theta + x_A \sin\theta$   
 $-3\sqrt{2} = 2\cos\theta - 4\sin\theta$  .....(1)  $- \sqrt{2} = 4\cos\theta + 2\sin\theta$  .....(2)

Multiply equation (1) by -2 and then add the equations

$$6\sqrt{2} = -4\cos\theta + 8\sin\theta$$

$$\underline{- \sqrt{2} = 4\cos\theta + 2\sin\theta}$$

$$5\sqrt{2} = 10\sin\theta$$

$$\sin\theta = \frac{\sqrt{2}}{2}$$

$$\therefore \theta = 45^\circ \text{ but since } \theta \text{ is obtuse } \theta = 135^\circ$$

**QUESTION 2**

2.1

$$(4)^2 + (3)^2 = r^2$$

$$\therefore r^2 = 25$$

$$\therefore r = 5$$

2.2

$$4 \cos \theta - 3 \sin \theta = \frac{4\sqrt{3} - 3}{2} \dots\dots A$$

$$3 \cos \theta + 4 \sin \theta = \frac{3\sqrt{3} + 4}{2} \dots\dots B$$

$$16 \cos \theta - 12 \sin \theta = 2(4\sqrt{3} - 3) \dots\dots A \times 4$$

$$9 \cos \theta + 12 \sin \theta = \frac{3(3\sqrt{3} + 4)}{2} \dots\dots B \times 3$$

$$\therefore 25 \cos \theta = 2(4\sqrt{3} - 3) + \frac{3(3\sqrt{3} + 4)}{2}$$

$$\therefore 25 \cos \theta = \frac{25\sqrt{3}}{2}$$

$$\therefore \cos \theta = \frac{\sqrt{3}}{2}$$

$$\therefore \theta = 30^\circ$$

2.3

$$AB^2 = (5)^2 + (5)^2 - 2(5)(5) \cos 30^\circ$$

$$\therefore AB^2 = 50 - 50 \left( \frac{\sqrt{3}}{2} \right)$$

$$\therefore AB^2 = 50 - 25\sqrt{3}$$

$$\therefore AB^2 = 25(2 - \sqrt{3})$$

$$\therefore AB = 5\sqrt{2 - \sqrt{3}}$$



2.4

$$\text{Area } \Delta OAB = \frac{1}{2} (5)(5) \sin 30^\circ$$

$$\therefore \text{Area } \Delta OAB = \frac{25}{4} \text{ units}^2$$

**QUESTION 3**

3.1 X(-6; 0) Y(3, 6) and Z(6; -6)

3.2 Here you will use Analytical geometry to help work out the angles of inclination

$$M_{xy} = \frac{2}{3} \quad \text{and}$$

$$M_{yz} = -4$$

$$\tan \theta = \frac{2}{3}$$

$$\tan \beta = -4$$

$$\theta = 33.69 \dots$$

$$\beta = 104.03 \dots$$

$$\therefore \alpha = 75.96 \dots$$

$$\therefore = 180 - (75.96 + 33.69) = 70.4^\circ$$



SESSION 17.1

TOPIC: FUNCTIONS



**Teacher Note:** Learners must be able to sketch all graphs from grades 10 & 11, including their inverses. It is important to understand how to find domain & range, intercepts, turning points, asymptotes and be able to interpret and identify all graphs. Vertical shifts affect the y values and horizontal shifts affect the x values.


SECTION A: TYPICAL EXAM QUESTIONS

QUESTION 1

On the following page, the diagram, which is drawn to scale, shows the graphs of  $f(x) = x^2 - 10x + 9$  and  $g(x) = 2x + c$ , where  $c \in \mathbb{R}$ .

The graph of  $f$  intersects the x-axis at P and Q and the y-axis at A. I and P are points of intersection of  $f$  and  $g$ . E is the turning point of  $f$ . F is a point on  $f$  such that AF is parallel to the x-axis and D is a point on  $g$  such that DE is parallel to the y-axis.





- 1.1 Show that the coordinates of P and Q are (1,0) and (9,0) respectively. (2)
- 1.2 Show that  $c=-2$  (2)
- 1.3 Determine the lengths of:
- 1.3.1 AF (2)
- 1.3.2 DE (4)
- 1.4 Determine the equation of  $p(x)$  if  $p$  is the mirror image of  $f$  in the line  $y=0$  (1)
- 1.5 Determine the value of  $x$  for which the tangent of  $f$  will be parallel to  $g$ . (4)
- [15]

**QUESTION 2**

If  $f(x) = \frac{2}{x}$  and  $g(x) = x^2 - 2$ , determine the value of the following:

2.1 p if  $g(-1) = p$  (2)


2.2 t if  $f(t) = -\frac{1}{2}$  (2)

2.3 k if  $g\left[f\left(\frac{1}{k}\right)\right] = -1$  (5)

[9]

**QUESTION 3**

In the diagram below, the graphs of  $f$  and  $g$  are shown. The graphs intersect at  $(1;3)$ . The asymptotes pass through the turning point of the parabola.



3.1 Determine the equation of  $f$  (3)

3.2 Write down the equations of the asymptotes (2)

3.3 Determine the equation of  $g$  (3)

3.4 Determine the coordinates of  $A$  (3)

3.5 Determine the values of  $x$  for which  $g(x) \leq 0$  (2)

3.6 Determine the equation of the graph formed if  $f$  is reflected about the  $y$ -axis (1)


[14]

**QUESTION 4**

- 4.1 Sketch the graph of  $f(x) = 2^x$  (2)
- 4.2 Determine the equation of the inverse  $f^{-1}$  (2)
- 4.3 Sketch the graph of the inverse on the same set of axes. (2)
- 4.4 If the graph of  $f$  is shifted 1 unit to the right to form the graph  $g$ , write down the equation of  $g$ . (1)
- 4.5 Sketch the graph of  $g$  on a separate set of axes (2)
- [9]

**QUESTION 5**

The graph of  $(x) = ax^2$ ,  $x \leq 0$  is sketched below. The point  $P(-6; -8)$  lies on the graph of  $f$ .



- 5.1 Calculate the value of  $a$  (2)
- 5.2 Determine the equation of  $f^{-1}$ , in the form  $y = \dots$  (3)
- 5.3 Write down the range of  $f^{-1}$  (1)
- 5.4 Draw the graph of  $f^{-1}$  on a set of axes. Indicate the coordinates of a point on the graph different from  $(0; 0)$  (2)
- 5.5 The graph of  $f$  is reflected across the line  $y = x$  and thereafter it is reflected across the  $x$ -axis. Determine the equation of the new function in the form  $y = \dots$  (3)
- [11]

**QUESTION 6**

Consider the function  $f(x) = \left(\frac{1}{3}\right)^x$

- 6.1 Is  $f$  an increasing or decreasing function? Give a reason for your answer. (2)
- 6.2 Determine  $f^{-1}(x)$  in the form  $y = \dots$  (2)
- 6.3 Write down the equation of the asymptote of  $f(x) - 5$ . (1)
- 6.4 Describe the transformation from  $f$  to  $g$  if  $g(x) = \log_3 x$  (2)
- [7]

**SECTION B: SOLUTIONS TO SECTION A****QUESTION 1**

- 1.1 x intercepts: P(1; 0) and Q(9; 0) (2)
- 1.2 graph g intersects f at P  $\therefore$  (1; 0) is a point on g  
 $g(x) = 2x + c$  Substitute point P.....  $2(1) + c = 0 \therefore c = -2$  (2)
- 1.3.1 can use symmetry and read off the graph itself OR:  
 Since AF  $\parallel$  x axis point F has co-ordinates (x; 9)  
 $\therefore x^2 - 10x + 9 = 9$   
 $x(x-10) = 0$   
 $\therefore x=10$  or  $x = 0 \therefore F(10; 9)$  so AF =  $10-0 = 10$  units (2)
- 1.3.2 Turning point of f:  $x = -\frac{b}{2a} = -\frac{(-10)}{2} = 5$   
 $f(5) = 25 - 50 + 9 = -16 \therefore E(5; -16)$   
 distance from the x axis to the T.P E is 16 units  
 D(5; y) lies on the line g(x):  $g(5) = 2(5) - 2 = 8$   
 Distance DE =  $8 + 16 = 24$  (4)
- 1.4  $p(x) = -f(x) = -x^2 + 10x - 9$  (1)
- 1.5 Parallel lines have equal gradients  $m$  g(x) = 2  
 $f(x) = 2x - 10 = 2$   
 $2x = 12$   
 $\therefore x = 6$  (4)
- [15]

**QUESTION 2**

- 2.1  $g(-1) = p$   
 $(-1)^2 - 2 = -1 = p$  (2)
- 2.2  $f(t) = -\frac{1}{2}$   
 $f(t) = \frac{2}{t} = -\frac{1}{2}$   
 $-t = 4 \therefore t = 4$  (2)

$$2.3 \quad g\left[f\left(\frac{1}{k}\right)\right] = -1$$

$$\left[f\left(\frac{1}{k}\right)\right] = \frac{2}{\frac{1}{k}} = 2k$$

$$g(2k) = (2k)^2 - 2 = 4k^2 - 2$$

$$4k^2 - 2 = -1$$

$$4k^2 = 1$$

$$k^2 = \frac{1}{4}$$

$$\therefore k = \pm \frac{1}{2}$$

(5)

[9]

**QUESTION 3**

3.1

$$y = a(x+1)^2 + 2$$

$$3 = a(-2+1)^2 + 2$$

$$\therefore 1 = a$$

$$\therefore f(x) = (x+1)^2 + 2$$

(3)

3.2

$$x = -1$$

$$y = 2$$

(2)

3.3

$$y = \frac{a}{x+1} + 2$$

$$\therefore 3 = \frac{a}{-2+1} + 2$$

$$\therefore 3 = -a + 2$$

(3)

$$\therefore a = -1$$

$$\therefore g(x) = \frac{-1}{x+1} + 2$$

3.4

$$0 = \frac{-1}{x+1} + 2$$

$$\therefore 0 = -1 + 2x + 2$$

$$\therefore -2x = 1$$

$$\therefore x = -\frac{1}{2}$$

$$\therefore A\left(-\frac{1}{2}; 0\right)$$

(3)

3.5

$$g(x) \leq 0$$

$$\therefore -1 < x \leq -\frac{1}{2}$$

(2)




3.6

$$y = (x-1)^2 + 2$$

(1)  
[14]

QUESTION 4

4.1



(2)

4.2

$$y = 2^x$$

$$\therefore x = 2^y$$

$$\therefore y = \log_x 2$$

$$f^{-1}(x) = \log_x 2$$



(2)


4.3 See diagram

(2)

4.4  $g(x) = 2^{x-1}$

(1)

4.5



(2)  
[9]


**QUESTION 5**

5.1  $f(x) = ax^2$   
 $a(-6)^2 = -8$   
 $36a = -8$   
 $\therefore a = -\frac{2}{9}$  (2)

5.2  $y = -\frac{2}{9}x^2$   
 $x = -\frac{2}{9}y^2$   
 $y^2 = -\frac{9}{2}x$   
 $y = \pm\sqrt{-\frac{9x}{2}}$ , since  $y \leq 0$   $y = -\sqrt{-\frac{9x}{2}}$  or  $y = -3\sqrt{-\frac{x}{2}}$  (3)

5.3  $y \leq 0$  (1)

5.4



(2)

5.5  $y = -f^{-1}(x) = -\left(-\sqrt{-\frac{9x}{2}}\right) = \sqrt{-\frac{9x}{2}}$  or  $3\sqrt{-\frac{x}{2}}$  (3)

[11]

**QUESTION 6**

6.1 Decreasing function  
 Since  $0 < a < 1$  OR As  $x$  increases,  $f(x)$  decreases (2)



MATHEMATICS

GRADE 12

SESSION 17


(TEACHER NOTES)

6.2

$$f^{-1}: x = \left(\frac{1}{3}\right)^y$$

$$y = \log_{\frac{1}{3}} x$$

OR



OR

$$f^{-1}: x = \left(\frac{1}{3}\right)^y$$

$$y = -\log_3 x$$

(2)

6.3  $y = -5$ 

(1)

6.4 Reflection about  $y = x$ .Reflection about the  $x$ -axis.**OR**Reflection about the  $y$ -axis.Then reflection about the line  $y = x$ .**OR**Reflection about the line  $y = -x$  followed by reflection about the  $y$ -axis.**OR**Rotation through  $90^\circ$  in a clockwise direction.**OR**Rotation through  $90^\circ$  in an anti-clockwise direction.

Reflection through the origin.

(2)

[7]

<b>SECTION C: HOMEWORK</b>
----------------------------

**QUESTION 1**Given  $f(x) = 2x$  find a simplified expression for:  $f(x) + f\left(\frac{1}{x}\right) + \frac{1}{f(x)} + f^{-1}(x)$ 

[6]

**QUESTION 2**Given that a function  $f$  satisfies the following conditions : $f(0) = 2$  ,  $f(-2) = 0$  ,  $f'(-1) = 0$  and  $f'(x) > 0$  for  $x \neq 0$ Draw a rough sketch of the graph  $f$ .

[5]

**QUESTION 3**Consider the function  $f(x) = (x + 1)^2 - 4$ 

3.1 Draw a neat sketch graph indicating the coordinates of the intercepts with the axes, the coordinates of the turning point and the equation of the axis of symmetry.

(6)

3.1.1 Give the range.

(2)

[8]

MATHEMATICS

GRADE 12

SESSION 17

(TEACHER NOTES)


**QUESTION 4**

Given:  $f(x) = a^x$  passing through the point  $(2; \frac{1}{4})$ ;  $g(x) = 4x^2$

- 4.1 Prove that  $a = \frac{1}{2}$  (2)
- 4.2 Determine the equation of  $y = f^{-1}(x)$  in the form  $y = \dots$  (2)
- 4.3 Determine the equation of  $y = h(x)$  where  $h(x)$  is the reflection of  $f(x)$  about the x-axis (1)
- 4.4 Determine the equation of the inverse of  $g$  in the form  $y =$  (2)
- 4.5 How must the domain of  $g(x)$  be restricted so that  $g^{-1}(x)$  will be a function? (2)
- [9]

**QUESTION 5**

The graph of  $g$  is shown where  $g(x) = \log_a x$  where  $0 < x < 1$   
 $(\frac{1}{2}; -1)$  is a point on  $g$



- 5.1 Determine the value of  $a$  (2)
- 5.2 Write down the domain of  $g(x)$  (2)
- 5.3 Write the equation of  $g^{-1}$  in the form  $g^{-1}(x) = \dots$  and state the domain (2)
- [6]


**SECTION D: HOMEWORK SOLUTIONS**

**QUESTION 1**

$$\begin{aligned}
 f(x) &= 2x \\
 f\left(\frac{1}{x}\right) &= 2\left(\frac{1}{x}\right) \\
 \frac{1}{f(x)} &= \frac{1}{2x} \\
 f^{-1}(x) &= \frac{1}{2}x \quad y=2x \text{ swop } x \text{ and } y \text{ to find inverse: } x = 2y \text{ so } y = \frac{1}{2}x \\
 f(x) + f\left(\frac{1}{x}\right) + \frac{1}{f(x)} + f^{-1}(x) &= 2x + \frac{2}{x} + \frac{1}{2x} + \frac{1}{2}x \\
 &= \frac{5x^2+5}{2x}
 \end{aligned}$$

[6]


**QUESTION 2**



[5]

**QUESTION 3**

3.1



(6)

3.1.1 Range:  $y \in [-4; \infty)$

(2)

**[8]****QUESTION 4**

4.1

$$y = a^x$$

$$\therefore \frac{1}{4} = a^2$$

$$\therefore a = \frac{1}{2}$$

(2)

4.2

$$y = \left(\frac{1}{2}\right)^x$$

$$\therefore x = \left(\frac{1}{2}\right)^y$$

$$\therefore y = \log_{\frac{1}{2}} x$$

(2)

4.3

$$y = \left(\frac{1}{2}\right)^x$$

(1)

4.4

$$y = 4x^2$$

$$\therefore x = 4y^2$$

$$\therefore \frac{x}{4} = y^2$$

$$\therefore y = \pm \sqrt{\frac{x}{4}}$$

(2)

4.5  $x > 0$  or  $x < 0$

(2)

**[9]****QUESTION 5**

5.1  $g(-\frac{1}{2}) = -1$

$$\log_a \frac{1}{2} = -1$$

$$\therefore a^{-1} = \frac{1}{2}$$

$$\therefore a = 2$$

(2)

5.2  $x > 0$  and  $x \neq 1$  (NB: The graph of  $g$  is only drawn for  $0 < x < 1$  but this is not the domain)

(2)

5.3  $g^{-1}(x) = 2^x$   
 $x \in \mathbb{R}, x \neq 0$

(NB: From the log graph  $x \neq 1$  so its' inverse will have  $y \neq 1$  the value that will make  $y=1$  in  $g^{-1}(x)$  is  $x=0$  so it must be excluded from the domain.)

(2)

**[6]**

**SESSION 17.2**

**TOPIC: CALCULUS**

**LESSON OVERVIEW**


- |    |                          |            |
|----|--------------------------|------------|
| 1. | Introduction session:    | 10 minutes |
| 2. | Typical exam questions:  |            |
|    | Question 1:              | 10 minutes |
|    | Question 2:              | 15 minutes |
|    | Question 3:              | 10 minutes |
|    | Question 4:              | 15 minutes |
|    | Question 5:              | 15 minutes |
| 3. | Discussion of solutions: | 15 minutes |

**SECTION A: TYPICAL EXAM QUESTIONS**

**QUESTION 1: 10 minutes**

A drinking glass, in the shape of a cylinder, must hold 200 ml of liquid when full.

- 1.1 Show that the height of the glass,  $h$ , can be expressed as  $h = \frac{200}{\pi r^2}$ . (2)
- 1.2 Show that the surface area of the glass can be expressed as  $S(r) = \pi r^2 + \frac{400}{r}$ . (2)
- 1.3 Hence determine the value of  $r$  for which the total surface area of the glass is a minimum. (5)



**QUESTION 2: 15 minutes**

- 2.1. Differentiate  $f(x) = -3x^2$  by first principles. (5)
- 2.2. If  $f(x) = (3x^2 - x)^2$ , find  $f'(x)$  (4)
- 2.3. If  $y = \frac{2x^2+3x-2}{x+2}$ , find  $\frac{dy}{dx}$  (3)
- 2.4. Find  $\frac{d}{dx} \left( \frac{4x^3-3x^2}{2\sqrt{x}} \right)$  (4)
- [16]

**QUESTION 3: 10 minutes**

3.1. Evaluate: (3)

$$\lim_{p \rightarrow 4} \frac{p^2 - 5p + 4}{4 - p}$$

3.2. Leaving your answer with positive exponents, find  $f'(x)$  if: (3)

$$f(x) = 4x^2 - 4x + \frac{1}{x^2} - \frac{1}{2}$$


3.3. Evaluate, leaving your answer in surd form, with positive exponents: (4)

$$D_x \left[ \frac{5x^2 - 3x}{\sqrt{x}} \right]$$

[10]

**QUESTION 4: 15 minutes**

Part of a rally track follows the path of a cubic curve. A plain view of this section of the track is shown in the diagram below. At a certain instant, cars A and B are at the turning points of the curve. Car C is at the point where  $x=150$ . The race starts at the origin, 0.



The function which describes this part of the track is:

$$f(x) = \frac{1}{900}x^3 - \frac{1}{5}x^2 + 9x$$

- 4.1. Determine the coordinates of cars A and B (6)
- 4.2. Find the average gradient between car A and car C. (2)
- 4.3. Find the equation of the tangent to the track at the starting flag. (2)
- 4.4. At which point between A and B does a car stop turning to the right and start turning to the left? (3)  
[13]

**QUESTION 5: 15 minutes**

The cross section of a hilly region can be drawn as the graph of  $y = x^3 - 8x^2 + 16x$ , for  $0 \leq x \leq 6$ , where  $x$  is measured in kilometres and  $y$  is the height above sea level in meters.

- 5.1. Draw the cross section for  $0 \leq x \leq 6$ . Show all calculations. (7)
- 5.2. Mark these the peak and the valley on your cross section and calculate the difference in height between the two. (2)
- 5.3. Determine the gradient of the hill at the point P where  $x = 1$ . (2)
- 5.4. Hence, determine the equation of the tangent to the hill at P. (4)  
[15]

**SECTION B: SOLUTIONS AND HINTS TO SECTION A**


**QUESTION 1**

1.1	$V = \pi r^2 h$ $200 = \pi r^2 h$ $h = \frac{200}{\pi r^2}$	$\checkmark V = \pi r^2 h$ $\checkmark 200 = \pi r^2 h$	(2)
1.2	$\text{Surface Area} = 2\pi rh + \pi r^2$ $S(r) = \pi r^2 + \frac{200}{\pi r^2} \cdot 2\pi r$ $S(r) = \pi r^2 + \frac{400}{r}$	$\checkmark S = 2\pi rh + \pi r^2$ $\checkmark S(r) = \pi r^2 + \frac{200}{\pi r^2} \cdot 2\pi r$	(2)

1.3	$S(r) = \pi r^2 + 400r^{-1}$ $\frac{dS}{dr} = 2\pi r - 400r^{-2}$ <p>At minimum: <math>\frac{dS}{dr} = 0</math></p> $2\pi r - \frac{400}{r^2} = 0$ $\pi r^3 - 200 = 0$ $r^3 = \frac{200}{\pi}$ $r = 3,99 \text{ cm}$	$\checkmark S(r) = \pi r^2 + 400r^{-1}$ $\checkmark \frac{dS}{dr} = 2\pi r - 400r^{-2}$ $\checkmark 2\pi r - \frac{400}{r^2} = 0$ $\checkmark r^3 = \frac{200}{\pi}$ $\checkmark r = 3,99 \text{ cm}$	(5)
-----	--	---	-----

[9]

**QUESTION 2**

2.1.	$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{-3x^2 - 6xh - 3h^2 + 3x^2}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{h(-6x - 3h)}{h}$ $f'(x) = \lim_{h \rightarrow 0} -6x - 3h$ $= -6x$	$f(x) = -3x^2$ $f(x+h) = -3(x+h)^2$ $= -3x^2 - 6xh - 3h^2$	$\checkmark -3x^2 - 6xh - 3h^2$ $\checkmark +3x^2$ $\checkmark h(-6x - 3h)$ $\checkmark -6x - 3h$ $\checkmark -6x$	(5)
2.2.	$f(x) = (3x^2 - x)^2$ $= 9x^4 - 6x^3 + x^2$ $f'(x) = 36x^3 - 18x^2 + 2x$		$\checkmark 9x^4 - 6x^3 + x^2$ $\checkmark 36x^3$ $\checkmark -18x^2$ $\checkmark +2x$	(4)
2.3.	$y = \frac{2x^2 + 3x - 2}{x + 2}$ $= \frac{(2x - 1)(x + 2)}{x + 2}$ $= 2x - 1$ $\frac{dy}{dx} = 2$		$\checkmark (2x - 1)(x + 2)$ $\checkmark 2x - 1$ $\checkmark \frac{dy}{dx} = 2$	(3)
2.4.	$\frac{d}{dx} \left( \frac{4x^3 - 3x^2}{2\sqrt{x}} \right)$ $= \frac{d}{dx} \left( \frac{4x^{\frac{3}{2}} - 3x^{\frac{5}{2}}}{2} \right)$ $= \frac{d}{dx} \left( 2x^{\frac{3}{2}} - \frac{3}{2}x^{\frac{5}{2}} \right)$ $= 5x^{\frac{3}{2}} - \frac{9}{4}x^{\frac{1}{2}}$		$\checkmark 2x^{\frac{5}{2}}$ $\checkmark -\frac{3}{2}x^{\frac{3}{2}}$ $\checkmark 5x^{\frac{3}{2}}$ $\checkmark -\frac{9}{4}x^{\frac{1}{2}}$	(4)

[16]



**QUESTION 3**

3.1.	$\lim_{p \rightarrow 4} \frac{p^2 - 5p + 4}{4 - p}$ $= \lim_{p \rightarrow 4} \frac{(p - 4)(p - 1)}{-(p - 4)}$ $= \lim_{p \rightarrow 4} -(p - 1)$ $= -(4 - 1)$ $= -3$	$\checkmark (p - 4)(p - 1)$ $\checkmark -(p - 4)$ $\checkmark -3$ <p style="text-align: right;">(3)</p>
3.2.	$f(x) = 4x^2 - 4x + \frac{1}{x^2} - \frac{1}{2}$ $f(x) = 4x^2 - 4x + x^{-2} - \frac{1}{2}$ $f'(x) = 8x - 4 - \frac{2}{x^3}$	$\checkmark 8x$ $\checkmark -4$ $\checkmark -\frac{2}{x^3}$ <p style="text-align: right;">(3)</p>
3.3.	$D_x \left[ \frac{5x^2 - 3x}{x} \right]$ $= D_x \left[ 5x^{\frac{3}{2}} - 3x^{\frac{1}{2}} \right]$ $= \frac{15\sqrt{x}}{2} - \frac{3}{2\sqrt{x}}$	$\checkmark 5x^{\frac{3}{2}}$ $\checkmark -3x^{\frac{1}{2}}$ $\checkmark \frac{15\sqrt{x}}{2}$ $\checkmark -\frac{3}{2\sqrt{x}}$ <p style="text-align: right;">(4)</p>

**[10]**

**QUESTION 4**

4.1.	$f(x) = \frac{1}{900}x^3 - \frac{1}{5}x^2 + 9x$ $f'(x) = \frac{1}{300}x^2 - \frac{2}{5}x + 9$ $0 = \frac{1}{300}x^2 - \frac{2}{5}x + 9$ $0 = x^2 - 120 + 2700$ $0 = (x - 30)(x - 90)$ $x = 30 \text{ or } x = 90$ $\therefore A(30; 12) \quad B(90; 0)$	<div style="text-align: center;"></div> $\checkmark f'(x) = \frac{1}{300}x^2 - \frac{2}{5}x + 9$ $\checkmark 0 = \frac{1}{300}x^2 - \frac{2}{5}x + 9$ $\checkmark 0 = x^2 - 120 + 2700$ $\checkmark 0 = (x - 30)(x - 90)$ $\checkmark A(30; 12)$ $\checkmark B(90; 0)$ <p style="text-align: right;">(6)</p>
4.2.	$C(150; 600)$ $m_{AC} = \frac{600 - 120}{150 - 30}$ $m_{AC} = 4$	$\checkmark m_{AC} = \frac{600 - 120}{150 - 30}$ $\checkmark m_{AC} = 4$ <p style="text-align: right;">(2)</p>
4.3.	$f'(x) = \frac{1}{300}(0)^2 - \frac{2}{5}(0) + 9$ $m = 9$ $\therefore y = 9x$	$\checkmark m = 9$ $\checkmark \therefore y = 9x$ <p style="text-align: right;">(2)</p>
4.4.	$f''(x) = \frac{1}{150}x - \frac{2}{5}$ $0 = \frac{1}{150}x - \frac{2}{5}$ $0 = x - 60$ $x = 60$ $\therefore (60; 60)$	$\checkmark \frac{1}{150}x - \frac{2}{5}$ $\checkmark 0 = \frac{1}{150}x - \frac{2}{5}$ $\checkmark (60; 60)$ <p style="text-align: right;">(3)</p>

**[13]**

**QUESTION 5**

<p>5.1.</p>	$y = x^3 - 8x^2 + 16x$ $0 = x(x^2 - 8x + 16)$ $0 = x(x - 4)(x - 4)$ $x = 0 \quad x = 4$ $(0,0)(4;0)$ $3x^2 - 16x + 16 = 0$ $(3x - 4)(x - 4) = 0$ $x = \frac{4}{3} \quad x = 4$ $y = 9,48 \quad y = 0$ $\left(1\frac{1}{3}; 9,48\right)$ $6x - 16 = 0$ $6x = 16$ $x = 2\frac{2}{3}$ $y = 4,74$ $\left(2\frac{2}{3}; 4,74\right)$ 	<p>x-intercepts:                  ✓(0,0)                  ✓(4;0)</p> <p>Turning points:                  ✓(4;0)                  ✓(1 1/3; 9,48)</p> <p>Inflection:                  ✓(2 2/3; 4,74)</p> <p>✓✓plotted function</p> <p style="text-align: right;">(7)</p>
<p>5.2.</p>	<p>Difference in height = 9,48km</p>	<p>✓marking the valley and peak on function                  ✓9,48km</p> <p style="text-align: right;">(2)</p>

5.3.	$m = 3x^2 - 16 + 16$ $m = 3(1)^2 - 16(1) + 16$ $m = 3$	$\checkmark m = 3x^2 - 16 + 16$ $\checkmark m = 3$ <span style="float: right;">(2)</span>
5.4.	$y = (1)^3 - 8(1)^2 + 16 = 9$ $(1; 9) \quad m = 3$ $\therefore y - 9 = 3(x - 1)$ $y = 3x + 6$	$\checkmark (1; 9)$ $\checkmark y - 9 = 3(x - 1)$ $\checkmark 3x$ $\checkmark +6$ <span style="float: right;">(4) [15]</span>


**SECTIONC: HOMEWORK**

**QUESTION 1**

A builder wishes to construct a steel window frame in the shape of a rectangle with a semi-circular part on top. The radius of the semi-circular part is  $r$  metres and the width of the rectangular part is  $h$  metres.

- 1.1 Write down, in terms of  $h$  and  $r$ 
  - 1.1.1 the steel perimeter ( $P$ ) of the frame. (2)
  - 1.1.2 the area enclosed by the frame. (2)
- 1.2 The area enclosed by the frame is to be 4 square metres.  
 Show that the perimeter ( $P$ ) is  


$$P = \left(\frac{\pi}{2} + 2\right)r + \frac{4}{r}$$
 (4)
- 1.3 If the steel for the frame costs R10 per metre, calculate the value of  $r$  for which the total cost of the steel will be a minimum. (4)



[12]

**QUESTION 2**

Refer to the figure. The graph (not drawn to scale) of  $f(x) = 4x^3 + 27x^2 - 30x - 1$  is shown with A and B the turning points of the graph.



- 2.1. Determine the coordinates of A and B. (6)
- 2.2. Calculate the average gradient of  $f$  between the points A and B. (2)
- 2.3. C is the  $y$ -intercept of the graph. Determine the equation of the tangents to  $f$  at C. (3)
- 2.4. Determine the  $x$ -coordinate of the point on  $f$  where this tangent cuts the graph again. (3)

[14]

## SECTION D: SOLUTIONS TO HOMEWORK

## QUESTION 1

1.1.1	$P = 2h + 2r + \frac{1}{2} \times 2\pi r$ $\therefore P = 2h + 2r + \pi r$	$\checkmark 2h + 2r$ $\checkmark \pi r$	(2)
1.1.2	$A = 2rh + \frac{1}{2} \pi r^2$	$\checkmark 2rh$ $\checkmark \frac{1}{2} \pi r^2$	(2)
1.2	$4 = 2rh + \frac{1}{2} \pi r^2$ $\therefore 8 = 4rh + \pi r^2$ $\therefore 8 - \pi r^2 = 4rh$ $\therefore \frac{8 - \pi r^2}{4r} = h$ $P = 2h + 2r + \pi r$ $\therefore P = 2 \left( \frac{8 - \pi r^2}{4r} \right) + 2r + \pi r$ $\therefore P = \frac{8 - \pi r^2}{2r} + 2r + \pi r$ $\therefore P = \frac{4}{r} - \frac{\pi r}{2} + 2r + \pi r$ $\therefore P = \frac{4}{r} + \frac{\pi r}{2} + 2r$ $\therefore P = \frac{4}{r} + \left( \frac{\pi}{2} + 2 \right) r$ $\therefore P = \left( \frac{\pi}{2} + 2 \right) r + \frac{4}{r}$	$\checkmark 4 = 2rh + \frac{1}{2} \pi r^2$ $\checkmark \frac{8 - \pi r^2}{4r} = h$ $\checkmark P = 2 \left( \frac{8 - \pi r^2}{4r} \right) + 2r + \pi r$ $\checkmark P = \left( \frac{\pi}{2} + 2 \right) r + \frac{4}{r}$	(4)



1.3	$C = 10\left(\frac{\pi}{2} + 2\right)r + \frac{40}{r}$ $\therefore C = 5\pi r + 20r + 40r^{-1}$ $\therefore C'(r) = 5\pi + 20 - 40r^{-2}$ $\therefore C'(r) = 5\pi + 20 - \frac{40}{r^2}$ $\therefore 0 = 5\pi + 20 - \frac{40}{r^2}$ $\therefore \frac{40}{r^2} = 5\pi + 20$ $\therefore \frac{40}{5\pi + 20} = r^2$ $\therefore \sqrt{\frac{40}{5\pi + 20}} = r$ $\therefore r = 1,06\text{m}$	$\checkmark C = 10\left(\frac{\pi}{2} + 2\right)r + \frac{40}{r}$ $\checkmark C = 5\pi r + 20r + 40r^{-1}$ $\checkmark 0 = 5\pi + 20 - \frac{40}{r^2}$ $\checkmark r = 1,06\text{m}$
-----	--	--

**[12]**

**QUESTION 2**

2.1.	<p>At A and B: <math>f'(x) = 0</math></p> $f'(x) = 12x^2 + 54x - 30 = 0$ $2x^2 + 9x - 5 = 0$ $(2x - 1)(x + 5) = 0$ $x = \frac{1}{2} \text{ or } x = -5$ $f\left(\frac{1}{2}\right) = 4\left(\frac{1}{2}\right)^3 + 27\left(\frac{1}{2}\right)^2 - 30\left(\frac{1}{2}\right) - 1$ $= \frac{-35}{4} (-8,75)$ $f(-5) = 4(-5)^3 + 27(-5)^2 - 30(-5) - 1$ $= 324$ $\therefore A(-5; 324), \quad B\left(\frac{1}{2}; \frac{-35}{4}\right)$	$\checkmark f'(x) = 0$ $\checkmark \text{substitution of } x \text{ values}$ $\checkmark \checkmark A(-5; 324)$ $\checkmark \checkmark B\left(\frac{1}{2}; \frac{-35}{4}\right)$
2.2.	$\text{Ave Grad} = \frac{324 - \left(\frac{-35}{4}\right)}{-5 - \frac{1}{2}}$ $= \frac{-121}{2} (-60,5)$	$\checkmark \text{subs } x \text{ and } y \text{ values}$ $\checkmark = \frac{-121}{2} (-60,5)$
2.3.	$C(0; -1)$ $f'(0) = -30$ $\text{Equ. of tangent: } y = -30x - 1$	$\checkmark C(0; -1)$ $\checkmark f'(0) = -30$ $\checkmark y = -30x - 1(3)$
2.4.	$4x^3 + 27x^2 - 30x - 1 = -30x - 1$ $4x^3 + 27x^2 = 0$ $x^2(4x + 27) = 0$ $x = 0 \text{ or } x = -\frac{27}{4}$ $\therefore x = \frac{-27}{4}$	$\checkmark \text{cubic=tangent}$ $\checkmark x^2(4x + 27) = 0$ $\checkmark x = \frac{-27}{4}$

**[14]**

**SESSION 18.1****TOPIC: LINEAR PROGRAMMING**

**Teacher Note:** This session covers the basic rules and methods for solving Linear Programming questions. An illustration of the different kinds of questions that may be asked, is given and the theory is addressed. There are 2 solved questions and 1 additional homework question.

**LESSON OVERVIEW**

- |    |                                     |            |
|----|-------------------------------------|------------|
| 1. | Introduction to Linear Programming: | 5 minutes  |
| 2. | Typical exam questions:             |            |
|    | Question 1:                         | 30 minutes |
|    | Question 2:                         | 30 minutes |
| 3. | Homework Question:                  | 25 minutes |

**SECTION A: TYPICAL EXAM QUESTIONS****QUESTION 1**

Magical Homes, a home decor company, must produce at least 90 bedroom lamps per week. Not more than 18 people can be employed. An artisan, who earns R600 per week, can produce 9 lamps per week, while an apprentice, who earns R300 per week, only 5 lamps per week. At least one apprentice must be employed for every five artisans. The ratio of apprentices to artisans must not exceed 1:2. Assume that the number of artisans is  $x$  and the number of apprentices is  $y$ .

- |     |   |      |
|-----|---|------|
| 1.1 | Write down the inequalities.  | (4)  |
| 1.2 | Sketch the feasible region.   | (5)  |
| 1.3 | How many artisans and apprentices should be employed in order to minimise the wages paid? | (7)  |
|     |   | [16] |

**QUESTION 2**


TV Mania is a company manufacturing television sets. The company produces two types of television sets: TX and TY. At least one set of both must be manufactured per week. At most 6 sets of TX and 5 sets of TY can be manufactured per week. It takes five hours to manufacture a TX set and 4 hours to manufacture a TY set. A working week is 40 hours. Twice as many workers are required to manufacture TX sets than TY sets, while at least 8 workers are constantly occupied in the manufacturing process.

- |     |   |      |
|-----|---|------|
| 1.1 | Write down the constraints if the company produces $x$ TX sets and $y$ TY sets.                                       | (5)  |
| 1.2 | Represent the system of constraints on graph paper. Indicate the whole number points in the feasible region.          | (5)  |
| 1.3 | The company makes a profit of R500 on TX and R250 on TY.  |      |
|     | 1.3.1 Determine the number of each television set to be manufactured weekly if the company wants to maximise profits. | (2)  |
|     | 1.3.2 What number of each set will yield a minimum profit per week?   | (2)  |
|     |   | [14] |

**SECTION B: SOLUTIONS AND HINTS TO SECTION A**

**QUESTION 1**


<p>1.1</p> <p>Let the number of artisans be <math>x</math> and the number of apprentices be <math>y</math>.</p> <p><math>x + y \leq 18</math></p> <p><math>9x + 5y \geq 90</math></p> <p><math>y \leq \frac{1}{2}x</math></p> <p><math>y \geq \frac{1}{5}x</math></p>		<p>✓ <math>x + y \leq 18</math></p> <p>✓ <math>9x + 5y \geq 90</math></p> <p>✓ <math>y \leq \frac{1}{2}x</math></p> <p>✓ <math>y \geq \frac{1}{5}x</math></p> <p>(4)</p>
<p>1.2</p>	<p>See diagram below</p>	<p>✓ <math>x + y \leq 18</math></p> <p>✓ <math>9x + 5y \geq 90</math></p> <p>✓ <math>y \leq \frac{1}{2}x</math></p> <p>✓ <math>y \geq \frac{1}{5}x</math></p> <p>✓ Feasible region (5)</p>




1.3	$C = 600x + 300y$ $\therefore 600x + 300y = C$ $\therefore 300y = -600x + C$ $\therefore y = -2x + \frac{C}{300}$ <p>Point A <math>\left(7\frac{19}{23}; 3\frac{21}{23}\right)</math> produces minimum cost.</p> <p>Therefore, the company will need to employ: 8 artisans and 4 apprentices</p>	<p>✓ <math>C = 600x + 300y</math></p> <p>✓ <math>y = -2x + \frac{C}{300}</math></p> <p>✓✓✓✓ <math>\left(7\frac{19}{23}; 3\frac{21}{23}\right)</math></p> <p>Note: Simultaneous equations must be used to obtain the value of x and y</p> <p>✓ 8 and 4 (7)</p>
-----	--	---

[16]

**QUESTION 2**

2.1	$1 \leq x \leq 6$ $1 \leq y \leq 5$ $5x + 4y \leq 40$ $2x + y \geq 8$ $x; y \in \mathbb{N}$	<p>✓ <math>1 \leq x \leq 6</math></p> <p>✓ <math>1 \leq y \leq 5</math></p> <p>✓ <math>5x + 4y \leq 40</math></p> <p>✓ <math>2x + y \geq 8</math></p> <p>✓ <math>x; y \in \mathbb{N}</math></p> <p style="text-align: right;">(5)</p>
2.2	<p>see diagram below</p> 	<p>✓ <math>1 \leq x \leq 6</math></p> <p>✓ <math>1 \leq y \leq 5</math></p> <p>✓ <math>5x + 4y \leq 40</math></p> <p>✓ <math>2x + y \geq 8</math></p> <p>✓ <math>x; y \in \mathbb{N}</math></p> <p style="text-align: right;">(5)</p>
2.3.1	$500x + 250y = P$ $\therefore 250y = -500x + P$ $\therefore y = -2x + \frac{P}{250}$ <p>Max at (6; 2)</p>	<p>✓ <math>500x + 250y = P</math></p> <p>✓ (6; 2)</p> <p style="text-align: right;">(2)</p>
2.3.2	<p>Min at (2; 4) and (3; 2)</p>	<p>✓ (2; 4)</p> <p>✓ (3; 2)</p> <p style="text-align: right;">(2)</p>





[14]

**SECTION C: HOMEWORK**

**QUESTION 1**

A shopkeeper intends buying up to 25 second-hand radios. He has a choice between two types, one without FM for R 30 each and one with FM costing R 40 each.


His total expenditure on the radios must not exceed R 840. If he can sell the first type at a profit of R 10 each and the second type at a profit of R 12 each, determine

- 1.1 The set of inequalities (3)
- 1.2 Sketch the graph and feasible region (5)
- 1.3 How many of each he should purchase to ensure a maximum profit. (4)

[12]

## SECTION D: HOMEWORK SOLUTIONS

1.1	$x + y \leq 25$ $30x + 40y \leq 840$ $x \leq 0$ $y \leq 0$ $x, y \in \mathbb{N}$	$\checkmark x + y \leq 25$ $\checkmark 30x + 40y \leq 840$ $\checkmark x \leq 0, y \leq 0, x, y \in \mathbb{N}$	(3)
1.2	see diagram on next page	$\checkmark x + y \leq 25$ $\checkmark 30x + 40y \leq 840$ $\checkmark x \leq 0$ $\checkmark y \leq 0$ $\checkmark x, y \in \mathbb{N}$	(5)
1.3	$10x + 12y = P$ $\therefore y = -1.2x + \frac{P}{12}$ Intersection $I(x, y)$ of $x + y = 25$ and $30x + 40y = 840$ $\therefore I(x, y) = (16, 9)$  Max at either $I(x, y)$ or $(25, 0)$ Max at $I(x, y), P = 268$  $\therefore x = 16, y = 9$	$\checkmark 10x + 12y = P$ $\checkmark$ Intersection $I(x, y)$ of $x + y = 25$ and $30x + 40y = 840$ $\checkmark$ Check $P$ at $I(x, y)$ and $(25, 0)$ $\checkmark$ Max at $I(x, y), P = 268$	(4)



[12]

## SESSION 18.2

## TOPIC: TRIGONOMETRY

**Teacher Note:** Learners need to know all of the previous Trigonometry they have studied as it will be examined as part of their knowledge in the final exam.

If specifically asked to simplify without a calculator, marks will not be awarded should calculator usage be apparent. The trig identities from grade 11 will not be given and must be learnt

$$\text{i.e. } \sin^2 x + \cos^2 x = 1 \quad \text{and} \quad \tan \theta = \frac{\sin \theta}{\cos \theta}$$

## SECTION A: TYPICAL EXAM QUESTIONS

## QUESTION 1

1.1 Simplify:

$$\frac{\cos(360+x) \cdot \cos(90+x) \cdot \tan^2(x-180)}{\sin(180-x) \cdot \sin(x-360) \cdot \tan(-x)} \quad [5]$$

## QUESTION 2

If  $\sin 12^\circ = k$  determine the following in terms of k:

2.1  $\sin 192^\circ$   (2)

2.2  $\cos 258^\circ$  (3)

2.3  $\sin 336^\circ$  (5)

[10]

## QUESTION 3

3.1 Prove that:  $\frac{\sin 2A + \cos 2A + 1}{\sin A + \cos A} = 2 \cos A$  (6)

3.2 For which values of A is the above identity not defined, for  $A \in [-180; 180]$  (3)  
[9]

## QUESTION 4

Prove without the use of a calculator, that if  $\sin 28^\circ = a$  and  $\cos 32^\circ = b$ , then:


$$b\sqrt{1-a^2} - a\sqrt{1-b^2} = \frac{1}{2} \quad [4]$$

## QUESTION 5

Determine the general solution of:  $\sin^2 x + \cos 2x - \cos x = 0$  [7]

**QUESTION 6**

The graph of  $f(x) = -2\sin x$  is drawn below



- 6.1 Write down the period of  $f$ . (1)
- 6.2 Write down the amplitude of  $h$  if  $h(x) = \frac{f(x)}{4}$ . (2)
- 6.3 Draw the graph of  $g(x) = \cos(x - 30^\circ)$  for  $x \in [-180^\circ; 180^\circ]$ . (3)
- 6.4 Use the graph to determine the number of solutions for  $-2\sin x = \cos(x - 30^\circ)$ ,  $x \in [-180^\circ; 180^\circ]$ . (1)
- 6.5 For which values of  $x$  is  $g(x) \geq 0$ ? (2)
- 6.6 For which values of  $x$  is  $f'(x) < 0$  and  $g'(x) > 0$ ? (3)
- [12]

## SECTION B: SOLUTIONS TO SECTION A

## QUESTION 1

$$\frac{\cos(360+x).\cos(90+x).\tan^2(x-180)}{\sin(180-x).\sin(x-360).\tan(-x)} = \frac{\cos x (-\sin x).\tan^2 x}{\sin x.\sin x.(-\tan x)}$$

$$= \frac{\cos x.\tan x}{\sin x}$$

$$= \frac{\cos x.\sin x}{\sin x.\cos x} = 1 \quad [6]$$

## QUESTION 2

$$2.1 \quad \sin 192 = \sin(180+12) = -\sin 12 = -k \quad (2)$$


$$2.2 \quad \cos 258 = \cos(180+78) = -\cos 78$$

$$= -\cos(90-12) = -\sin 12 = -k \quad (3)$$

$$2.3 \quad \sin 336 = \sin(360-24)$$

$$= -\sin 24 = -\sin(2 \times 12) = -2\sin 12.\cos 12$$

$$= -2k(\sqrt{1-k^2})$$



(5)  
[10]

## QUESTION 3



$$3.1 \quad \frac{\sin 2A + \cos 2A + 1}{\sin A + \cos A} = \frac{2\sin A \cos A + 2\cos^2 A - 1 + 1}{\sin A + \cos A}$$

$$= \frac{2\cos A(\sin A + \cos A)}{(\sin A + \cos A)} = 2\cos A \quad (6)$$

$$3.2 \quad \text{Identity is undefined if } \sin A + \cos A = 0$$

$$\sin A = -\cos A$$

$$\tan A = -1 \quad \cos A \neq 0$$

$$\text{undefined when } A = 135 + k.180 \quad k \in \mathbb{Z}$$

(3)  
[9]

## QUESTION 4

$$b\sqrt{1-a^2} - a\sqrt{1-b^2}$$

$$= \cos 32^\circ.\sqrt{1-\sin^2 28^\circ} - \sin 28^\circ.\sqrt{1-\cos^2 32^\circ}$$

$$= \cos 32^\circ.\cos 28^\circ - \sin 28^\circ.\sin 32^\circ$$

$$= \cos(32^\circ + 28^\circ)$$

$$= \cos 60^\circ$$

$$= \frac{1}{2}$$

[4]

**QUESTION 5**

$$\sin^2 x + \cos 2x - \cos x = 0$$

$$\sin^2 x + (\cos^2 x - \sin^2 x) - \cos x = 0$$

$$\cos^2 x - \cos x = 0$$

$$\cos x(\cos x - 1) = 0$$

$$\cos x = 0 \text{ or } \cos x = 1$$

$$x = \pm 90^\circ + k.360^\circ \text{ or } x = 0^\circ + k.360^\circ \quad k \in Z$$

$$= k.360^\circ$$

(i.e.  $x = 90^\circ + k.180^\circ$  or  $x = k.360^\circ \pm 90^\circ, k \in Z$ )


[7]

**QUESTION 6**

6.1 Period =  $360^\circ$  (1)

6.2 Amplitude =  $\frac{1}{2}$  (2)

6.3 (3)



6.4 2 solutions (1)

6.5  $-60^\circ \leq x \leq 120^\circ$  or  $x \in [-60^\circ; 120^\circ]$  (2)

6.6  $-90^\circ < x < 30^\circ$  or  $x \in (-90^\circ; 30^\circ)$  (3)

[12]

## SECTION B: HOMEWORK

## QUESTION 1

Determine the value of the following without using a calculator:

$$\frac{\sin(-145^\circ) \cdot \cos(-215^\circ)}{\sin 510^\circ \cdot \cos 340^\circ} \quad [8]$$

## QUESTION 2

If  $\sin 2A = \frac{\sqrt{5}}{3}$ , with  $2A \in [90^\circ : 270^\circ]$  determine without the use of a calculator:

2.1  $\cos 2A$  (4)

2.2  $\tan 2A$  (1)

2.3  $\sin A$  (4)

[9]

## QUESTION 3

Simplify completely:  $\frac{\sin(90+\theta) + \cos(180+\theta) \sin(-\theta)}{\sin 180 - \tan 135}$  [5]

## QUESTION 4

Prove that for any angle A:

$$\frac{4 \sin A \cdot \cos A \cdot \cos 2A \cdot \sin 15^\circ}{\sin 2A (\tan 225^\circ - 2 \sin^2 A)} = \frac{\bar{6} - \sqrt{2}}{2} \quad [6]$$



## QUESTION 5

Determine the general solution of:

$$6 \cos x - 5 = \frac{4}{\cos x}; \cos x \neq 0 \quad [6]$$

## QUESTION 6

Determine the value of the following without using a calculator:

$$\cos^4 375^\circ - \sin^4 345^\circ \quad [6]$$

## QUESTION 7

If  $\sin 19^\circ = t$ , express the following in terms of  $t$ . (Make use of a sketch)

7.1  $\sin 79^\circ$  (7)

7.2  $\tan 71^\circ$  (3)

[10]



**SECTION D: HOMEWORK SOLUTIONS**

**QUESTION 1**


$$\begin{aligned} & \frac{\sin(-145^\circ) \cdot \cos(-215^\circ)}{\sin 510^\circ \cdot \cos 340^\circ} \\ &= \frac{(-\sin 145^\circ)(\cos 215^\circ)}{(\sin 150^\circ)(\cos 20^\circ)} \\ &= \frac{(-\sin 35^\circ)(-\cos 35^\circ)}{(\sin 30^\circ)(\cos 20^\circ)} \\ &= \frac{\sin 35^\circ \cos 35^\circ}{\left(\frac{1}{2}\right)(\cos 20^\circ)} \\ &= \frac{2 \sin 35^\circ \cos 35^\circ}{\cos 20^\circ} \\ &= \frac{\sin 70^\circ}{\cos 20^\circ} \\ &= \frac{\cos 20^\circ}{\cos 20^\circ} \\ &= 1 \end{aligned}$$



[8]

**QUESTION 2**

$$\begin{aligned} \sin 2A &= \frac{\sqrt{5}}{3} \\ x^2 &= r^2 - y^2 \\ x^2 &= 3^2 - (\sqrt{5})^2 \\ x^2 &= 4 \\ x &= \pm 2 \\ \therefore x &= -2 \\ \cos 2A &= \frac{-2}{3} \end{aligned}$$



[9]

**QUESTION 3**

$$\begin{aligned} & \frac{\sin(90^\circ + \theta) + \cos(180^\circ + \theta) \sin(-\theta)}{\sin 180^\circ - \tan 135^\circ} \\ &= \frac{\cos \theta + (-\cos \theta)(-\sin \theta)}{0 + 1} \\ &= \cos \theta + \cos \theta \cdot \sin \theta \\ &= \cos \theta(1 + \sin \theta) \end{aligned}$$

[5]

## QUESTION 4

$$\begin{aligned}
 & \frac{4 \sin A \cos A \cos 2A \cdot \sin 15^\circ}{\sin 2A(1 - 2 \sin^2 A)} \\
 &= \frac{4 \sin A \cos A \cos 2A \cdot \sin 15^\circ}{2 \sin A \cos A(1 - 2 \sin^2 A)} \\
 &= \frac{2 \cos 2A \cdot \sin 15^\circ}{\cos 2A} \\
 &= 2 \sin 15^\circ \\
 &= 2 \sin(45^\circ - 30^\circ) \\
 &= 2[\sin 45^\circ \cos 30^\circ - \cos 45^\circ \sin 30^\circ] \\
 &= 2 \left[ \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2} - \frac{\sqrt{2}}{2} \cdot \frac{1}{2} \right] \\
 &= 2 \left[ \frac{\sqrt{6}}{4} - \frac{\sqrt{2}}{4} \right] \\
 &= \frac{\sqrt{6} - \sqrt{2}}{2}
 \end{aligned}$$

[6]

## QUESTION 5

$$\begin{aligned}
 6 \cos x - 5 &= \frac{4}{\cos x} \\
 6 \cos^2 x - 5 \cos x &= 4 \\
 6 \cos^2 x - 5 \cos x - 4 &= 0 \\
 (3 \cos x - 4)(2 \cos x + 1) &= 0 \\
 \cos x = \frac{4}{3} \quad \text{or} \quad \cos x = \frac{-1}{2} \\
 \text{no solution} \quad \text{or} \quad x = 120^\circ + k \cdot 360^\circ, k \in Z \\
 &\text{or} \\
 x &= 240^\circ + k \cdot 360^\circ, k \in Z
 \end{aligned}$$

$$\text{Alternative solution for } \cos x = \frac{-1}{2}$$

$$x = k \cdot 360^\circ \pm 120^\circ \quad k \in Z$$

**Note:**

If candidate puts  $\pm k \cdot 360$  then  $k \in \mathbb{N}_0$

[6]

**QUESTION 6**

$$\begin{aligned} & \cos^4 375^\circ - \sin^4 345^\circ \\ &= \cos^4 15^\circ - \sin^4 15^\circ \\ &= (\cos^2 15^\circ + \sin^2 15^\circ)(\cos^2 15^\circ - \sin^2 15^\circ) \\ &= (1)(\cos 30^\circ) \\ &= \frac{\sqrt{3}}{2} \end{aligned}$$

[6]

**QUESTION 7**

7.1

$$\begin{aligned} \sin 19^\circ &= \frac{t}{1} \\ x^2 + t^2 &= 1^2 \\ \therefore x^2 &= 1 - t^2 \\ \therefore x &= \sqrt{1 - t^2} \\ \sin 79^\circ &= \sin(19^\circ + 60^\circ) \\ &= \sin 19^\circ \cos 60^\circ + \cos 19^\circ \sin 60^\circ \\ &= (t) \left( \frac{1}{2} \right) + \left( \frac{\sqrt{1 - t^2}}{1} \right) \left( \frac{\sqrt{3}}{2} \right) \\ &= \frac{t + \sqrt{3}\sqrt{1 - t^2}}{2} = \frac{t + \sqrt{3 - 3t^2}}{2} \end{aligned}$$



(7)

7.2

$$\begin{aligned} \tan 71^\circ &= \frac{\sin 71^\circ}{\cos 71^\circ} \\ &= \frac{\cos 19^\circ}{\sin 19^\circ} \\ &= \frac{\sqrt{1 - t^2}}{1} \\ &= \frac{1}{t} \\ &= \frac{\sqrt{1 - t^2}}{t} \end{aligned}$$

(3)

[10]

SESSION 19.1


TOPIC: 2D TRIGONOMETRY

**Teacher Note:** Learners must revise the sine, cos and area rules to help them 'solve' triangles.

**SECTION A: TYPICAL EXAM QUESTIONS**

**QUESTION 1**

AB and CD are two towers in the same horizontal plane, the angle of depression of D from A is  $\alpha$  and the angle of elevation of C from B is  $\beta$ .




1.1 If  $AB = x$  prove that  $CD = \frac{x \tan \beta}{\tan \alpha}$  (4)

1.2 Hence find CD without a calculator when  $x = 50\sqrt{3}$ ,  $\beta = 45$  and  $\alpha = 30$  (2)  
[6]

**QUESTION 2**

In the diagram below; AC is a diameter of the circle with  $AB = x$ ,  $AC = BC = y$  and  $\angle ABC = \alpha$  and  $\angle ADC = 90$




a) Show that  $\cos \alpha = \frac{x}{2y}$  (3)

b) Determine DC in terms of x and y (4)  
[7]

**QUESTION 3**

From a point P, on top of a building the angle of elevation to the top of a cell phone tower is  $24^\circ$  and the angle of depression to the foot of the tower is  $19^\circ$ , as shown below:




If the height of the tower is 28m how far is the building from the tower, if they lie in the same horizontal plane? (Round off to the nearest whole number). [8]

**QUESTION 4**

A piece of land has the form of a quadrilateral ABCD with  $AB = 20\text{m}$ ,  $BC = 12\text{m}$ ,  $CD = 7\text{m}$  and  $AD = 28\text{m}$ ,  $\hat{B} = 110^\circ$

The owner decides to divide the land into two plots by erecting a fence from A to C.



- (a) Calculate the length of the fence AC correct to one decimal place. [2]
  - (b) Calculate the size of  $\hat{BAC}$  correct to the nearest degree. [2]
  - (c) Calculate the size of  $\hat{D}$ , correct to the nearest degree. [3]
  - (d) Calculate the area of the entire piece of land ABCD, correct to one decimal place. [3]
- [10]

**QUESTION 5**

$\triangle ABC$  is an isosceles triangle with  $AB = BC = m$ ,  $AB = c$ ,  $AC = b$  and  $BC = a$ .

Prove that  $\cos B = 1 - \frac{b^2}{2a^2}$  [4]

**SECTION B: SOLUTIONS TO SECTION A****QUESTION 1**

1.1  $\tan \alpha = x/DB$   
 $DB = x/\tan \alpha$

In  $\triangle CBD$ :  $\tan \beta = CD/DB$   
 $CD = DB \tan \beta = \frac{x \tan \beta}{\tan \alpha}$  (4)

1.2  $CD = \frac{50\sqrt{3} \tan 45}{\tan 30} = \frac{50\sqrt{3} \left(\frac{1}{\sqrt{2}}\right)}{\frac{1}{\sqrt{3}}} = 150 \left(\frac{1}{\sqrt{2}}\right) = \frac{150\sqrt{2}}{2} = 75\sqrt{2}$  (2)

[6]

**QUESTION 2**

(a)  $AC^2 = AB^2 + BC^2 - 2AB \cdot BC \cos \alpha$   
 $y^2 = x^2 + y^2 - 2xy \cos \alpha$   
 $-2xy \cos \alpha = -x^2$   
 $\therefore \cos \alpha = \frac{x^2}{2xy} = \frac{x}{2y}$  (3)



(b) In  $\triangle ABD$ :  $\cos \alpha = BD/AB$   
 $BD = AB \cos \alpha = x \left(\frac{x}{2y}\right) = \frac{x^2}{2y}$   
 $\therefore DC = y - \frac{x^2}{2y} = \frac{2y^2 - x^2}{2y}$  (4)

[7]

**QUESTION 3**

In  $\triangle PQS$ :  $\angle QSP = 71^\circ$ ,  $QS \perp SO$ ,  $\angle Q = 66^\circ$  (angles in triangle PQS)

$$\frac{28}{\sin 43} = \frac{PS}{\sin 66}$$

$$PS = \frac{28 \sin 66}{\sin 43} = 38m$$

In  $\triangle PSO$ :  $\cos 19 = \frac{OS}{PS}$

$$\therefore OS = 38 \cos 19 = 36$$

[8]

**QUESTION 4**

(a)

$$AC^2 = (12m)^2 + (20m)^2 - 2(12m)(20m) \cos 110^\circ$$

$$\therefore AC^2 = 708,1696688$$

$$\therefore AC = 26,6m$$

(3)

b)

$$\frac{\sin \hat{BAC}}{12m} = \frac{\sin 110^\circ}{26,6m}$$

$$\therefore \sin \hat{BAC} = \frac{12 \times \sin 110^\circ}{26,6m}$$

$$\therefore \sin \hat{BAC} = 0,4239214831$$

$$\therefore \hat{BAC} = 25^\circ$$

(3)

**OR**

$$(12m)^2 = (20m)^2 + (26,6m)^2 - 2(20m)(26,6m) \cos \hat{BAC}$$

$$\therefore 1064 \cos \hat{BAC} = 963,56m^2$$

$$\therefore \cos \hat{BAC} = 0,9056015038$$

$$\therefore \hat{BAC} = 25^\circ$$

c)

$$(26,6m)^2 = (7m)^2 + (28m)^2 - 2(7m)(28m) \cos \hat{D}$$

$$\therefore 392 \cos \hat{D} = 125,44$$

$$\therefore \cos \hat{D} = 0,32$$

$$\therefore \hat{D} = 71^\circ$$

(2)

d)

Area ABCD

$$= \frac{1}{2}(12m)(20m) \sin 110^\circ + \frac{1}{2}(7m)(28m) \sin 71^\circ$$

$$= 205,4m^2$$

(2)

**[10]**

**QUESTION 5**

$$b^2 = a^2 + c^2 - 2ac \cos B$$


$$\therefore b^2 = a^2 + a^2 - 2a \times a \times \cos B$$

$$\therefore b^2 = 2a^2 - 2a^2 \cos B$$

$$\therefore b^2 = 2a^2(1 - \cos B)$$

$$\therefore \frac{b^2}{2a^2} = 1 - \cos B$$

$$\therefore \cos B = 1 - \frac{b^2}{2a^2}$$



**[4]**

**SECTION C: HOMEWORK**


**QUESTION 1**

A mathematician stands on a bank of a river and finds that the angle of elevation to the top of a tree on the opposite bank is  $18,3^\circ$ . If she moves 45m backwards in line with her first position and the tree, she finds the angle of elevation is now  $13,7^\circ$ . Calculate the height of the tree and the width of the river (to the nearest metre).

(Tip: A diagram must be used in this question)

**QUESTION 2**


Two ships, A and B, are 120 km apart. Ship A is at a bearing of  $67^\circ$  from D and 97 km away from D. DN points due north. Ship B is at a bearing of  $208^\circ$  from D.



- (a) Determine the bearing of Ship A from Ship B ( $\angle MBA$ ), when  $BM \parallel DN$
- (b) If Ship B travels due north, and Ship B travels due south, then at some instant of time, Ship A is due east of Ship B. Calculate the distance between the two ships at that instant.

**QUESTION 3**


If  $b=c$  and  $a^2 = 7b^2$ , show why it is impossible to construct  $\triangle ABC$ .





**SECTION D: SOLUTIONS TO HOMEWORK**

**QUESTION 1**



In  $\triangle CDA$   $\angle DAC = 180 - 18,3 = 161,7^\circ$   
 $\angle DCA = 180 - (13,7 + 161,7) = 4,6^\circ$

$$\frac{AC}{\sin 13,7} = \frac{45}{\sin 4,6}$$

$$\therefore AC = \frac{45 \sin 13,7}{\sin 4,6} = 133m$$

In  $\triangle ABC$   $\sin 18,3 = \frac{BC}{AC} = \frac{BC}{132,89}$

$$\therefore BC = 132,89 \times \sin 18,3 = 42m$$

Tree is 42m



Using Pythagoras:  $AB = \sqrt{((132,89 \dots)^2 - (41,7 \dots)^2)} = 126m = \text{width of the river}$

**QUESTION 2**

2a)

$$\hat{NDB} = 360^\circ - 208^\circ = 152^\circ$$

$$\therefore \hat{MBD} = 28^\circ$$

$$\hat{BDA} = 208^\circ - 67^\circ = 141^\circ$$

$$\frac{\sin \hat{DBA}}{97} = \frac{\sin 141^\circ}{120}$$

$$\therefore \hat{DBA} = 30,58^\circ$$

$$\therefore \sin \hat{DBA} = \frac{97 \sin 141^\circ}{120}$$


$$\therefore \hat{MBA} = 30,58^\circ + 28^\circ$$

$$\therefore \sin \hat{DBA} = 0,5087006494$$

$$\therefore \hat{MBA} = 58,58^\circ$$

2b)

$$\hat{B} = 30,58^\circ$$



$$\frac{EA}{120} = \sin(28^\circ + 30,58^\circ)$$

$$EA = 120 \sin(28^\circ + 30,58^\circ)$$

$$EA = 102,4 \text{ km}$$

**QUESTION 3**

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\therefore 7b^2 = b^2 + b^2 - 2b^2 \cos A$$

$$\therefore 5b^2 = -2b^2 \cos A$$

$$\therefore \frac{5b^2}{-2b^2} = \cos A$$

$$\therefore \cos A = -\frac{5}{2}$$

This equation has no solution since  $-1 \leq \cos A \leq 1$



## SESSION 24 SELF STUDY

## TOPIC: 3D TRIGONOMETRY



**Teacher Note:** Encourage learners to take time to understand all given information, including the diagram. They must fill in as much information as possible on the diagram and then work from the triangle with the most information. Learners must revise 2D Trigonometry as they will need all the rules learned thus far to solve triangles.

## SECTION A: TYPICAL EXAM TYPE QUESTIONS


## QUESTION 1

In the figure A, B & C are three points in the horizontal plane, such that  $\angle ABC = 120^\circ$ .

D is a point directly above A

$AD = BC = y$

The angle of elevation of D from B is  $\theta$



a) Show that  $AC = y\sqrt{\left(\frac{1}{\tan^2 \theta} + \frac{1}{\tan \theta} + 1\right)}$  (7)

b) If  $y=15$  and  $\theta = 22$ , calculate:


i) AC (2)

ii)  $\angle ADC$  (3)

[12]

**QUESTION 2**

In the diagram below, AB is a straight line 1 500 m long. DC is a vertical tower 158 metres high with C, A and B points in the same horizontal plane. The angles of elevation of D from A and B are  $25^\circ$  and  $\theta$ .  $\hat{CAB} = 30^\circ$ .



- (a) Determine the length of AC. (3)
  - (b) Find the value of  $\theta$ . (5)
  - (c) Calculate the area of  $\triangle ABC$ . (2)
  - (d) Calculate the size of  $\hat{ADB}$ . (6)
- [16]**




**QUESTION 3**

(DOE Nov 2010 P2)

The angle of elevation from a point C on the ground, at the centre of the goalpost, to the highest point A of the arc, directly above the centre of the Moses Mabida soccer stadium, is  $64,75^\circ$ .

The soccer pitch is 100 metres long and 64 metres wide as prescribed by FIFA for world cup stadiums. Also  $AC \perp PC$ . In the figure below  $PQ = 100$  metres and  $PC = 32$  metres.




MATHEMATICS      GRADE 12      SESSION 19 SELF STUDY      (TEACHER NOTES)

- 3.1 Determine AC (3)
- 3.2 Calculate  $\angle PAC$  (3)
- 3.3 A camera is positioned at point D, 40 metres directly below A. Calculate the distance from D to C. (4)  
[10]

**QUESTION 4***(DOE March 2011)*

The sketch below shows one side of the elevation of a house. Some dimensions (in metres) are indicated on the figure.



Calculate, rounded off to ONE decimal place:

- 4.1 EC (3)
- 4.2  $\hat{DCE}$  (3)
- 4.3 Area of  $\triangle DEC$  (2)
- 4.4 The height EF (3)  
[11]

## SECTION B: SOLUTIONS TO SECTION A

**QUESTION 1**(a) In  $\triangle ADB$ 

$$\angle ADB = 90 - \theta \quad \tan \theta = \frac{DA}{AB} = \frac{y}{AB} \quad \therefore AB = \frac{y}{\tan \theta}$$

In  $\triangle ABC$ 

$$\begin{aligned} AC^2 &= AB^2 + BC^2 - 2AB \cdot BC \cos 120 = \left(\frac{y}{\tan \theta}\right)^2 + y^2 - 2\left(\frac{y}{\tan \theta}\right)(y)(-\cos 60) \\ &= y^2 \frac{1}{\tan^2 \theta} + y^2 + y^2 \left(\frac{1}{\tan \theta}\right) \\ &= y^2 \left[\left(\frac{1}{\tan^2 \theta}\right) + 1 + \frac{1}{\tan \theta}\right] \end{aligned}$$

$$\therefore AC = \sqrt{y^2 \left[\left(\frac{1}{\tan^2 \theta}\right) + 1 + \frac{1}{\tan \theta}\right]} = y \sqrt{\left[\left(\frac{1}{\tan^2 \theta}\right) + 1 + \frac{1}{\tan \theta}\right]} \quad (7)$$

(b)

$$(i) \quad AC = 46,4\text{m} \quad (2)$$

$$(ii) \quad \text{In } \triangle ADC: \tan(\angle ADC) = \frac{AC}{AD} = \frac{46,48}{15} = 3,0985 \dots$$

$$\angle ADC = 72,11^\circ \quad (3)$$

**[12]****QUESTION 2**(a) In  $\triangle ADC$ :

$$\hat{D} = 65^\circ \quad (\angle \text{ of } \triangle)$$

$$\frac{AC}{\sin 65^\circ} = \frac{158}{\sin 25^\circ}$$

$$\therefore AC \cdot \sin 25^\circ = 158 \cdot \sin 65^\circ$$

$$\therefore AC = \frac{158 \cdot \sin 65^\circ}{\sin 25^\circ}$$

$$\therefore AC = 338,83\text{m} \quad (3)$$

(b) In  $\triangle ACB$ :

$$BC^2 = 338,83^2 + 1500^2 - 2(338,83)(1500)\cos 30^\circ$$

$$\therefore BC^2 = 1\,484\,499,606$$

$$\therefore BC = 1218,4\text{m}$$

In  $\triangle DCB$ :

$$\tan \theta = \frac{DC}{BC}$$

$$\therefore \tan \theta = \frac{158}{1218,4}$$

$$\therefore \theta = 7,39^\circ \quad (5)$$

$$(c) \quad \text{Area } \triangle ABC = \frac{1}{2}(338,83)(1500) \sin 30^\circ$$

$$\therefore \text{Area } \triangle ABC = 127061,25m^2 \quad (2)$$

$$(d) \quad AD^2 = (338,83)^2 + (158)^2$$

$$\therefore AD^2 = 139769,7689$$

$$\therefore AD = 373,86m$$

$$BD^2 = (1218,4)^2 + (158)^2$$

$$\therefore BD^2 = 1509462,56$$

$$\therefore BD = 1228,60m$$

$$(1500)^2 = (373,86)^2 + (1228,60)^2 - 2(373,86)(1228,60) \cos \hat{A}DB$$

$$\therefore 2(373,86)(1228,60) \cos \hat{A}DB = (373,86)^2 + (1228,60)^2 - (1500)^2$$

$$\therefore 918648,792 \cos \hat{A}DB = -600770,7404$$

$$\therefore \cos \hat{A}DB = -0,6539721661$$

$$\therefore \hat{A}DB = 130,84^\circ \quad (6)$$

**[16]**

**QUESTION 3**


$$3.1 \quad \cos 64,75^\circ = \frac{50}{AC}$$

$$AC = \frac{50}{\cos 64,75^\circ}$$

$$= 117,21m \quad (3)$$



$$3.2 \quad PC \text{ is given to be } \frac{1}{2}(64) = 32m$$



$$\tan \hat{P}AC = \frac{32}{117,21}$$

$$\theta = 15,27^\circ \quad (15,27042173\dots)$$


**Note:** If the candidate takes the unrounded answer for AC, then the answer is  $15,27^\circ$  (15.26987495...) (3)

$$\begin{aligned}
 3.3 \quad CD^2 &= 117,21^2 + 40^2 - 2(117,21)(40)\cos 25,25 \\
 &= 6857,289092 \\
 \therefore CD &= 82,81 \text{ m}
 \end{aligned}$$

**Note:**

If don't use the rounded off then  $CD = 82,81 \text{ m}$ . Accept this answer.

**OR**



$$\begin{aligned}
 AM &= AC \sin 64,75^\circ \quad \text{OR} \quad AM = CM \tan 64,75^\circ \quad \text{OR} \quad AM = AC \cos 25,25^\circ \\
 &= 106,0111876 \qquad \qquad \qquad = 50 \tan 64,75^\circ \qquad \qquad \qquad = 117,21 \cdot \cos 25,25^\circ \\
 &= 106,01 \qquad \qquad \qquad = 106,01 \qquad \qquad \qquad = 106,01 \\
 DM &= 106,01 - 40 \\
 &= 66,01 \\
 CD^2 &= CM^2 + DM^2 \\
 &= (50)^2 + (66,01)^2 \\
 &= 6857,3201 \\
 CD &= 82,81 \text{ metres}
 \end{aligned}$$

(4)

**[10]**

**QUESTION 4**

$$\begin{aligned}
 4.1 \quad EC^2 &= DE^2 + DC^2 - 2DE \cdot DC \cos \hat{C} \\
 &= (7,5)^2 + (9,4)^2 - 2 \cdot (7,5)(9,4) \cos 32^\circ \\
 &= 25,03521844... \\
 EC &= 5,0 \text{ metres}
 \end{aligned}$$

(3)

$$\begin{aligned}
 4.2 \quad \frac{\sin DCE}{7,5} &= \frac{\sin 32^\circ}{5,0} \\
 \sin \hat{DCE} &= \frac{7,5 \cdot \sin 32^\circ}{5,0} \\
 &= 0,7948788963 \\
 \hat{DCE} &= 52,6^\circ
 \end{aligned}$$

(3)



$$\begin{aligned}
 4.3 \quad & \text{Area of } \triangle DEC \\
 &= \frac{1}{2} DE \cdot DC \sin \hat{D} \\
 &= \frac{1}{2} (7,5)(9,4) \sin 32^\circ \\
 &= 18,7m^2
 \end{aligned}$$

OR

Area of  $\triangle DEC$

$$\begin{aligned}
 &= \frac{1}{2} CE \cdot DC \sin 52,6^\circ \\
 &= \frac{1}{2} (5,0)(9,4) \sin 52,6^\circ \\
 &= 18,7m^2
 \end{aligned}$$

(2)

$$\begin{aligned}
 4.4 \quad & \sin 32^\circ = \frac{EG}{7,5} \\
 & EG = 7,5 \cdot \sin 32^\circ \\
 & \quad = 4,0 \\
 & EF = (4 + 3,5) \\
 & \quad = 7,5 \text{ metres}
 \end{aligned}$$



OR

$$\begin{aligned}
 EG &= EC \cdot \sin 52,6^\circ \\
 &= (5,0) \cdot \sin 52,6^\circ \\
 &= 4,0 \\
 EF &= 4,0 + 3,5 \\
 &= 7,5
 \end{aligned}$$


(3)

[11]

**SECTION C: SOLUTIONS TO HOMEWORK**

**QUESTION 1**

In the diagram below the points ABC lie in the same horizontal plane. A hot air balloon is stationary at point P, which is directly above C. The angle of elevation of P from A is  $\theta$ .  $\angle ABC = 90 - \alpha$  and the distance from B to C is half the distance of BA, where BA is d units.




a) Show that the height of the balloon above C is  $h = \frac{d\sqrt{(5-\sin\alpha)}}{2} \tan\theta$

b) Calculate h if  $d = 300$ ,  $\alpha = 32$  and  $\theta = 63$

**QUESTION 2**



In the figure A,B & C are three points in the same horizontal plane. AD represents a lamp pole that is perpendicular to the horizontal plane.



Given that  $\angle BDA = \angle ABC = \theta$  and  $\angle BCA = \beta$ ,  $BC = x$

a) Write  $\angle BAC$  in terms of  $\theta$  and  $\beta$

b) Show that  $AB = \frac{x \sin \beta}{\sin(\theta + \beta)}$


c) IF  $AB = AC$  show:

i)  $AB = \frac{x}{2 \cos \theta}$

ii)  $AD = \frac{x}{2 \sin \theta}$

### QUESTION 3

Thandi is standing at point P on the horizontal ground and observes two poles, AC and BD, of different heights. P, C and D are in the same horizontal plane. From P the angles of inclination to the top of the poles A and B are  $23^\circ$  and  $18^\circ$  respectively. Thandi is 18 m from the base of pole AC. The height of pole BD is 7 m.




Calculate, correct to TWO decimal places:

- The distance from Thandi to the top of pole BD.
- The distance from Thandi to the top of pole AC.
- The distance between the tops of the poles, that is the length of AB, if  $\hat{APB} = 42^\circ$

**QUESTION 4**

A rectangular block of wood has a breadth of 6 metres, height of 8 metres and a length of 15 metres. A plane cut is made through the block as shown in the diagram revealing the triangular plane that has been formed. Calculate the size of  $\hat{E}BG$ .



**SECTION D: SOLUTIONS TO HOMEWORK**

**QUESTION 1**



a) In  $\triangle ABC$

$$AC^2 = AB^2 + BC^2 - 2AB \cdot BC \cos(90 - \alpha)$$

$$= d^2 + \left(\frac{1}{2}d\right)^2 - 2d\left(\frac{1}{2}d\right)\sin\alpha$$

$$= \frac{5}{4}d^2 - d^2\sin\alpha = d^2\left(\frac{5}{4} - \sin\alpha\right)$$

$$\therefore AC = \frac{d\sqrt{(5-\sin\alpha)}}{2}$$

In  $\triangle ACP$

$$\tan\theta = \frac{PC}{AC}$$

$$PC = h = AC \tan\theta = \frac{d\sqrt{(5-\sin\alpha)}}{2} \tan\theta$$

b)  $h = \frac{300(\sqrt{5-4\sin 32})}{2} \tan 63 = 500m$

**QUESTION 2**

a)  $\angle BAC = 180 - (\theta + \beta)$

b) 
$$\frac{AB}{\sin \beta} = \frac{\text{opposite}}{\sin(180 - (\theta + \beta))}$$

$$AB = \frac{x \sin \beta}{\sin(\theta + \beta)}$$

c) i) IF  $AB = AC$  Then  $\theta = \beta$   

$$AB = \frac{x \sin \theta}{\sin 2\theta} = \frac{x \sin \theta}{2 \sin \theta \cos \theta} = \frac{x}{2 \cos \theta}$$

ii) In  $\triangle BDA$   
 $\angle B = 90 - \theta$   

$$\frac{AB}{\sin \theta} = \frac{AD}{\sin(90 - \theta)} \quad \therefore AD = \frac{\cos \theta \left( \frac{x}{2 \cos \theta} \right)}{\sin \theta} = \frac{x}{2 \sin \theta}$$

**QUESTION 3**

a)

$$\frac{7}{PB} = \sin 18^\circ$$

$$\therefore PB = \frac{7}{\sin 18^\circ}$$

$$\therefore PB = 22,65247584..$$



b)

$$\frac{18}{PA} = \cos 23^\circ$$

$$\therefore PA = \frac{18}{\cos 23^\circ}$$

$$\therefore PA = 19,55448679....$$

c)

$$AB^2 = (22,65)^2 + (19,55)^2 - 2(22,65)(19,55) \cdot \cos 42^\circ$$

$$\therefore AB^2 = 237,0847954...$$

$$\therefore AB = 15,40 \text{ m}$$

## QUESTION 4

In  $\triangle AEB$ :

$$EB^2 = 8^2 + 6^2$$

$$\therefore EB^2 = 100$$

$$\therefore EB = 10$$

In  $\triangle GBC$ :

$$BC^2 = 15^2 + 8^2$$

$$\therefore BC^2 = 289$$

$$\therefore BC = 17$$

In  $\triangle ACB$ :

$$EG^2 = 15^2 + 6^2$$

$$\therefore EG^2 = 261$$

$$\therefore EG = \sqrt{261}$$

In  $\triangle EGB$ :

$$\therefore (\sqrt{261})^2 = 17^2 + 10^2 - (2(17)(10)\cos \hat{E}BG)$$

$$\therefore 261 = 389 - (340\cos \hat{E}BG)$$

$$\therefore -128 = -340\cos \hat{E}BG$$

$$\therefore \frac{32}{85} = \cos \hat{E}BG$$

$$\therefore \hat{E}BG = 67,88^\circ$$

