Download more resources like this on ECOLEBOOKS.COM



# NATIONAL SENIOR CERTIFICATE

# GRADE 12

# **SEPTEMBER 2020**

# **MATHEMATICS P2**

**MARKS: 150** 

TIME: 3 hours

This question paper consists of 15 pages, including a 1-page information sheet and an answer book of 25 pages.

#### **INSTRUCTIONS AND INFORMATION**

Read the following instructions carefully before answering the questions.

- 1. This question paper consists of 10 questions.
- 2. Answer ALL the questions in the SPECIAL ANSWER BOOK provided.
- 3. Clearly show ALL calculations, diagrams, graphs, et cetera that you have used in determining the answers.
- 4. Answers only will NOT necessarily be awarded full marks.
- 5. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
- 6. If necessary, round off answers to TWO decimal places, unless stated otherwise.
- 7. Diagrams are NOT necessarily drawn to scale.
- 8. An information sheet with formulae is included at the end of the question paper.
- 9. Write neatly and legibly.



(EC/SEPTEMBER 2020)

MATHEMATICS P2

## **QUESTION 1**

The following table shows a comparison of a school's Grade 12 final marks in 2019 and the learners' School Based Assessment (SBA) marks for the year.

LE	ARNERS	1	2	3	4	5	6	7	8	9	10	
SB.	A MARK	99	93	77	74	63	62	63	63	47	37	
FIN	NAL MARK	94	81	73	65	59	58	55	49	43	31	
1.1	Determine the ec answer correct to	quation 0 4 deci	of the	least sc aces.)	juares 1	egressi	ion line	e for the	e data.	(Round	off you	ır (
1.2	Determine the correlation coefficient between the SBA mark and the final mark.						(					
1.3	Comment on the	correla	ation be	etween	the SB	A marl	k and th	ne final	mark.			
1.4	Learner 11 scored 51% for SBA. Predict the final mark he should get, correct to the nearest unit.						ie					
1.5	Given that the m one deviation of	ean for the me	the fin an.	al mar	k is 60,	8, calc	ulate ho	ow mar	ny learr	ners we	re withi	n ( [1

### **QUESTION 2**

The speeds, in kilometres per hour, of cyclists that passed a point on the route of the Ironman Race were recorded and summarised in the table below:

ÉcoleBooks					
Speed (km/h)	Frequency (f)	Cumulative Frequency			
$0 < x \le 10$	10	10			
$10 < x \le 20$		30			
$20 \! < \! x \! \le \! 30$	45				
$30 < x \le 40$	72				
$40 < x \le 50$		170			

2.1	Complete the above table in the ANSWER BOOK provided.	(2)
2.2	Make use of the axes provided in the ANSWER BOOK to draw a cumulative frequency curve for the above data.	(3)
2.3	Indicate clearly on your graph where the estimates of the lower quartile $(Q_1)$ and median (M) speeds can be read off. Write down these estimates.	(2)
2.4	Draw a box and whisker diagram for the data. Use the number line in the ANSWER BOOK.	(2)
2.5	Use your graph to estimate the number of cyclists that passed the point with speeds greater than 35 km/h.	(1) [ <b>10</b> ]

3

In the diagram, P, Q (0; -2) and R (6; -4) are the vertices of triangle PQR. The equation of PQ is 3x - y - 2 = 0. The equation of PR is y = -x + 2. RS is the perpendicular from R to the *y*-axis. QPR =  $\theta$ .



In the diagram below, C is the centre of the circle defined by  $x^2 - 6x + y^2 - 4y + 9 = 0$ . T (-1; y) is a point outside the circle. Two tangents are drawn to the circle from T. STB is tangent to the circle at B and has equation y = -2x + k. TA is tangent to the circle at A and is parallel to the x-axis. BV is a diameter of the circle.



4.6	Calculate the size of $\widehat{ACB}$ . Give reason(s).	(4) [ <b>18</b> ]
4.5	Determine the value of <i>k</i> .	(2)
4.4	Calculate the length of TB. Give reason(s).	(4)
4.3	Determine the equation of line TA.	(1)
4.2	Determine the equation of BV.	(3)
4.1	Determine the coordinates of C.	(4)

5

- 5.1 If  $\cos 22^\circ = p$ ; determine the following in terms of *p*:
  - $5.1.1 \cos 158^{\circ}$  (2)

      $5.1.2 \sin 112^{\circ}$  (2)

      $5.1.3 \sin 38^{\circ}$  (4)
- 5.2 Determine all the values of P in the interval  $[0^\circ; 360^\circ]$  which satisfy the equation: sin P = sin 2P. (4)
- 5.3 If  $\triangle ABC$  is a scalene triangle, show that:  $\cos(A + B) = -\cos C$ . (2)
- 5.4 Prove the following identity:

$$\frac{\cos^2 x - \cos x - \sin^2 x}{2\sin x \cdot \cos x + \sin x} = \frac{1}{\tan x} - \frac{1}{\sin x}$$
(5)

5.5 Determine the general solution of:  $4 + 7\cos\theta + \cos 2\theta = 0$ . (6)

[25]



In the diagram below, the graphs of  $f(x) = \tan bx$  and  $g(x) = \cos(x - 30^\circ)$  are drawn on the same set of axes for  $-180^\circ \le x \le 180^\circ$ . The points P(90°;1) and Q lie on f and g respectively. Use the diagram to answer the following questions.



Points L, M and N are in the same horizontal plane. KL is a vertical tower. The angle of elevation of K from M is  $q^{\circ}$ . NL =  $p^{\circ}$ ; NL = NM = d and KL = h.



7.1	Determine the size of $L\widehat{N}M$ in terms of <i>p</i> .	(2)
7.2	Prove that $LM = \frac{d \sin 2p}{\sin p}$ .	(2)
7.3	Hence, show that $h = 2d \cos p \tan q$ .	(3) [ <b>7</b> ]

8.1 Complete the following theorem statement:

The line drawn from the centre of a circle perpendicular to a chord ...

8.2 In the diagram below, circle ABC with centre O is given. OB = 8 units and AB = BC = 10 units. E is the midpoint of AC. Let OE = y and AE = x.



Calculate, with reasons, the length of OE.

8.3 Complete the following theorem statement:

The angle subtended by an arc at the centre of a circle is ... at the circle (on the same side of the chord as the centre).

(5)

(1)

9

(1)

8.4 In the diagram, O is the centre of a circle ABCD. AOD is the diameter and OC is a radius. AB, BC, CD, AC and BD are straight lines.



Write down, with reasons, an equation that expresses the relationship between each of the given groups of angles.

	ANGLES	EQUATION / RELATIONSHIP	REASON	
e.g.	$\widehat{M}_3; \widehat{P}$	$\hat{\mathbf{M}}_3 = 2 \times \hat{\mathbf{P}}$	$\angle$ at centre = 2 $\lor \angle$ at circum.	
8.4.1	$\hat{O}_2; \hat{B}_2$			
8.4.2	$\widehat{D}_1; \widehat{C}_3; \widehat{D}_2$			
8.4.3	$\hat{B}_1; \hat{B}_2; \hat{D}_1; \hat{D}_2$			
8.4.4	$\widehat{D}_1; \widehat{C}_1$			(8
	•		·	[15

9.1 Given that PC is a tangent to the circle ACB; BAP and BCQ are straight lines. PC = PQ and  $C\hat{P}Q = 20^{\circ}$ .



Prove, stating reasons, that BC is NOT a diameter.

(5)

9.2 In the diagram below, O is the centre of circle ABC. The tangent PA to the circle and diameter AB meets at A. OCQ and BCP are straight lines. PQ||AB.



Prove, stating reasons, that PQ = QC.

(6)

9.3 In the diagram below, chords AB and FE of circles with centre O are produced to points C and D. BE||CD.



Prove that ACDF is a cyclic quadrilateral.

(5) [**16**]



10.1 In the diagram,  $\Delta$ MON is drawn. S is a point on MO and T is a point on MN such that ST||ON. SN and OT are drawn.



Use the diagram to prove the theorem which states that a line parallel to one side of a triangle divides the other two sides proportionally. In other words, prove that:  $\frac{MS}{SO} = \frac{MT}{TN}.$ 

(5)

10.2 In the diagram, O is the centre of the circle. PQRS is a cyclic quadrilateral. The tangent through P intersects RS produced at A. OB⊥PR and PA⊥AS.



#### Prove that:

		TOTAL:	150
10.2.4	$BR \cdot RQ = RS \cdot RP$		(6) [ <b>19</b> ]
10.2.3	$\mathbf{P}_4 = \mathbf{K}_2$		(4)
1023			
10.2.2	$AP \cdot RS = BR \cdot PS$		(1)
10.2.1	$\Delta APS \parallel \Delta BRS$		(3)

Copyright reserved

(EC/SEPTEMBER 2020)

# **INFORMATION SHEET: MATHEMATICS**

**MATHEMATICS P2** 

<u>15</u>

$$x = \frac{-b \pm \sqrt{b^{2} - 4ac}}{2a}$$

$$A = P(1+ni) \qquad A = P(1-ni) \qquad A = P(1-i)^{n} \qquad A = P(1+i)^{n}$$

$$\sum_{i=1}^{n} 1 = n \qquad \sum_{i=1}^{n} i = \frac{n(n+1)}{2} \qquad T_{n} = a + (n-1)d \qquad S_{n} = \frac{n}{2}(2a + (n-1)d)$$

$$T_{n} = ar^{n-1} \qquad S_{n} = \frac{a(r^{n}-1)}{r-1} ; \qquad r \neq 1 \qquad S_{n} = \frac{a}{1-r} ; -1 < r < 1$$

$$F = \frac{x[(1+i)^{n}-1]}{i} \qquad P = \frac{x(1-(1+i)^{-n})}{i}$$

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

$$d = \sqrt{(x_{2}-x_{1})^{2} + (y_{2}-y_{1})^{2}} \qquad M\left(\frac{x_{1}+x_{2}}{2}; \frac{y_{1}+y_{2}}{2}\right)$$

$$y = mx + c \qquad y - y_{1} = m(x-x_{1}) \qquad m = \frac{y_{2}-y_{1}}{x_{2}-x_{1}} \qquad m = \tan\theta$$

$$(x-a)^{2} + (y-b)^{2} = r^{2}$$

$$In \ ABC: \qquad \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} \qquad a^{2} = b^{2} + c^{2} - 2bc \cos A \qquad area \ ABC = \frac{1}{2}ab \sin C$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta \qquad \sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \qquad \sin(\alpha - \beta) = \sin \alpha \cos \beta$$

$$\frac{1}{2\cos^{2}\alpha - 1} \qquad \sigma^{2} = \frac{\sum_{i=1}^{n} (x_{i} - \overline{x})^{2}}{n}$$

$$P(A) = \frac{n(A)}{n(S)} \qquad P(A \ or B) = P(A) + P(B) - P(A \ and B)$$

$$\hat{y} = a + bx \qquad \qquad b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$

Copyright reserved