



KWAZULU-NATAL PROVINCE

EDUCATION
REPUBLIC OF SOUTH AFRICA

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

PHYSICAL SCIENCES

COMMON TEST

APRIL 2021

MARKS : 100

TIME : 2 Hours

This question paper consists of 11 pages and 2 data sheets

INSTRUCTIONS AND INFORMATION

1. Write your centre number and examination number in the appropriate spaces on the ANSWER BOOK.
2. This question paper consists of SEVEN questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two sub-questions, for example between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. You may use appropriate mathematical instruments.
8. You are advised to use the attached DATA SHEETS.
9. Show ALL formulae and substitutions in ALL calculations.
10. Round off your final numerical answers to a minimum of TWO decimal places.
11. Give brief motivations, discussions et cetera where required.
12. Write neatly and legibly.

QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Choose the answer and write only the letter (A-D) next to the question number (1.1-1.6) in the ANSWER BOOK, for example 1.11 D.


1.1 A car of mass 1000 kg pulls a trailer of mass 500 kg as shown.




According to Newton's third law of motion, which ONE of the following statements is TRUE?

- A The car and trailer pull each other with a force that is equal in magnitude but opposite in direction. Therefore, the net force is zero and the trailer cannot move.
- B The force that the car exerts on the trailer is greater than the force that the trailer exerts on the car. Therefore, the trailer moves forward.
- C The action force from the car is quicker than the reaction force from the trailer, so they move forward.
- D The action-reaction forces are equal in magnitude, but the force between the ground and wheels pushes them forward. (2)
- 1.2 When the momentum of an object of constant mass is doubled, then its kinetic energy will be ...
- A halved.
- B doubled.
- C three times greater.
- D four times greater. (2)
- 1.3 Two objects are released from the same height at the same time. One object has TWICE the weight of the other. Neglecting friction, which ONE of the following statements is CORRECT for the motion?
- A The heavier object hits the ground first
- B The lighter object hits the ground first.
- C Both objects hit the ground at the same time
- D On hitting the ground, the heavier object has a greater velocity than the lighter object (2)



1.4 Which ONE of the following compounds is a KETONE?



1.5 P and Q represent two organic compounds in the reactions below:



Which ONE of the following represents compound Q?



1.6 $\text{C}_n\text{H}_{2n}\text{O}_2$ is the general formula for both . . .

A A ketone and an aldehyde.

B An ester and an aldehyde.


C A ketone and a carboxylic acid.

D An ester and a carboxylic acid. (2)


[12]

QUESTION 2

- 2.1 A box of mass of 50 kg, placed on a ROUGH horizontal surface, is pulled by a force of 120 N that acts at an angle of 60° to the horizontal. The box moves at a CONSTANT VELOCITY along the surface.



- 2.1.1 State *Newton's First Law of Motion* in words. (2)
- 2.1.2 Draw a free-body diagram of ALL the forces acting on the box while it moves. (4)
- 2.1.3 The box is travelling at constant velocity. What can be deduced about the forces acting on the box? (1)
- 2.1.4 Calculate the normal force acting on the box. (3)
- 2.1.5 The angle of the applied force is now decreased. How will this affect the magnitude of the normal force?
Choose from INCREASES, DECREASES or REMAINS THE SAME
EXPLAIN the answer. (3)
- 2.2 The Sun and the Earth exert a gravitational force on each other.




- 2.2.1 State Newton's law of Universal Gravitation in words. (2)
- 2.2.2 The mass of the Sun is 332 600 times greater than that of the Earth. The distance between the centres of the Sun and the Earth is $1,496 \times 10^9 \text{ m}$. Calculate the gravitational force that the Sun exerts on the Earth. (4)

[19]

QUESTION 3


A goods train is being assembled in a yard. Car 2, of mass of 92×10^3 kg and moving with a velocity of $1.3 \text{ m}\cdot\text{s}^{-1}$ to the right collides with Car 1 of mass 65×10^3 kg that is moving to the right with a velocity of $0.80 \text{ m}\cdot\text{s}^{-1}$. The two cars are joined together after the collision. Ignore the effects of friction.



- 3.1 State the principle of conservation of momentum in words (2)
- 3.2 Calculate the magnitude of the velocity of the cars after the collision. (4)
- 3.3 Determine, by means of appropriate calculations, whether the collision between the two cars is elastic or inelastic. (5)
- [11]**

QUESTION 4

A hot air balloon is rising vertically at a constant velocity of $15 \text{ m}\cdot\text{s}^{-1}$. When the balloon is at some unknown height above the ground, a stone is released from the balloon. The stone is observed to hit the ground with a velocity of $45 \text{ m}\cdot\text{s}^{-1}$.



- 4.1 Define free fall (2)
- 4.2 Write down the magnitude and direction of the initial velocity of the stone when it is released (2)

4.3 Determine how high above the ground the hot air balloon was when the stone was released from it. (3)

4.4 Determine how long the stone took to hit the ground after being released. (3)

4.5 Sketch the position-time graph for the entire motion of the stone from the moment it was released from the hot air balloon to the moment it hits the ground. Use the ground as a zero reference point.

Indicate the following on your graph:

- Height when the stone is released.
- Time taken to hit the ground.

(4)
[14]

QUESTION 5 (Start on a new page.)

The letters A to G in the table below represent organic compounds.

A			
B			
C			
D C ₄ H ₈ O	E CH ₃ CH(CH ₃)CH(CH ₃)CH ₂ CCCH ₃	F	CH ₃ (CH ₂) ₃ CH ₂ OH

Use the above table to answer the following questions:

5.1 Write down the:

- 5.1.1 IUPAC name of compound **A**. (2)
- 5.1.2 IUPAC name of compound **C**. (2)
- 5.1.3 Name of the homologous series to which compound **B** belongs. (1)

- 5.2 Compound E is a hydrocarbon.
- 5.2.1 Define the term *hydrocarbon*. (2)
- 5.2.2 Is compound E a saturated or unsaturated compound? Give a reason for the answer. (2)
- 5.2.3 Write down the general formula of the homologous series to which compound E belongs. (1)
- 5.3 Write down the IUPAC name of a POSITIONAL ISOMER of compound F. (2)
- 5.4 D is the molecular formula of TWO functional isomers.
- 5.4.1 Define the term *functional isomer*. (2)
- 5.4.2 Draw the STRUCTURAL FORMULA of ONE of the FUNCTIONAL isomers of D. (2)
- 5.4.3 Write down the IUPAC name of the OTHER FUNCTIONAL isomer of D. (Hint: This is NOT the same compound as mentioned in Question 5.4.2) (2)

[18]

QUESTION 6 (Start on a new page.)

The boiling points of TWO organic compounds, A and B, were determined.

	FORMULA	MOLECULAR MASS (g.mol ⁻¹)
A	CH ₃ (CH ₂) ₂ COOH	88,1
B	CH ₃ (CH ₂) ₃ CH ₂ OH	88,1

6.1 State the definition of *boiling point*. (2)

The following boiling points were obtained:

137 °C	163 °C
--------	--------

6.2 Write down the boiling point that is most likely to be that of compound A. (2)

6.3 Explain FULLY how you arrived at the answer. (4)

6.4 Write down the IUPAC name of the compound with the LOWER vapour pressure. (2)


6.5 The boiling point of another compound, CH₃(CH₂)₂CH₂OH was also determined. How will the boiling point of this compound compare to that of compound B? Choose from GREATER THAN, LESS THAN or EQUAL TO. Explain the answer. (3)

[13]

QUESTION 7 (Start on a new page.)

The flow diagram shows how COMPOUND X can be used to prepare other organic compounds. The letters A, B, C and D represent different organic reactions.

Compound X is a HALOALKANE.



Use the information in the flow diagram to answer the following questions:

- 7.1 Write down the type of:
- 7.1.1 Substitution reaction represented by A. (1)
- 7.1.2 Elimination reaction represented by C. (1)
- 7.2 Write down the type of reaction represented by:
- 7.2.1 B. (1)
- 7.2.2 D. (1)
- 7.3 Consider REACTION C. Write down
- 7.3.1 TWO reaction conditions for this reaction. (2)
- 7.3.2 IUPAC name of the alkene formed. (1)
- 7.4 Write down the name or formula of the inorganic reactant for REACTION A. (1)
- 7.5 Reaction B involves the reaction of an organic compound with a PRIMARY ALCOHOL.
- 7.5.1 Define the term *primary alcohol*. (2)
- 7.5.2 Write down the name or formula of the catalyst used in REACTION B. (1)
- 7.5.3 Write down the structural formula of the ORGANIC COMPOUND that reacts with the primary alcohol in REACTION B. (2)

[13]**TOTAL MARKS: [100]**

TABLE 3: THE PERIODIC TABLE OF ELEMENTS
TABEL 3: DIE PERIODIEKE TABEL VAN ELE

TABLE 3: THE PERIODIC TABLE OF ELEMENTS

1 (I)	2 (II)	3	4	5	6	7	8	9	10	11	12	13 (III)	14 (IV)	15 (V)	16 (VI)	17 (VII)	18 (VIII)														
1 H 1	2 He 4	3 Li 7	4 Be 9	5 B 11	6 C 12	7 N 14	8 O 16	9 F 19	10 Ne 20	11 Na 23	12 Mg 24	13 Al 27	14 Si 28	15 P 31	16 S 32	17 Cl 35,5	18 Ar 40														
19 K 39	20 Ca 40	21 Sc 45	22 Ti 48	23 V 51	24 Cr 52	25 Mn 55	26 Fe 56	27 Co 59	28 Ni 59	29 Cu 63,5	30 Zn 65	31 Ga 70	32 Ge 73	33 As 75	34 Se 79	35 Br 80	36 Kr 84														
37 Rb 86	38 Sr 88	39 Y 89	40 Zr 91	41 Nb 92	42 Mo 96	43 Tc 98	44 Ru 101	45 Rh 103	46 Pd 106	47 Ag 108	48 Cd 112	49 In 115	50 Sn 119	51 Sb 122	52 Te 128	53 I 127	54 Xe 131														
55 Cs 133	56 Ba 137	57 La 139	72 Hf 179	73 Ta 181	74 W 184	75 Re 186	76 Os 190	77 Ir 192	78 Pt 195	79 Au 197	80 Hg 201	81 Tl 204	82 Pb 207	83 Bi 209	84 Po	85 At	86 Rn														
87 Fr	88 Ra 226	89 Ac	90 Th 232	91 Pa	92 U 238	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr	104 Rf	105 Db	106 Sg	107 Bh	108 Hs	109 Mt	110 Ds	111 Rg	112 Cn	113 Nh	114 Fl	115 Mc	116 Lv	117 Ts	118 Og

KEY/SLEUTEL

Atomic number
Atoomgetal

Electronegativity
Elektronegatiwiteit

Symbol
Simbool

Approximate relative atomic mass
Benaderde relatiewe atoommassa

29 Cu 63,5

**DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 1 (PHYSICS)
GEGEWENS VIR FISIESTE WETENSAPPE GRAAD 12
VRAESTEL 1 (FISIKA)**

TABLE 1: PHYSICAL CONSTANTS / TABEL 1: FISIESTE KONSTANTES

NAME / NAAM	SYMBOL / SIMBOOL	VALUE / WAARDE
Acceleration due to gravity <i>Swaartekragversnelling</i>	g	9,8 m·s ⁻²
Universal gravitational constant <i>Universele gravitasiekonstante</i>	G	6,67 × 10 ⁻¹¹ N·m ² ·kg ⁻²
Speed of light in a vacuum <i>Spoed van lig in 'n vakuum</i>	c	3,0 × 10 ⁸ m·s ⁻¹
Planck's constant <i>Planck se konstante</i>	h	6,63 × 10 ⁻³⁴ J·s
Coulomb's constant <i>Coulomb se konstante</i>	k	9,0 × 10 ⁹ N·m ² ·C ⁻²
Charge on electron <i>Lading op electron</i>	e ⁻	-1,6 × 10 ⁻¹⁹ C
Electron mass <i>Elektronmassa</i>	m _e	9,11 × 10 ⁻³¹ kg
Mass of Earth <i>Massa van Aarde</i>	M	5,98 × 10 ²⁴ kg
Radius of Earth <i>Radius van Aarde</i>	R _E	6,38 × 10 ⁶ m

**TABLE 2: FORMULAE / TABEL 2: FORMULES
MOTION / BEWEGING**

$v_f = v_i + a \Delta t$	$\Delta x = v_i \Delta t + \frac{1}{2} a \Delta t^2$ or/of $\Delta y = v_i \Delta t + \frac{1}{2} a \Delta t^2$
$v_f^2 = v_i^2 + 2a\Delta x$ or/of $v_f^2 = v_i^2 + 2a\Delta y$	$\Delta x = \left(\frac{v_i + v_f}{2} \right) \Delta t$ or/of $\Delta y = \left(\frac{v_i + v_f}{2} \right) \Delta t$

FORCE / KRAG

$F_{net} = ma$	$p = mv$
$f_{s(max)} = \mu_s N$	$f_k = \mu_k N$
$F_{net} \Delta t = \Delta p$ $\Delta p = mv_f - mv_i$	$w = mg$
$F = \frac{Gm_1 m_2}{r^2}$	$g = \frac{Gm}{r^2}$

WORK, ENERGY AND POWER / ARBEID, ENERGIE EN DRYWING

$W = F \Delta x \cos \theta$	$U = mgh$ or/of $E_p = mgh$
$K = \frac{1}{2} mv^2$ or/of $E_k = \frac{1}{2} mv^2$	$W_{net} = \Delta K$ or/of $W_{net} = \Delta E_k$ $\Delta K = K_f - K_i$ or/of $\Delta E_k = E_{kf} - E_{ki}$
$W_{nc} = \Delta K + \Delta U$ or/of $W_{nc} = \Delta E_k + \Delta E_p$	$P = \frac{W}{\Delta t}$
$P_{av} = F \cdot v_{av}$ / $P_{gem} = F \cdot v_{gem}$	



KWAZULU-NATAL PROVINCE

EDUCATION
REPUBLIC OF SOUTH AFRICA

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

PHYSICAL SCIENCES

COMMON TEST

APRIL 2021

MARKING GUIDELINE

MARKS : 100

TIME : 2 hours

This Marking Guideline consists of 8 pages.


QUESTION 1

- 1.1 D✓✓ (2)
 - 1.2 D✓✓ (2)
 - 1.3 C✓✓ (2)
 - 1.4 A✓✓ (2)
 - 1.5 C✓✓ (2)
 - 1.6 D✓✓ (2)
- [12]**

QUESTION 2

2.1.1 A body will remain in its state of rest or motion at constant velocity unless a non-zero resultant/net force acts on it. ✓✓ (2)

2.1.2



CRITERIA FOR MARKING	
• Mark is awarded for label and arrow.	
• Do not penalise for length of arrows.	
• Deduct 1 mark for any additional force.	
• If force(s) do not make contact with body/dot	
: Max:3/4	
• If arrows missing but labels are there:	
Max:3/4	

Acceptable Labels	
N	Normal/ F_N / F_{Normal}
f_k	Friction/ F_f / $f_{kinetic}$
F_A	F /120 N/ $F_{applied}$
W	F_g / $F_{gravity}$ / $Gravitational\ force$ /490 N

2.1.3 Sum of all forces acting on the box is zero/ No net force acting on the box✓ (4)

2.1.4 $F_{net} = ma$
 $F_{net} = 0$
 $N + F_{Ay} = W$ } ✓ (1)

$N + 120\sin 60^\circ = 50(9,8)$ ✓

$N = 386,08\text{ N}$ ✓ (3)

2.1.5 INCREASES✓, If the angle of the applied force decreases, the vertical component of the applied force will also decrease✓. This will affect the normal force since the sum of the normal force and vertical component of the applied force must be equal to the weight of the object✓. (3)

- 2.2.1 Each body in the universe attracts every other body with a force that is directly proportional to the product of their masses and inversely proportional to the square of the distance between their centres. ✓✓ (2)

OR

Each particle in the universe attracts every other particle with a force that is directly proportional to the product of their masses and inversely proportional to the square of the distance between them. ✓✓


2.2.2 $F = G \frac{M_1 \times M_2}{r^2}$ ✓

$$F = 6,67 \times 10^{-11} \frac{(5,98 \times 10^{24})(332600)(5,98 \times 10^{24})}{(1,496 \times 10^9)^2}$$
 ✓

$$F = 3,55 \times 10^{26} \text{ N}$$
 ✓

(4)
[19]**QUESTION 3**

- 3.1 The total linear momentum of a closed system remains constant (is conserved) ✓✓



OR

The total linear momentum before collision is equal to the total linear momentum after the collision. ✓✓ (2)

3.2 $\Sigma p_f = \Sigma p_i$ } ✓
 $(m_1 + m_2)v = m_1 v_{i1} + m_2 v_{i2}$ }

$$(92 \times 10^3 + 65 \times 10^3)v \checkmark = 65 \times 10^3 (0,8) + 92 \times 10^3 (1,3) \checkmark$$

$$v = 1,09 \text{ m} \cdot \text{s}^{-1} \checkmark$$

∴ the velocity of the cars is $1,09 \text{ m} \cdot \text{s}^{-1}$

(4)

POSITIVE MARKING FROM Q. 3.2

3.3	$\Sigma K_f = \frac{1}{2} (m_1 + m_2) v^2 \checkmark$ $= \frac{1}{2} (65 \times 10^3 + 92 \times 10^3) 1,09^2 \checkmark$ $= 93265,85 \text{ J}$ $\Sigma K_f \neq \Sigma K_i \checkmark$ ∴ Collision is Inelastic ✓	$\Sigma K_i = \frac{1}{2} m_1 v_{i1}^2 + \frac{1}{2} m_2 v_{i2}^2$ $= \frac{1}{2} (65 \times 10^3) (0,8)^2 + \frac{1}{2} (92 \times 10^3) (1,3)^2 \checkmark$ $= 98540 \text{ J}$
-----	---	---


(5)
[11]

QUESTION 4

4.1 Motion upon which the only force acting is the force of gravity. ✓✓ (2)

4.2 $15 \text{ m}\cdot\text{s}^{-1}$ ✓upwards✓ (2)

4.3

Stone Trajectory**OPTION 1: From A to D**

UPWARDS POSITIVE	UPWARDS NEGATIVE
$v_f^2 = v_i^2 + 2a\Delta y$	$v_f^2 = v_i^2 + 2a\Delta y$
$-45^2 = 15^2 + 2(-9,8)\Delta y$	$45^2 = -15^2 + 2(9,8)\Delta y$
$\Delta y = -91,84 \text{ m}$	$\Delta y = 91,84 \text{ m}$
\therefore Height above the ground when the stone is released is $91,84 \text{ m}$ ✓	\therefore Height above the ground when the stone is released is $91,84 \text{ m}$ ✓

OPTION 2: From A to D

UPWARDS POSITIVE	UPWARDS NEGATIVE
$v_f = v_i + a\Delta t$	$v_f = v_i + a\Delta t$
$-45 = 15 + (-9,8) \Delta t$	$45 = -15 + (9,8) \Delta t$
$\Delta t = 6,12 \text{ s}$	$\Delta t = 6,12 \text{ s}$
$\Delta y = v_i\Delta t + \frac{1}{2}a\Delta t^2$	$\Delta y = v_i\Delta t + \frac{1}{2}a\Delta t^2$
$\Delta y = (15)(6,12) + \frac{1}{2}(-9,8)(6,12)^2$ ✓	$\Delta y = (-15)(6,12) + \frac{1}{2}(9,8)(6,12)^2$ ✓
$\Delta y = -91,73 \text{ m}$	$\Delta y = 91,73 \text{ m}$
\therefore Height above the ground when the stone is released is $91,73 \text{ m}$ ✓	\therefore Height above the ground when the stone is released is $91,73 \text{ m}$ ✓

OPTION 3: From A to D

UPWARDS POSITIVE	UPWARDS NEGATIVE
$v_f = v_i + a\Delta t$	$v_f = v_i + a\Delta t$
$-45 = 15 + (-9,8) \Delta t$	$45 = -15 + (9,8) \Delta t$
$\Delta t = 6,12 \text{ s}$	$\Delta t = 6,12 \text{ s}$
$\Delta y = \left(\frac{v_i + v_f}{2}\right)\Delta t$ ✓	$\Delta y = \left(\frac{v_i + v_f}{2}\right)\Delta t$ ✓
$\Delta y = \left\{\left(\frac{15 + (-45)}{2}\right) 6,12\right\}$ ✓	$\Delta y = \left\{\left(\frac{15 + (-45)}{2}\right) 6,12\right\}$ ✓
$\Delta y = -91,80 \text{ m}$	$\Delta y = -91,80 \text{ m}$
\therefore Height above the ground when the stone is released is $91,80 \text{ m}$ ✓	\therefore Height above the ground when the stone is released is $91,80 \text{ m}$ ✓

OPTION 3: From C to D

UPWARDS POSITIVE	UPWARDS NEGATIVE
$v_f = v_i + a\Delta t$ $-45 = -15 + (-9,8) \Delta t$ $\Delta t = 3,06 \text{ s}$ $\Delta y = \left(\frac{v_i + v_f}{2}\right) \Delta t \checkmark$ $\Delta y = \left\{\left(\frac{-15 + (-45)}{2}\right) 3,06\right\} \checkmark$ $\Delta y = -91,80 \text{ m}$ \therefore Height above the ground when the stone is released is 91,80 m \checkmark $\Delta y = v_i \Delta t + \frac{1}{2} a \Delta t^2 \checkmark$ $\Delta y = (-15)(3,06) + \frac{1}{2}(-9,8)(3,06)^2 \checkmark$ $\Delta y = -91,78 \text{ m}$ \therefore Height above the ground when the stone is released is 91,78 m \checkmark	$v_f = v_i + a\Delta t$ $45 = 15 + (9,8) \Delta t$ $\Delta t = 3,06 \text{ s}$ $\Delta y = \left(\frac{v_i + v_f}{2}\right) \Delta t \checkmark$ $\Delta y = \left\{\left(\frac{15 + (45)}{2}\right) 3,06\right\} \checkmark$ $\Delta y = 91,80 \text{ m}$ \therefore Height above the ground when the stone is released is 91,80 m \checkmark $\Delta y = v_i \Delta t + \frac{1}{2} a \Delta t^2 \checkmark$ $\Delta y = (15)(3,06) + \frac{1}{2}(9,8)(3,06)^2 \checkmark$ $\Delta y = 91,78 \text{ m}$ \therefore Height above the ground when the stone is released is 91,78 m \checkmark

(3)

4.4

OPTION 1: From A to D

UPWARDS POSITIVE	UPWARDS NEGATIVE
$v_f = v_i + a\Delta t \checkmark$ $-45 = 15 + (-9,8) \Delta t \checkmark$ $\Delta t = 6,12 \text{ s} \checkmark$	$v_f = v_i + a\Delta t \checkmark$ $45 = -15 + (9,8) \Delta t \checkmark$ $\Delta t = 6,12 \text{ s} \checkmark$

OPTION 2: From A to B and then C - D

UPWARDS POSITIVE	UPWARDS NEGATIVE
$v_f = v_i + a\Delta t$ $0 = 15 + (-9,8) \Delta t$ $\Delta t = 1,53 \text{ s}$ $v_f = v_i + a\Delta t$ $-45 = -15 + (-9,8) \Delta t$ $\Delta t = 3,06 \text{ s}$ $\Delta t = \underline{2(1,53) + 3,06} \checkmark$ $\Delta t = 6,12 \text{ s} \checkmark$	$v_f = v_i + a\Delta t$ $0 = -15 + (9,8) \Delta t$ $\Delta t = 1,53 \text{ s}$ $v_f = v_i + a\Delta t$ $45 = 15 + (9,8) \Delta t$ $\Delta t = 3,06 \text{ s}$ $\Delta t = \underline{2(1,53) + 3,06} \checkmark$ $\Delta t = 6,12 \text{ s} \checkmark$


OPTION 3: POSITIVE MARKING FROM 7.2

From A to D


UPWARDS POSITIVE	UPWARDS NEGATIVE
$\Delta y = v_i \Delta t + \frac{1}{2} a \Delta t^2 \checkmark$ $-91,73 = (15\Delta t + \frac{1}{2}(-9,8)\Delta t^2) \checkmark$ $\Delta t = 6,12 \text{ s} \checkmark$	$\Delta y = v_i \Delta t + \frac{1}{2} a \Delta t^2 \checkmark$ $91,73 = (-15\Delta t + \frac{1}{2}(9,8)\Delta t^2) \checkmark$ $\Delta t = 6,12 \text{ s} \checkmark$

(3)

4.5 POSITIVE MARKING FROM 7.2 AND 7.3



OR



NOTE: LEARNERS MAY CHOOSE HEIGHT WHEN THE BALL IS RELEASED AS THE REFERENCE POINT

Criteria for marking	
Correct shape	✓
Indication of height (9,73 – 9,84 m)	✓
Indication of the correct end time	✓
Correct choice of the zero point of reference	✓

(4)
[14]

QUESTION 55.1.1 2,4-dimethylhexane ✓ (2)5.1.2 1-chloro-2-methylpropane ✓ (2)

5.1.3 Esters ✓ (1)

5.2.1 Compound that consists of carbon and hydrogen only. ✓✓ (2)



5.2.2 Unsaturated ✓. Contains a multiple bond/triple bond between carbon atoms in the carbon chain. ✓ (2)

5.2.3 C_nH_{2n-2} ✓ (1)


5.3 pentan-2-ol ✓✓ OR pentan-3-ol ✓✓ (2)

5.4.1 Compounds with same molecular formula but different functional groups. ✓✓ (2)

5.4.2


 OR Books


OR



- | | |
|----------------------------------|-----|
| • Whole structure correct: | 2/2 |
| • Only functional group correct | 1/2 |
| • More than one functional group | 0/2 |

(2)

5.4.3 **POSITIVE MARKING FROM Q5.4.2**

butan-2-one ✓✓ OR butanal ✓✓ OR 2 - Methylpropanal (2)

[18]

QUESTION 6

6.1 The temperature at which the vapour pressure of a substance equals the atmospheric pressure. ✓✓ (2)

6.2 163 °C ✓✓ (2)

6.3 ✓

The acid (A) has TWO sites for hydrogen bonding ✓ while the alcohol (B) has only one site. ✓

The carboxylic acid molecules require more energy to overcome the intermolecular forces ✓

Carboxylic acid will have a higher boiling point. ✓ (4)

6.4 Butanoic acid. ✓✓ (2)

6.5 LOWER ✓.



CH₃(CH₂)₂CH₂OH or butan-1-ol has a shorter carbon chain/smaller surface area than compound B ✓. Strength of the intermolecular forces in CH₃(CH₂)₂CH₂OH or butan-1-ol is weaker than that in compound B. ✓

Therefore lesser energy needed to separate the molecules. ✓ **OR**

Compound B has a longer carbon chain/ larger surface area than CH₃(CH₂)₂CH₂OH or butan-1-ol ✓ The intermolecular forces between molecules of compound B are therefore stronger ✓ and require more energy to separate the molecules. ✓ (3)

[13]

QUESTION 7

7.1.1 Hydrolysis ✓ (1)

7.1.2 Dehydrohalogenation ✓ (1)

7.2.1 Esterification ✓ (1)

7.2.2 Addition/hydration ✓ (1)

7.3.1 Heat ✓

Concentrated strong base in ethanol ✓ (2)


7.3.2 Propene/Prop-1-ene ✓ (1)

7.4 Water/H₂O ✓ (1)

7.5.1 The carbon to which the hydroxyl (OH) group is bonded to, is bonded to ONE other carbon atom. ✓✓ (2)

7.5.2 sulphuric acid/H₂SO₄ ✓ (1)

7.5.3



- | | |
|----------------------------------|-----|
| • Whole structure correct: | 2/2 |
| • Only functional group correct | 1/2 |
| • More than one functional group | 0/2 |

(2)

[13]