



# basic education

Department:  
Basic Education  
**REPUBLIC OF SOUTH AFRICA**

**NATIONAL  
SENIOR CERTIFICATE/  
NASIONALE  
SENIOR SERTIFIKAAT**

**GRADE/GRAAD 12**

**MATHEMATICS P2/WISKUNDE V2  
NOVEMBER 2018  
MARKING GUIDELINES/NASIENRIGLYNE**

**MARKS/PUNTE: 150**

**These marking guidelines consist of 24pages.  
Hierdie nasienriglyne bestaan uit 24 bladsye.**

**NOTE:**

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- If a candidate has crossed out an attempt of a question and not redone the question, mark the crossed out version.
- Consistent accuracy applies in ALL aspects of the marking memorandum. Stop marking at the second calculation error.
- Assuming answers/values in order to solve a problem is NOT acceptable.

**NOTA:**

- *As 'n kandidaat 'n vraag TWEE KEER beantwoord, sien slegs die EERSTE poging na.*
- *As 'n kandidaat 'n antwoord van 'n vraag doodtrek en nie oordoen nie, sien die doodgetrekte poging na.*
- *Volgehoue akkuraatheid word in ALLE aspekte van die nasienriglyne toegepas. Hou op nasien by die tweede berekeningsfout.*
- *Om antwoorde/waardes te aanvaar om 'n probleem op te los, word NIE toegelaat NIE.*

<b>GEOMETRY • MEETKUNDE</b>	
<b>S</b>	<b>A mark for a correct statement (A statement mark is independent of a reason)</b>
	<i>'n Punt vir 'n korrekte bewering ( 'n Punt vir 'n bewering is onafhanklik van die rede)</i>
<b>R</b>	<b>A mark for the correct reason (A reason mark may only be awarded if the statement is correct)</b>
	<i>'n Punt vir 'n korrekte rede ( 'n Punt word slegs vir die rede toegeken as die bewering korrek is)</i>
<b>S/R</b>	<b>Award a mark if statement AND reason are both correct</b>
	<i>Ken 'n punt toe as die bewering EN rede beide korrek is</i>


**QUESTION/VRAAG 1**


1.1.1	140 items	✓ answer	(1)
1.1.2	Modal class/modale klas: $20 < x \leq 30$ minutes <b>OR/OF</b> $20 \leq x < 30$ minutes	✓ answer ✓ answer	(1) (1)
1.1.3	Number of minutes taken = 20 minutes	✓ answer	(1)
1.1.4	140 – 126 [Accept: 124 to 128] 14 orders (12 to 16) <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;">Answer only: Full marks</div>	✓ 126 ✓ answer	(2)
1.1.5	75 <sup>th</sup> percentile is at 105 items =37 minutes [accept 36 – 38 minutes] <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;">Answer only: Full marks</div>	✓ 105 ✓ answer	(2)
1.1.6	Lower quartile is at 35 items =21,5 min [accept 21 – 23 min] IQR = 37 – 21,5 = 15,5 min [accept 13 – 17 min] <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;">Answer only: Full marks</div>	✓ lower quartile (Q <sub>1</sub> ) ✓ answer	(2)

35	70	75	80	80
90	100	100	105	105
110	110	115	120	125

1.2.1(a)	$\bar{x} = \frac{1420}{15}$ = R94,666... = R94,67 <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;">Answer only: Full marks</div>	✓ 1420 ✓ answer	(2)
1.2.1(b)	$\sigma = R22,691... = R22,69$	✓✓ answer	(2)
1.2.2(a)	They both collected the <b>same (equal) amount</b> in tips, i.e. R1 420 over the 15-day period.  <i>Hulle albei het dieselfde bedrag met footjies ontvang, nl. R1 420 oor die 15 dae-tydperk</i>	✓ answer	(1)
1.2.2(b)	Mary's standard deviation is smaller than Reggie's which suggests that there was <b>greater variation in the amount of tips that Reggie collected</b> each day compared to the number of tips that Mary collected each day.  <i>Marie se standaardafwyking is kleiner as Reggie s'n wat beteken dat daar groter variasie/verspreiding in die footjies was wat Reggie elke dag ontvang het in vergelyking met die getal footjies wat Marie elke dag ontvang het.</i>	✓ explanation	(1)
<b>[15]</b>			

**QUESTION/VRAAG 2**





2.1	251 km/h		✓ answer	(1)
2.2.1	$r = 0,52$ OR C		✓ answer	(1)
2.2.2	The points are <b>fairly scattered</b> and the least squares regression line is increasing.  <i>Die punte is redelik verspreid en die kleinstekwadrate-regressielyn neem toe.</i>		✓ reason	(1)
2.3	There is a weak positive relation hence the height could have an influence  <i>Daar is 'n swak positiewe verband, tog kan die lengte 'n invloed hê.</i>		✓ answer	(1)
	<b>OR/OF</b> There is no conclusive evidence that the height of a player will influence his/her tennis serve speed.  <i>Daar is geen duidelike bewys dat die lengte van die speler sy/haar afslaanspoed kan beïnvloed nie.</i>		✓ answer	(1)
	<b>OR/OF</b> There is no conclusive evidence that a taller person will serve faster than a shorter person.  <i>Daar is geen duidelike bewys dat 'n langer speler vinniger sal afslaan as 'n korter een nie.</i>		✓ answer	(1)

2.4	<p>For <math>(0 ; 27,07)</math>, it means that the player has a height of 0 m but can serve at a speed of 27,07 km/h.  <b>It is impossible for a person to have a height of 0 m.</b></p> <p><i>(0 ; 27,07) beteken dat 'n speler 'n lengte van 0 m kan hê en teen 'n spoed van 27,07 km/h kan afslaan. Dit is onmoontlik om 'n lengte van 0 m te hê.</i></p> <p><b>OR/OF</b></p> <p>This means that the <b>player does not exist and therefore cannot serve and have a serve speed.</b></p> <p><i>Dit beteken dat die speler nie bestaan nie en daarom nie kan afslaan en 'n afslaanspoed hê nie.</i></p>	<p>✓ explanation (1)</p> <p>✓ explanation (1)</p>
<b>[5]</b>		



**QUESTION/VRAAG 3**



<p>3.1.1</p>	$m_{KN} = \frac{y_2 - y_1}{x_2 - x_1}$ $m_{KN} = \frac{2 - (-1)}{-1 - 1}$ $= -\frac{3}{2}$ <div style="text-align: center;">  <p>Answer only: Full marks</p> </div>	<p>✓ correct substitution</p> <p>✓ answer</p> <p style="text-align: right;">(2)</p>
<p>3.1.2</p>	$\tan \theta = m_{KN} = -\frac{3}{2}$ $\theta = 180^\circ - 56,31^\circ$ $\theta = 123,69^\circ$ <div style="text-align: center;"> <p>Answer only: Full marks</p> </div>	<p>✓ <math>\tan \theta = m_{KN} = -\frac{3}{2}</math></p> <p>✓ answer</p> <p style="text-align: right;">(2)</p>
<p>3.2</p>	<p>Inclination <math>KL = 123,69^\circ - 78,69^\circ = 45^\circ</math> [ext <math>\angle \Delta</math>]</p> $\tan 45^\circ = m_{KL} = 1$	<p>✓ S</p> <p>✓ <math>\tan 45^\circ = m_{KL} = 1</math></p> <p style="text-align: right;">(2)</p>
<p>3.3</p>	$y = x + c$ $2 = -1 + c$ $c = 3$ $y = x + 3$ <p><b>OR/OF</b></p> $y - y_1 = 1(x - x_1)$ $y - 2 = 1(x - (-1))$ $y = x + 3$	<p>✓ substitute <math>(-1; 2)</math> and <math>m</math></p> <p>✓ equation</p> <p style="text-align: right;">(2)</p> <p>✓ substitute <math>(-1; 2)</math> and <math>m</math></p> <p>✓ equation</p> <p style="text-align: right;">(2)</p>

3.4	$KN = \sqrt{(1+1)^2 + (-1-2)^2}$ $KN = \sqrt{13} \text{ or } 3,61$ <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 100px;">Answer only: Full marks</div>	✓ substitute K and N into distance formula ✓ answer (2)
3.5.1	$(x+3)^2 + (y+5)^2 = 13 \quad \dots(1)$ L is a point on KL $y = x + 3 \quad \dots(2)$ (2) in (1): $(x+3)^2 + (x+3+5)^2 = 13$ $x^2 + 6x + 9 + x^2 + 16x + 64 = 13$ $2x^2 + 22x + 60 = 0$ $x^2 + 11x + 30 = 0$ $(x+5)(x+6) = 0$ $x = -5 \text{ or } x = -6$ $y = -2 \text{ or } y = -3$ $L(-5 ; -2) \text{ or } (-6 ; -3)$ <p><b>OR/OF</b></p> $(x+3)^2 + (y+5)^2 = 13 \quad \dots(1)$ L is a point on KL $y = x + 3 \quad \therefore x = y - 3 \quad \dots(2)$ (2) in (1): $(y-3+3)^2 + (y+5)^2 = 13$ $y^2 + y^2 + 10y + 25 = 13$ $2y^2 + 10y + 12 = 0$ $y^2 + 5y + 6 = 0$ $(y+2)(y+3) = 0$ $y = -2 \text{ or } y = -3$ $x = -5 \text{ or } x = -6$ $L(-5 ; -2) \text{ or } (-6 ; -3)$	✓ equation (1)  ✓ substituting eq (2)  ✓ standard form  ✓ x-values ✓ y-values (5)  ✓ equation (1)  ✓ substituting eq (2)  ✓ standard form  ✓ y-values (both) ✓ x-values (both) (5)
3.5.2	Midpoint of KM: $(-2 ; -1,5)$ $\therefore \frac{x_L + 1}{2} = -2 \text{ and } \frac{y_L - 1}{2} = -\frac{3}{2}$ $\therefore L(-5 ; -2)$ <p><b>OR/OF</b></p> $m_{KN} = m_{LM}$ $\frac{y - (-5)}{x - (-3)} = -\frac{3}{2}$ $2(x+3+5) = -3(x+3)$ $2x+16 = -3x-9$ $5x = -25$ $x = -5$ $\therefore L(-5 ; -2)$ <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 100px;">Answer only: Full marks</div>	✓ midpoint of KM  ✓ x value ✓ y value (3)  ✓ $m_{LM} = m_{KN}$  ✓ x value ✓ y value (3)


	<p><b>OR/OF</b>  <math>N \rightarrow M:</math>  <math>(x; y) \rightarrow (x - 4; y - 4)</math>  <math>\therefore L(-1 - 4; 2 - 4)</math>  <math>\therefore L(-5; -2)</math></p> <p><math>N \rightarrow K:</math>  <math>(x; y) \rightarrow (x - 2; y + 3)</math>  <math>\therefore L(-3 - 2; -5 + 3)</math>  <math>\therefore L(-5; -2)</math></p>	
	<p><b>OR/OF</b>  <math>N \rightarrow M:</math>  <math>(x; y) \rightarrow (x - 4; y - 4)</math>  <math>\therefore L(-1 - 4; 2 - 4)</math>  <math>\therefore L(-5; -2)</math></p> <p><math>N \rightarrow K:</math>  <math>(x; y) \rightarrow (x - 2; y + 3)</math>  <math>\therefore L(-3 - 2; -5 + 3)</math>  <math>\therefore L(-5; -2)</math></p>	
	<p><b>OR/OF</b>  <math>N \rightarrow M:</math>  <math>(x; y) \rightarrow (x - 4; y - 4)</math>  <math>\therefore L(-1 - 4; 2 - 4)</math>  <math>\therefore L(-5; -2)</math></p> <p><math>N \rightarrow K:</math>  <math>(x; y) \rightarrow (x - 2; y + 3)</math>  <math>\therefore L(-3 - 2; -5 + 3)</math>  <math>\therefore L(-5; -2)</math></p>	
	<p><b>OR/OF</b>  <math>N \rightarrow M:</math>  <math>(x; y) \rightarrow (x - 4; y - 4)</math>  <math>\therefore L(-1 - 4; 2 - 4)</math>  <math>\therefore L(-5; -2)</math></p> <p><math>N \rightarrow K:</math>  <math>(x; y) \rightarrow (x - 2; y + 3)</math>  <math>\therefore L(-3 - 2; -5 + 3)</math>  <math>\therefore L(-5; -2)</math></p>	
	<p><b>OR/OF</b>  <math>N \rightarrow M:</math>  <math>(x; y) \rightarrow (x - 4; y - 4)</math>  <math>\therefore L(-1 - 4; 2 - 4)</math>  <math>\therefore L(-5; -2)</math></p> <p><math>N \rightarrow K:</math>  <math>(x; y) \rightarrow (x - 2; y + 3)</math>  <math>\therefore L(-3 - 2; -5 + 3)</math>  <math>\therefore L(-5; -2)</math></p>	<p>✓ transformation</p> <p>✓ x value ✓ y value</p> <p>(3)</p>
3.6	<p>T(-6; -3) (from Question 3.5.1)</p> $KT = \sqrt{(-1 - (-6))^2 + (2 - (-3))^2}$ $= \sqrt{50}$ <p>KN = <math>\sqrt{13}</math> (CA from 3.4)</p> $\text{Area of } \Delta KTN = \frac{1}{2} KT \cdot KN \sin \hat{LKN}$ $= \frac{1}{2} \sqrt{50} \cdot \sqrt{13} \sin 78,69^\circ$ $= 12,50 \text{ square units}$	<p>✓ coordinates of T</p> <p>✓ length of KT</p> <p>✓ substitution into area rule</p> <p>✓ answer</p> <p>(4)</p>



	<p><b>OR/OF</b></p> <p>In <math>\Delta KLM</math>:</p> $\frac{TL}{\sin 22,62^\circ} = \frac{\sqrt{13}}{\sin 78,69^\circ}$ $TL = 1,414..$ $KL = \sqrt{(-1 - (-5))^2 + (2 - (-2))^2}$ $= \sqrt{32}$ $\therefore KT = 7,0708...$ $\text{Area of } \Delta KTN = \frac{1}{2} KT \cdot KN \sin \hat{LKN}$ $= \frac{1}{2} (7,0708) \cdot \sqrt{13} \sin 78,69^\circ$ $= 12,50 \text{ square units}$	<p>✓ length of TL</p> <p>✓ length of KT</p> <p>✓ substitution into area rule</p> <p>✓ answer</p> <p style="text-align: right;">(4)</p>
<b>[22]</b>		




**QUESTION/VRAAG 4**



4.1	F(3;1)	✓ x value ✓ y value (2)
4.2	$FS = \sqrt{(6-3)^2 + (5-1)^2}$ $FS = 5$	✓ substitution of F & S ✓ answer (2)
4.3	$FH(FS) : HG = 1 : 2$ $\therefore HG = 2 FH$ $= 10$	✓ $HG = 10$ (1)
4.4	Tangents from common/same point / <i>Raaklyne vanaf gemeenskaplike of dieselfde punt</i>	✓ answer (1)
4.5.1	$\hat{F}HJ = 90^\circ$ [tan $\perp$ radius / rkl $\perp$ radius] $FJ^2 = 20^2 + 5^2$ [Pyth theorem/stelling] $FJ = \sqrt{425}$ or $5\sqrt{17}$ or 20,62	✓ S ✓ R ✓ S ✓ answer (4)
4.5.2	$(x - m)^2 + (y - n)^2 = 100$	✓ answer (1)

<p>4.5.3</p>	<p>K(22; n)                  GK = HG = 10                  FH = FS = 5  <math>m = 22 - 10</math>  <math>m = 12</math>                  F, H and G are collinear  <i>F, H en G is saamlynig</i>  <math>FG^2 = (12 - 3)^2 + (n - 1)^2</math>  <math>15^2 = 81 + (n - 1)^2</math>  <math>(n - 1)^2 = 144</math>  <math>n - 1 = \pm 12</math>  <math>n \neq 13</math> or <math>n = -11</math>  <math>\therefore G(12; -11)</math></p> <p><b>OR/OF</b></p> <p>K(22; n)                  GK = HG = 10                  FH = FS = 5  <math>m = 22 - 10</math>  <math>m = 12</math>                  Let J(22 ; y):  <math>FJ^2 = (22 - 3)^2 + (y - 1)^2</math>  <math>425 = 361 + y^2 - 2y + 1</math>  <math>0 = y^2 - 2y - 63</math>  <math>0 = (y - 9)(y + 7)</math>  <math>\therefore y = 9</math> or/of <math>y \neq -7</math>  <math>\therefore n = 9 - 20 = -11</math>  <math>\therefore G(12; -11)</math></p>	<p>[radius <math>\perp</math> tangent]                  [radii]                  [radii]</p> <p>[HJ is a common tangent]  <i>[HJ is 'n gemeenskaplike raaklyn]</i></p> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <math>n^2 - 2n - 143 = 0</math>  <math>(n + 11)(n - 13) = 0</math>  <math>n = -11</math> or <math>n \neq 13</math> </div> <p>[radius <math>\perp</math> tangent]                  [radii]                  [radii]</p>	<p>✓ K(22; n)</p> <p>✓ value of <math>m</math></p> <p>✓ subst. of F and G in distance formula                  ✓ <math>FG = 15</math></p> <p>✓ simplification/                  standard form                  ✓ value of <math>n</math>                  ✓ coordinates of G                  (7)</p> <p>✓ K(22; n)</p> <p>✓ value of <math>m</math></p> <p>✓ subst. of F and J in distance formula                  ✓ <math>FJ = \sqrt{425}</math>                  ✓ standard form</p> <p>✓ value of <math>n</math>                  ✓ coordinates of G                  (7)</p>
<b>[18]</b>			


**QUESTION/VRAAG 5**



5.1.1	$k^2 = (\sqrt{5})^2 - 1^2$ $= 4$ $k = -2$ <div style="border: 1px solid black; padding: 5px; display: inline-block; margin-left: 100px;">Answer only: full marks</div>	✓ substitution into theorem of Pythagoras ✓ answer (2)
5.1.2(a)	$\tan \theta = -\frac{1}{2}$	✓ answer (1)
5.1.2(b)	$\cos(180^\circ + \theta) = -\cos \theta$ $= \frac{2}{\sqrt{5}}$ <div style="border: 1px solid black; padding: 5px; display: inline-block; margin-left: 100px;">Answer only: full marks</div>	✓ reduction ✓ answer (2)
5.1.2(c)	$\sin(\theta + 60^\circ) = \frac{a+b}{\sqrt{20}}$ $\text{LHS} = \sin \theta \cos 60^\circ + \cos \theta \sin 60^\circ$ $= \left(\frac{1}{\sqrt{5}}\right)\left(\frac{1}{2}\right) + \left(-\frac{2}{\sqrt{5}}\right)\left(\frac{\sqrt{3}}{2}\right)$ $= \frac{1-2\sqrt{3}}{2\sqrt{5}}$ $= \frac{1-2\sqrt{3}}{\sqrt{20}}$	✓ expansion ✓ subst of $\sin \theta$ ✓ subst of $\cos \theta$ ✓ both special $\angle$ s ✓ $\frac{1-2\sqrt{3}}{2\sqrt{5}}$ (5)
5.1.3	$\tan \theta = -\frac{1}{2}$ $\therefore \theta = 180^\circ - 26,57^\circ$ $\therefore \theta = 153,43^\circ$ $\tan(2\theta - 40^\circ) = \tan[(2 \times 153,43^\circ) - 40^\circ]$ $= \tan 266,87^\circ$ $= 18,3$	✓ $\theta$ ✓ substitution ✓ answer (3)


5.2	$\text{LHS} = \frac{\cos x + \sin x}{\cos x - \sin x} - \frac{\cos x - \sin x}{\cos x + \sin x} \qquad \text{RHS} = 2 \tan 2x$ $= \frac{(\cos x + \sin x)^2 - (\cos x - \sin x)^2}{(\cos x - \sin x)(\cos x + \sin x)}$ $= \frac{\cos^2 x + 2 \sin x \cos x + \sin^2 x - \cos^2 x + 2 \sin x \cos x - \sin^2 x}{\cos^2 x - \sin^2 x}$ $= \frac{2(2 \sin x \cos x)}{\cos^2 x - \sin^2 x}$ $= \frac{2 \sin 2x}{\cos 2x}$ $= 2 \tan 2x$ $= \text{RHS}$ <p><b>OR/OF</b></p> $\text{LHS} = \frac{\cos x + \sin x}{\cos x - \sin x} - \frac{\cos x - \sin x}{\cos x + \sin x} \qquad \text{RHS} = 2 \tan 2x$ $= \frac{(\cos x + \sin x)^2 - (\cos x - \sin x)^2}{(\cos x - \sin x)(\cos x + \sin x)}$ $= \frac{(\cos x + \sin x + \cos x - \sin x)(\cos x + \sin x - \cos x + \sin x)}{\cos^2 x - \sin^2 x}$ $= \frac{(2 \cos x)(2 \sin x)}{\cos^2 x - \sin^2 x}$ $= \frac{2(2 \sin x \cos x)}{\cos^2 x - \sin^2 x}$ $= \frac{2 \sin 2x}{\cos 2x}$ $= 2 \tan 2x$ $= \text{RHS}$ <p><b>OR/OF</b></p> $\text{RHS} = 2 \tan 2x$ $= \frac{2 \sin 2x}{\cos 2x}$ $= \frac{2(2 \sin x \cdot \cos x)}{\cos^2 x - \sin^2 x}$ $= \frac{4 \sin x \cdot \cos x}{\cos^2 x - \sin^2 x}$ $= \frac{1 + 2 \sin x \cdot \cos x - (1 - 2 \sin x \cdot \cos x)}{\cos^2 x - \sin^2 x}$ $= \frac{(\cos x + \sin x)^2 - (\cos x - \sin x)^2}{(\cos x + \sin x)(\cos x - \sin x)}$ $= \frac{(\cos x + \sin x)^2}{(\cos x + \sin x)(\cos x - \sin x)} - \frac{(\cos x - \sin x)^2}{(\cos x + \sin x)(\cos x - \sin x)}$ $= \frac{\cos x + \sin x}{\cos x - \sin x} - \frac{\cos x - \sin x}{\cos x + \sin x} = \text{LHS}$	<p>✓ single fraction</p> <p>✓ expansion</p> <p>✓ simplification (both)</p> <p>✓ double ∠ identity</p> <p>✓ double ∠ identity</p> <p style="text-align: right;">(5)</p> <p>✓ single fraction</p> <p>✓ difference of two squares</p> <p>✓ simplification (both)</p> <p>✓ double ∠ identity</p> <p>✓ double ∠ identity</p> <p style="text-align: right;">(5)</p> <p>✓ double ∠ identity</p> <p>✓ double ∠ identity</p> <p>✓ identity &amp; method</p> <p>✓ factorising numerator and denominator</p> <p>✓ writing as 2 terms</p> <p style="text-align: right;">(5)</p>
-----	---	---




<p>5.3</p>	$\sum_{A=38^\circ}^{52^\circ} \cos^2 A$ $= \cos^2 38^\circ + \cos^2 39^\circ + \cos^2 40^\circ + \dots + \cos^2 51^\circ + \cos^2 52^\circ$ $= \sin^2 52^\circ + \sin^2 51^\circ + \sin^2 50^\circ + \dots + \cos^2 51^\circ + \cos^2 52^\circ$ $= 7(1) + \cos^2 45^\circ$ $= 7 + \left(\frac{\sqrt{2}}{2}\right)^2 \quad \text{or} \quad = 7 + \left(\frac{1}{\sqrt{2}}\right)^2$ $= 7\frac{1}{2}$ <p><b>OR/OF</b></p> $\sum_{A=38^\circ}^{52^\circ} \cos^2 A$ $= \cos^2 38^\circ + \cos^2 39^\circ + \cos^2 40^\circ + \dots + \cos^2 51^\circ + \cos^2 52^\circ$ $= (\cos^2 38^\circ + \sin^2 52^\circ) + (\cos^2 39^\circ + \sin^2 51^\circ) \dots + \cos^2 45^\circ$ $= 7(1) + \cos^2 45^\circ$ $= 7 + \left(\frac{\sqrt{2}}{2}\right)^2 \quad \text{or} \quad = 7 + \left(\frac{1}{\sqrt{2}}\right)^2$ $= 7\frac{1}{2}$ 	<ul style="list-style-type: none"> <li>✓ expansion</li> <li>✓ co ratio</li> <li>✓ <math>\cos^2 45^\circ</math></li> <li>✓ <math>7 \times</math> identity</li>   <li>✓ answer</li> </ul> <p style="text-align: right;">(5)</p> <ul style="list-style-type: none"> <li>✓ expansion</li> <li>✓ pairing</li> <li>✓ <math>\cos^2 45^\circ</math></li> <li>✓ <math>7 \times</math> identity</li>   <li>✓ answer</li> </ul> <p style="text-align: right;">(5)</p>
------------	---	---

[23]

**QUESTION/VRAAG 6**

6.1	Period = $120^\circ$	✓ answer (1)
6.2	$2 = -2 \tan \frac{3}{2}x$ $\tan \left( \frac{3}{2}t \right) = -1$ $\frac{3}{2}t = 135^\circ + k.180^\circ \quad \text{OR/OR} \quad \frac{3}{2}t = -45^\circ + k.180^\circ$ $t = 90^\circ + k.120^\circ ; k \in Z \quad \quad \quad t = -30^\circ + k.120^\circ ; k \in Z$ <p><b>OR/OR</b></p> $2 = -2 \tan \frac{3}{2}x$ $\tan \left( \frac{3}{2}t \right) = -1$ $\frac{3}{2}t = 135^\circ + k.360^\circ \quad \text{or/of} \quad \frac{3}{2}t = 315^\circ + k.360^\circ$ $t = 90^\circ + k.240^\circ \quad \text{or/of} \quad t = 210^\circ + k.240^\circ ; k \in Z$	✓ equating  ✓ general solution of $\frac{3}{2}t$ ✓ general solution of $t$ (3)  ✓ equating  ✓ general solution of $\frac{3}{2}t$ ✓ general solution of $t$ (3)
6.3		✓ asymptotes: $x = \pm 60^\circ ; x = 180^\circ$ ✓ x-intercepts $0^\circ ; \pm 120^\circ$ ✓ negative shape  ✓ $(90^\circ ; 2)$ or $(-30^\circ ; 2)$ or $(30^\circ ; -2)$ or $(-90^\circ ; -2)$  (4)
6.4	$x \in (-60^\circ ; -30^\circ] \text{ or } (60^\circ ; 90^\circ]$ <p><b>OR/OR</b></p> $-60^\circ < x \leq -30^\circ \text{ or } 60^\circ < x \leq 90^\circ$	✓ interval ✓ interval ✓ notation (3)  ✓ interval ✓ interval ✓ notation (3)
6.5	$g(x) = -2 \tan \left[ \frac{3}{2}(x + 40^\circ) \right] = f(x + 40^\circ)$ Translation of $40^\circ$ to the left / skuif met $40^\circ$ links	✓ Translation of $40^\circ$ ✓ to the left  (2)
<b>[13]</b>		


**QUESTION/VRAAG 7**



<p>7.1</p>	<p><math>\hat{A}BD = 30^\circ</math>  <math>\sin 30^\circ = \frac{h}{AB}</math>  <math>AB = \frac{h}{\sin 30^\circ}</math> <b>OR</b> <math>AB = \frac{h}{\frac{1}{2}}</math> <b>OR</b> <math>AB = 2h</math></p> <p><b>OR/OF</b></p> <p><math>\hat{B}AD = 60^\circ</math>  <math>\cos 60^\circ = \frac{h}{AB}</math>  <math>AB = \frac{h}{\cos 60^\circ}</math> <b>OR</b> <math>AB = \frac{h}{\frac{1}{2}}</math> <b>OR</b> <math>AB = 2h</math></p>	<p>✓ <math>\hat{A}BD = 30^\circ</math></p> <p>✓ answer (2)</p> <p>✓ <math>\hat{B}AD = 60^\circ</math></p> <p>✓ answer (2)</p>
<p>7.2</p>	<p><math>BC^2 = AB^2 + AC^2 - 2AB \cdot AC \cos \hat{B}AC</math>  <math>= (2h)^2 + (3h)^2 - 2(2h)(3h) \cos 2x</math>  <math>= 13h^2 - 12h^2 (2 \cos^2 x - 1)</math>  <math>= 13h^2 - 24h^2 \cos^2 x + 12h^2</math>  <math>= 25h^2 - 24h^2 \cos^2 x</math>  <math>BC = h\sqrt{25 - 24 \cos^2 x}</math></p>	<p>✓ use of cosine rule in <math>\triangle ABC</math></p> <p>✓ substitution</p> <p>✓ double angle identity</p> <p>✓ <math>25h^2 - 24h^2 \cos^2 x</math></p> <p>(4)</p>
<p><b>[6]</b></p>		




**QUESTION/VRAAG 8**



8.1.1	$\hat{P} = \hat{M}_1 = 66^\circ$	[tan chord theorem/raaklyn koordst]	✓S ✓R (2)
8.1.2	$\hat{M}_2 = 90^\circ$	[∠ in semi circle/∠ in halfsirkel]	✓S ✓R (2)
8.1.3	$\hat{N}_1 = 180^\circ - (90^\circ + 66^\circ)$ $= 24^\circ$	[sum of ∠s of /som van ∠e ΔMNP]	✓S (1)
8.1.4	$\hat{O}_2 = \hat{P} = 66^\circ$	[corres. ∠s;/ooreenk ∠e, PM    OR]	✓S ✓R (2)
8.1.5	$\hat{R} + \hat{N}_1 + \hat{N}_2 = 180^\circ - 66^\circ$ $= 114^\circ$ $\hat{R} = \hat{N}_1 + \hat{N}_2 = 57^\circ$ $\therefore \hat{N}_2 = 33^\circ$	[sum of ∠s of /som van ∠e ΔRNO]  [∠s opposite = radii/∠e teenoor = radii]	✓S ✓S/R ✓S (3)
	<b>OR/OF</b> $\hat{P}\hat{O}\hat{R} = 114^\circ$ $\hat{P}\hat{N}\hat{R} = 57^\circ$ $\therefore \hat{N}_2 = 33^\circ$	[∠s on straight line/∠e op reguitlyn] [∠ at centre = twice ∠ at circumference/ midpts∠ = 2 × omtreks∠]	✓S ✓S/R ✓S (3)


8.2



8.2.1	$FC \parallel AB \parallel GH$ [opp sides of <b>rectangle</b> / <i>teenoorst sye v reghoek</i> ]	✓ R (1)
8.2.2	$\frac{AC}{CH} = \frac{AF}{FG}$ [line $\parallel$ one side of $\Delta$ ] <b>OR</b> [prop theorem; $FC \parallel GH$ ] <i>[lyn <math>\parallel</math> een sy van <math>\Delta</math>] <b>OF</b> [eweredighst; <math>FC \parallel GH</math>]</i>  $\frac{AC}{21} = \frac{20}{15}$ $AC = \frac{20 \times 21}{15}$ $= 28$ $DB = AC = 28$ [diags of rectangle =/ <i>hoeklyne v reghoek</i> = ] $DM = \frac{1}{2}DB = 14$ [diags of rectangle bisect/ <i>hoekl v reghoek halveer</i> ]	✓ S ✓ R   ✓ AC ✓ S ✓ S (5)
<b>[16]</b>		


**QUESTION/VRAAG 9**

9.1



9.1	<p>Constr/Konstr.: Draw KO and MO/Trek KO en MO</p> <p>Proof:</p> $\hat{O}_1 = 2\hat{J} \quad [\angle \text{ at centre} = \text{twice } \angle \text{ at circumference}]$ <p><i>[midpts <math>\angle = 2 \times \text{omtreks } \angle</math>]</i></p> $\hat{O}_2 = 2\hat{L} \quad [\angle \text{ at centre} = \text{twice } \angle \text{ at circumference}]$ $\hat{O}_1 + \hat{O}_2 = 360^\circ \quad [\angle \text{ s around a point / } \angle \text{ e om 'n punt}]$ $\therefore 2\hat{J} + 2\hat{L} = 360^\circ$ $\therefore 2(\hat{J} + \hat{L}) = 360^\circ$ $\therefore \hat{J} + \hat{L} = 180^\circ$ <p><b>OR/OF</b></p> <p>Constr/Konstr.: Draw KO and MO/Trek KO en MO</p> <p>Proof:</p> <p>Let <math>\hat{J} = x</math></p> $\hat{O}_1 = 2x \quad [\angle \text{ at centre} = \text{twice } \angle \text{ at circumference}]$ <p><i>[midpts <math>\angle = 2 \times \text{omtreks } \angle</math>]</i></p> $\hat{O}_2 = 360^\circ - 2x \quad [\angle \text{ s around a point / } \angle \text{ e om 'n punt}]$ $\therefore \hat{L} = 180^\circ - x \quad [\angle \text{ at centre} = \text{twice } \angle \text{ at circumference}]$ $\therefore \hat{J} + \hat{L} = 180^\circ$	<p>✓ construction</p> <p>✓ S/R</p> <p>✓ S</p> <p>✓ S/R</p> <p>✓ S</p> <p style="text-align: right;">(5)</p> <p>✓ construction</p> <p>✓ S ✓ R</p> <p>✓ S/R</p> <p>✓ S</p> <p style="text-align: right;">(5)</p>
-----	---	--

9.2




9.2.1(a)	$\hat{B}_1 = x$ [∠s in same seg/∠e in dieselfde segm]	✓ S ✓ R	(2)
9.2.1(b)	$\hat{B}_2 = y$ [ext ∠ of cyclic quad/buite∠ koordevh]	✓ S ✓ R	(2)
9.2.2	$\hat{C} = 180^\circ - (x + y)$ [sum of ∠s of/som v ∠e, ΔACR] $\hat{SBD} + \hat{C} = x + y + 180^\circ - (x + y)$ $\hat{SBD} + \hat{C} = 180^\circ$ SCDB is a cyclic quad [converse opp angles of cyclic quad] [omgekeerde teenoorst ∠e koordevh]	✓ S ✓ S ✓ R	(3)
	<b>OR/OF</b> $\hat{S}_1 = \hat{T}_2$ [∠s in same segment/∠e in dies. segment] $\hat{T}_2 = \hat{D}_1 + \hat{D}_2 = \hat{BDR}$ [ext ∠ of cyc quad/buite∠ koordevh] $\therefore \hat{S}_1 = \hat{BDR}$ $\therefore$ SCDB is cyc quad [ext ∠ of quad = opp ∠/buite∠ = tos ∠]	✓ S ✓ S ✓ R	(3)

9.2.3	$\hat{T}_4 = y - 30^\circ$ [ext $\angle$ of/buite $\angle$ $\Delta$ TDR] $\hat{T}_1 = y - 30^\circ$ [vert opp $\angle$ s =/regoorst $\angle$ e =] $y - 30^\circ + x + 100^\circ = 180^\circ$ [sum of $\angle$ s of/som v $\angle$ e, $\Delta$ AST] $\therefore x + y = 110^\circ$ $\hat{S}\hat{B}\hat{D} = 110^\circ$ $\therefore$ SD not diameter [line does not subtend $90^\circ \angle$ ] <i>SD nie 'n middellyn [lyn onderspan nie <math>90^\circ \angle</math> ]</i> <b>OR/OF</b> $\hat{A}\hat{S}\hat{T} = \hat{C} + \hat{D}_2$ [ext $\angle$ of/buite $\angle$ $\Delta$ SCD] $\hat{C} = 100^\circ - 30^\circ = 70^\circ$ $\hat{S}\hat{B}\hat{D} = 180^\circ - 70^\circ$ [opp $\angle$ s cyclic quad/ teenoorst $\angle$ e kdvh] $= 110^\circ$ $\therefore$ SD not diameter [line does not subtend $90^\circ \angle$ ] <i>SD nie 'n middellyn [lyn onderspan nie <math>90^\circ \angle</math> ]</i>	✓ S ✓ S  ✓ S ✓ R  ✓ S ✓ S ✓ S ✓ R	(4)          (4)
			<b>[16]</b>



**QUESTION/VRAAG 10**



<p>10.1.1</p>	<p><math>\hat{A}_2 = \hat{A}_1 = 90^\circ - x</math> [= chords subtend = <math>\angle s</math> = <i>kde onderspan</i> = <math>\angle e</math>]</p> <p><math>\hat{D}_2 = x</math> [exterior angle of cyclic quad/<i>buite</i> <math>\angle</math> <i>koordevh.</i>]</p> <p><math>\therefore \hat{C}_2 = 90^\circ - x</math> [sum of <math>\angle s</math> of/<i>som v</i> <math>\angle e</math>, <math>\Delta DCM</math>]</p> <p><math>\therefore \hat{C}_2 = \hat{A}_1 = 90^\circ - x</math></p> <p><math>\therefore MC</math> is a tangent to the circle at <math>C</math> [converse: tan chord th] <i>MC is 'n raaklyn by C [omgekeerde raakl koordst]</i></p> <p><b>OR/OF</b></p> <p><math>\hat{A}_2 = \hat{A}_1 = 90^\circ - x</math> [= chords subtend = <math>\angle s</math>/ = <i>kde onderspan</i> = <math>\angle e</math>]</p> <p><math>\hat{C}_1 + \hat{C}_2 = x</math> [sum of <math>\angle s</math> of/<i>som v</i> <math>\angle e</math>, <math>\Delta ACM</math>]</p> <p><math>\therefore \hat{C}_1 + \hat{C}_2 = \hat{B} = x</math></p> <p><math>\therefore MC</math> is a tangent to the circle at <math>C</math> [converse : tan chord th] <i>MC is 'n raaklyn by C [omgekeerde raakl koordst]</i></p> <p><b>OR/OF</b></p> <p>In <math>\Delta AMC</math> and <math>\Delta ACB</math>:</p> <p><math>\hat{A}_2 = \hat{A}_1 = 90^\circ - x</math> [= chords subtend = <math>\angle s</math>/ = <i>kde onderspan</i> = <math>\angle e</math>]</p> <p><math>\hat{A}MC = \hat{A}CB = 90^\circ</math> [given]</p> <p><math>\therefore \hat{C}_1 + \hat{C}_2 = \hat{B} = x</math></p>	<p>✓ S ✓ R</p> <p>✓ S/R</p> <p>✓ <math>\hat{C}_2 = 90^\circ - x</math></p> <p>✓ R</p> <p>(5)</p> <p>✓ S ✓ R</p> <p>✓✓ <math>\hat{C}_1 + \hat{C}_2 = x</math></p> <p>✓ R</p> <p>(5)</p> <p>✓ S ✓ R</p> <p>✓✓ <math>\hat{C}_1 + \hat{C}_2 = x</math></p>
---------------	---	--

	<p><math>\therefore MC</math> is a tangent to the circle at <math>C</math> [converse : tan chord th]  <i>MC is 'n raaklyn by C [omgekeerde raakl koordst]</i></p>	<p>✓ R (5)</p>
	<p>In <math>\triangle ACB</math> and/en <math>\triangle CMD</math>  <math>\hat{B} = \hat{D}_2 = x</math> [proved <b>OR</b> exterior <math>\angle</math> of cyclic quad.]  <i>[bewys OF buite <math>\angle</math> v koordevh]</i>  <math>\hat{A}_2 = \hat{C}_2 = 90^\circ - x</math> [proved <b>OR</b> sum of <math>\angle</math>s in <math>\triangle</math>]  <i>[Bewys OF som v <math>\angle</math>e in <math>\triangle</math>]</i>  <math>\triangle ACB \parallel \triangle CMD</math> [<math>\angle, \angle, \angle</math>]</p> <p><b>OR/OF</b>          In <math>\triangle ACB</math> and/en <math>\triangle CMD</math>  <math>\hat{B} = \hat{D}_2 = x</math> [proved <b>OR</b> exterior <math>\angle</math> of cyclic quad.]  <i>[bewys OF buite <math>\angle</math> v koordevh]</i>  <math>\hat{A}\hat{C}B = \hat{A}\hat{M}C = 90^\circ</math> [given/gegee]  <math>\triangle ACB \parallel \triangle CMD</math> [<math>\angle, \angle, \angle</math>]</p> <p><b>OR/OF</b>          In <math>\triangle ACB</math> and/en <math>\triangle CMD</math>  <math>\hat{B} = \hat{D}_2 = x</math> [proved <b>OR</b> exterior <math>\angle</math> of cyclic quad]  <i>[bewys OF buite <math>\angle</math> v koordevh]</i>  <math>\hat{A}_2 = \hat{C}_2 = 90^\circ - x</math> [proved <b>OR</b> sum of <math>\angle</math>s in <math>\triangle</math>]  <i>[Bewys OF som v <math>\angle</math>e in <math>\triangle</math>]</i>  <math>\hat{A}\hat{C}B = \hat{A}\hat{M}C = 90^\circ</math> [given <b>OR</b> sum of <math>\angle</math>s in <math>\triangle</math>]  <i>[gegee OF som v <math>\angle</math>e in <math>\triangle</math>]</i>  <math>\triangle ACB \parallel \triangle CMD</math></p>	<p>✓ S          ✓ S          ✓ R (3)          ✓ S          ✓ S          ✓ R (3)          ✓ S          ✓ S          ✓ S          ✓ S (3)</p>
10.2.1	<p><math>\frac{BC}{MD} = \frac{AB}{DC}</math> [<math>\triangle ACB \parallel \triangle CMD</math>]  <math>\frac{DC}{MD} = \frac{AB}{DC}</math> [<math>BC = DC</math>]  <math>\therefore DC^2 = AB \times MD</math></p> <p>In <math>\triangle AMC</math> and/en <math>\triangle CMD</math>  <math>\hat{M}</math> is common/<i>gemeen</i>  <math>\hat{A}_1 = \hat{C}_2</math> [tan chord th /<i>raaklyn koordst</i>]</p> <p><b>OR/OF</b>  <math>\hat{C}_1 + \hat{C}_2 = \hat{B} = \hat{D} = x</math> [tan chord th /<i>raaklyn koordst</i> <b>OR/OF</b>          exterior <math>\angle</math> of cyclic quad/ <i>buite <math>\angle</math> v kdvh</i>]</p> <p><math>\triangle AMC \parallel \triangle CMD</math> [<math>\angle, \angle, \angle</math>]  <math>\frac{AM}{CM} = \frac{CM}{MD}</math>  <math>\therefore CM^2 = AM \times MD</math>  <math>\therefore \frac{CM^2}{DC^2} = \frac{AM \times MD}{AB \times MD}</math>  <math>= \frac{AM}{AB}</math></p>	<p>✓ <math>\frac{BC}{MD} = \frac{AB}{DC}</math>          ✓ <math>DC^2 = AB \times MD</math>          ✓ S          ✓ S          ✓ <math>CM^2 = AM \times MD</math>          ✓ <math>\frac{AM \times MD}{AB \times MD}</math> (6)</p>

	<p><b>OR/OF</b></p> $\frac{AC}{MC} = \frac{AB}{DC} \quad [\Delta ACB \parallel \Delta CMD]$ $\therefore CM \times AB = AC \times DC$ <p>In <math>\Delta AMC</math> and/en <math>\Delta ACB</math>  <math>\hat{C} = \hat{M} = 90^\circ</math> [given]  <math>\hat{A}_1 = \hat{A}_2</math> [proven]</p> <p><b>OR/OF</b></p> $\hat{A}\hat{C}\hat{M} = \hat{B} = x$ [proven] $\Delta AMC \parallel \Delta ACB$ [ $\angle, \angle, \angle$ ] $\frac{AC}{AM} = \frac{BC}{MC}$ $\therefore AC \times MC = AM \times BC$ $\therefore AC = \frac{BC \cdot AM}{MC}$ $CM \times AB = \frac{BC \cdot AM}{MC} \times DC$ $CM^2 = \frac{DC \cdot AM}{AB} \times DC \quad [BC = DC]$ $\frac{CM^2}{DC^2} = \frac{AM}{AB}$	<p>✓ <math>\frac{AC}{MC} = \frac{AB}{DC}</math></p> <p>✓ S</p> <p>✓ S</p> <p>✓ <math>AC \cdot MC = AM \cdot BC</math></p> <p>✓ equating</p> <p>✓ S</p> <p>(6)</p>
<p>10.2.2</p>	<p>In <math>\Delta DMC</math>:</p> $\frac{CM}{DC} = \sin x$ $\frac{CM^2}{DC^2} = \sin^2 x \quad \frac{AC}{AB} = \frac{CM}{DC}$ $\therefore \frac{AM}{AB} = \sin^2 x$ <p><b>OR/OF</b></p> <p>In <math>\Delta ABC</math>:</p> $\sin x = \frac{AC}{AB}$ <p>In <math>\Delta AMC</math>:</p> $\sin x = \frac{AM}{AC}$ $\sin x \cdot \sin x = \frac{AC}{AB} \times \frac{AM}{AC} = \frac{AM}{AB}$	<p>✓ trig ratio</p> <p>✓ square both sides</p> <p>(2)</p> <p>✓ 2 equations for <math>\sin x</math></p> <p>✓ product</p> <p>(2)</p>
<p><b>[16]</b></p>		

**TOTAL/TOTAAL: 150**